

W-TRIVIALITY OF LOW DIMENSIONAL MANIFOLDS

ARITRA C BHATTACHARYA, BIKRAMJIT KUNDU, AND ANIRUDDHA C NAOLEKAR

ABSTRACT. A space X is W -trivial if for every real vector bundle α over X the total Stiefel-Whitney class $w(\alpha)$ is 1. It follows from a result of Milnor that if X is an orientable closed smooth manifold of dimension 1, 2, 4 or 8, then X is not W -trivial. In this note we completely characterize W -trivial orientable connected closed smooth manifolds in dimensions 3, 5 and 6. In dimension 7, we describe necessary conditions for an orientable connected closed smooth 7-manifold to be W -trivial.

1. INTRODUCTION.

A theorem of Milnor [8] (see also [1, Theorem 1, page 223]) states the following.

Theorem 1.1. [8, Theorem 1, page 444] There exists a real vector bundle α over the sphere S^n with $w_n(\alpha) \neq 0$ if and only if $n = 1, 2, 4, 8$. \square

For $n \in \{1, 2, 4, 8\}$, the bundles in Milnor's theorem are the famous Hopf bundles. Since $H^i(S^n; \mathbb{Z}_2) = 0$ for $i \neq 0, n$, for a real vector bundle α over S^n , $w(\alpha) = 1$ if and only if $w_n(\alpha) = 0$.

We recall the following definition from [12].

Definition 1.2. A space X is said to be W -trivial if for every real vector bundle α over X , the total Stiefel-Whitney class $w(\alpha)$ is 1.

In view of the above definition, Milnor's theorem can be restated as: S^n is W -trivial if and only if $n \neq 1, 2, 4, 8$.

In this note we address the question if there are orientable connected closed smooth manifolds other than the spheres that are W -trivial. We completely characterize W -trivial manifolds in dimensions 3, 5, 6 and prove a necessary condition for a manifold to be W -trivial in dimension 7. Before stating the main results we make a few remarks.

Remark 1.3. (1) If $n = 1, 2, 4, 8$ and X is an orientable connected closed smooth n -manifold, then X is not W -trivial. This is because if $f : X \rightarrow S^n$ is a degree 1 map, then

$$w_n(f^* \alpha) \neq 0$$

where α is the Hopf bundle on S^n .

- (2) Recall that, for a commutative ring R , a R -homology n -sphere is an orientable connected closed smooth n -manifold X with $H_i(X; R) \cong H_i(S^n; R)$ for all i . It is known [8, page 94] that for a real vector bundle α the smallest positive integer n for which $w_n(\alpha) \neq 0$ is a power of 2. Thus if X is a \mathbb{Z}_2 -homology n -sphere and n is not a power of 2, then X is W -trivial.
- (3) It would be interesting to know if Theorem 1.1 is true for \mathbb{Z}_2 -homology spheres.

2010 *Mathematics Subject Classification.* 57R20.

Key words and phrases. Stiefel-Whitney class, W -trivial.

We now state our main results. We assume that all manifolds that appear in the sequel are connected.

Theorem 1.4. Let X be an orientable closed smooth manifold of dimension either 3 or 5. Then X is W -trivial if and only if X is a \mathbb{Z}_2 -homology sphere. \square

In dimension 6 there exist W -trivial manifolds X that are not \mathbb{Z}_2 -homology spheres. For example $S^3 \times S^3$ is W -trivial (see Remark 1.3 (2) above). In dimension 6 we prove the following.

Theorem 1.5. Let X be an orientable closed smooth manifold of dimension 6. Then X is W -trivial if and only if

$$H_i(X; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0 \\ F & i = 1 \\ F' & i = 2 \\ \mathbb{Z}^t \oplus F' & i = 3 \\ F & i = 4 \\ 0 & i = 5 \\ \mathbb{Z} & i = 6 \end{cases}$$

where F, F' are finite abelian groups neither of which contain any element of order 2 and $t \geq 0$. \square

For example $S^3 \times S^3$ is a 6-manifold whose homology groups are of the form described in the above theorem and hence is W -trivial. In dimension 7 we derive the following homological restriction on W -trivial manifolds.

Theorem 1.6. Let X be an orientable closed smooth 7-manifold. If X is W -trivial, then the integral homology groups of X are of the form

$$H_i(X; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0 \\ F & i = 1 \\ F' & i = 2 \\ F'' & i = 3 \\ F' & i = 4 \\ F & i = 5 \\ 0 & i = 6 \\ \mathbb{Z} & i = 7 \end{cases}$$

where F, F', F'' are finite abelian groups with F, F'' not having any element of order 2. The converse is true if in addition F' does not contain any element of order 2. \square

In particular, if X is a W -trivial orientable closed smooth 7-manifold, then X is a rational homology 7-sphere. We have not been able to decide whether or not there is a W -trivial oriented closed smooth 7-manifold X such that $H_2(X; \mathbb{Z})$ contains an element of order 2.

We end this section by looking at some examples and non-examples. Note that in dimensions 3, 5 and 7 every W -trivial manifold turns out to be a rational homology sphere. We begin by discussing the W -triviality of certain familiar rational homology spheres.

Example 1.7. Since $H^1(\mathbb{R}P^n; \mathbb{Z}_2) \neq 0$, the real projective spaces $\mathbb{R}P^n$ are not W -trivial (see Proposition 2.2 below). Let

$$L_{n,p} = S^{2n-1} / \mathbb{Z}_p$$

$n > 1$ and p a prime, denote the Lens space obtained as a quotient of S^{2n-1} by the diagonal action of \mathbb{Z}_p . Here we regard \mathbb{Z}_p as the group of p -th roots of unity and S^{2n-1} as the unit sphere in \mathbb{C}^n . The integral homology groups of $L_{n,p}$ are zero in even degrees i , $1 < i < 2n-1$ and are equal to \mathbb{Z}_p in odd degrees i , $1 \leq i < 2n-1$. Hence if p odd, then the Lens space $L_{n,p}$ is W -trivial.

Example 1.8. Let $V_2(\mathbb{R}^5)$ denote the Stiefel manifold of orthonormal 2-frames in \mathbb{R}^5 . $V_2(\mathbb{R}^5)$ is a rational homology 7-sphere. The integral homology of $V_2(\mathbb{R}^5)$ is of the form as in Theorem 1.6 with $F = F' = 0$ and $F'' = \mathbb{Z}_2$ and hence $V_2(\mathbb{R}^5)$ is not W -trivial. Another way to prove this is to consider the sphere bundle

$$S^3 \hookrightarrow V_2(\mathbb{R}^5) \xrightarrow{p} V_1(\mathbb{R}^5) = S^4.$$

A part of the Gysin sequence associated to the above sphere bundle is

$$\rightarrow H^0(S^4; \mathbb{Z}_2) \rightarrow H^4(S^4; \mathbb{Z}_2) \xrightarrow{p^*} H^4(V_2(\mathbb{R}^5); \mathbb{Z}_2) \rightarrow H^1(S^4; \mathbb{Z}_2) \rightarrow .$$

It is well known [3] that $H^4(V_2(\mathbb{R}^5); \mathbb{Z}_2) \cong \mathbb{Z}_2$. Hence p^* is an isomorphism. Thus if α is the Hopf bundle over S^4 , then $w_4(p^*(\alpha)) = p^*(w_4(\alpha)) \neq 0$ and hence $V_2(\mathbb{R}^5)$ is not W -trivial.

Example 1.9. Examples of W -trivial manifolds in dimension 7 are given by certain rational homology spheres constructed by Ruberman [11, Example 7, page 232]. In each dimension $n > 5$ Ruberman has constructed examples X_k of simply connected rational homology n -spheres with

$$H_i(X_k; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0, n \\ \mathbb{Z}_k & i = 2, n-3 \\ 0 & \text{otherwise.} \end{cases}$$

In dimension 7, these rational homology spheres X_k have the homology as in Theorem 1.6 with $F = F'' = 0$, $F' = \mathbb{Z}_k$, and hence are W -trivial if k is odd. For a general $n > 5$, when k is odd, it is clear that X_k is a \mathbb{Z}_2 -homology sphere. If in addition n is not a power of 2, then X_k is W -trivial. In each dimension $n \geq 5$ there exist, by a construction of Kervaire [7, Theorem 1, page 67], a non simply connected integral homology sphere K . If the Ruberman manifold X_k is W -trivial, then the connected sum $X_k \# K$ is W -trivial. This gives us examples of non simply connected W -trivial manifolds.

Example 1.10. In dimension 5, there exist simply connected manifolds B_k (see [2]) whose integral homology is of the form

$$H_i(B_k; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0, 5 \\ \mathbb{Z}_k \oplus \mathbb{Z}_k & i = 2 \\ 0 & \text{otherwise.} \end{cases}$$

If k is odd, then B_k is W -trivial. One may now take connected sum with the integral homology spheres of Kervaire to obtain a non simply connected W -trivial 5-manifold.

This note is organized as follows. In section 2 we collect some preliminary facts about W -trivial manifolds and then prove the main theorems. Throughout, we follow the conventions stated below.

Conventions. In this note we follow the conventions of [10] which we recall briefly. The notations F, F', F'', \dots will denote finite abelian groups. Given F (resp. $F', F'' \dots$), the integer s (resp. s', s'', \dots) will denote the number of primes p_i that are equal to 2 in a direct sum decomposition

$$F = \oplus_i \mathbb{Z}_{p_i}^{k_i}$$

of F (resp. F', F'', \dots) with p_i not necessarily distinct. Thus we have

$$\text{Ext}(F, \mathbb{Z}) = F$$

and

$$\text{Ext}(F, \mathbb{Z}_2) = \mathbb{Z}_2^s = \text{Hom}(F, \mathbb{Z}_2).$$

Given s (resp. s', s'', \dots) the integers u, v (resp. $u', v'; u'', v''; \dots$) will denote the number of primes p_i such that k_i is equal to 1 and the number of primes p_i such that $k_i > 1$ respectively. Thus $s = u + v$. In particular,

$$\text{Ext}(F, \mathbb{Z}_4) = \mathbb{Z}_2^u \oplus \mathbb{Z}_4^v.$$

Finally, all vector bundles that we consider are real unless stated otherwise.

2. PROOFS

In this section we prove the main theorems. We begin by fixing some notations and proving some preliminary results that we shall use. We shall denote by ρ_k the mod- k reduction homomorphism

$$\rho_k : H^i(X; \mathbb{Z}) \longrightarrow H^i(X; \mathbb{Z}_k)$$

and by

$$\rho_{4,2} : H^4(X; \mathbb{Z}_4) \longrightarrow H^4(X; \mathbb{Z}_2)$$

the homomorphism induced by the coefficient surjection.

We shall denote by \mathfrak{P} the Pontryagin square (see [13])

$$\mathfrak{P} : H^{2k}(X; \mathbb{Z}_2) \longrightarrow H^{4k}(X; \mathbb{Z}_4).$$

The Pontryagin square satisfies

$$\rho_{4,2} \circ \mathfrak{P}(x) = x^2$$

for every $x \in H^{2k}(X; \mathbb{Z}_2)$.

We begin with some observations that we shall use in the sequel. Recall that given a finite abelian group $F = \bigoplus_i \mathbb{Z}_{p_i^{k_i}}$, the integer s denotes the number of primes p_i that are equal to 2.

Remark 2.1. Let F be a finite abelian group and

$$\beta : \mathbb{Z}_2^\ell \oplus \mathbb{Z}_2^s \longrightarrow F$$

be a monomorphism. Then $\ell = 0$. This is because the number of elements of order 2 in $\mathbb{Z}_2^\ell \oplus \mathbb{Z}_2^s$ and F are $2^{\ell+s} - 1$ and $2^s - 1$ respectively. Since $2^{\ell+s} - 1 \leq 2^s - 1$, it follows that $\ell = 0$.

Proposition 2.2. Let X be a closed smooth n -manifold. If X is W -trivial, then

- (1) X is orientable,
- (2) X is unoriented nullbordant,
- (3) $H^1(X; \mathbb{Z}_2) = 0$,
- (4) $H_1(X; \mathbb{Z})$ is finite and does not have elements of order 2,
- (5) $\rho_2 : H^2(X; \mathbb{Z}) \longrightarrow H^2(X; \mathbb{Z}_2)$ is the zero homomorphism,
- (6) $H_2(X; \mathbb{Z})$ is finite, and
- (7) $Sq^k : H^{n-k}(X; \mathbb{Z}_2) \longrightarrow H^n(X; \mathbb{Z}_2)$ is the zero homomorphism for all $k \geq 1$.

Proof. If TX is the tangent bundle of X , then as $w_1(TX) = 0$, we must have that X is orientable proving (1). To prove (2) we note that as $w(TX) = 1$ all Stiefel-Whitney numbers of X must be zero. Hence by a deep theorem of Thom [8, Theorem 4.10, page 53] X is unoriented nullbordant. Since for every $x \in H^1(X; \mathbb{Z}_2)$ there exists a line bundle α with $w_1(\alpha) = x$ we must have that $H^1(X; \mathbb{Z}_2) = 0$ proving (3). That $H_1(X; \mathbb{Z})$ is finite and does not contain elements of order two follows from (3). Next, if

$$\rho_2 : H^2(X; \mathbb{Z}) \longrightarrow H^2(X; \mathbb{Z}_2)$$

is non-zero we choose $x \in H^2(X; \mathbb{Z})$, $x \neq 0$ such that $\rho_2(x) \neq 0$. If α is a complex line bundle with $c_1(\alpha) = x$, then

$$w_2(\alpha_{\mathbb{R}}) = \rho_2(c_1(\alpha)) = \rho_2(x) \neq 0.$$

Here $\alpha_{\mathbb{R}}$ denotes the underlying real bundle of α . This contradiction proves (5). To prove (6), we assume that the dimension of X is not equal to 1, 2, 4, 8 (see Remark 1.3 above). If dimension of X is 3, then $H_2(X; \mathbb{Z}) = 0$ and we are done. So assuming that dimension of X is greater than 4, we have

$$H_i(X; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0 \\ F & i = 1 \\ \mathbb{Z}^\ell \oplus F' & i = 2 \\ \mathbb{Z}^t \oplus F'' & i = 3 \end{cases}$$

where F has no elements of order 2. Since $\rho_2 : H^2(X; \mathbb{Z}) \rightarrow H^2(X; \mathbb{Z}_2)$ is the zero homomorphism and F has no elements of order 2, we have that

$$\beta : H^2(X; \mathbb{Z}_2) \cong \mathbb{Z}_2^\ell \oplus \mathbb{Z}_2^{s'} \rightarrow H^3(X; \mathbb{Z}) \cong \mathbb{Z}^t \oplus F'$$

is a monomorphism where β is the Bockstein associated to

$$0 \rightarrow \mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}_2 \rightarrow 0.$$

This forces $\ell = 0$ and completes the proof of (6). Finally, as X is W -trivial the Wu classes v_k , $k \geq 1$, of X are all zero. As

$$Sq^k x = x \smile v_k = 0$$

for all $x \in H^{n-k}(X; \mathbb{Z}_2)$ we have that (7) is true. This completes the proof of the proposition. \square

Proposition 2.3. Let X be an orientable closed smooth manifold. Then X is W -trivial if any one of the following conditions is satisfied.

- (1) $\widetilde{KO}(X) = 0$,
- (2) X is a \mathbb{Z}_2 -homology n -sphere and n is not a power of 2.

Proof. Recall that $\widetilde{KO}(X)$ is the abelian group of stable isomorphism classes of real vector bundles over X . Two vector bundles α, β over X are stably equivalent if there exist trivial bundles ϵ, ϵ' (possibly of different ranks) such that

$$\alpha \oplus \epsilon \cong \beta \oplus \epsilon'.$$

By the Whitney product theorem we have that $w_i(\alpha) = w_i(\alpha \oplus \epsilon)$ for all $i \geq 0$. Thus stably isomorphic vector bundles have the same Stiefel-Whitney classes. This completes the proof of (1) as $\widetilde{KO}(X) = 0$ implies that every vector bundle on X is stably trivial. The proof of (2) has been outlined in Remark 1.3 (2). This completes the proof of the proposition. \square

Remark 2.4. We mention two remarks related to statements in the Proposition 2.2-2.3.

- (1) As noted after Theorem 1.5, $S^3 \times S^3$ is W -trivial. This also follows from Proposition 2.3 since $\widetilde{KO}(S^3 \times S^3) = 0$.
- (2) The converse of Proposition 2.2 (2) is not true. Examples are provided by the odd dimensional real projective spaces.

In dimension 6, a W -trivial manifold can be a product as we have seen. We note that this is not possible in dimensions 3, 5 and 7.

Proposition 2.5. Let X be a non-empty W -trivial closed smooth manifold of dimension 3, 5, or 7. Then X is not a product of manifolds of positive dimension. \square

Proof. Suppose that X splits as a product of positive dimensional manifolds

$$X = X_1 \times X_2$$

with X_i closed and smooth for $i = 1, 2$. As X is W -trivial so is X_i for $i = 1, 2$. This is because the projection

$$\pi_i : X_1 \times X_2 \longrightarrow X_i$$

induces a monomorphism $\pi_i^* : H^*(X_i; \mathbb{Z}_2) \longrightarrow H^*(X_1 \times X_2; \mathbb{Z}_2)$ for $i = 1, 2$.

As the dimension of X is 3, 5 or 7 it follows that, in each case, one of X_1 or X_2 must have dimension equal to 2^j where $j = 0, 1, 2$. But by Remark 1.3 (1) no manifold of dimension 1, 2 or 4 is W -trivial. This contradiction proves the proposition. \square

We mention that if X, Y are \mathbb{Z}_2 -homology spheres of dimensions m, n respectively with none of $m, n, m+n$ a power of 2, then the product $X \times Y$ is W -trivial. This is because the cohomology of $X \times Y$ is concentrated in degrees 0, m, n and $m+n$. Since none of $m, n, m+n$ are a power of 2, by Remark 1.3 (2), all the Stiefel-Whitney classes of $X \times Y$ in positive degrees are zero.

In general the product of two W -trivial manifolds is not W -trivial. For example, the spheres S^3, S^5 are W -trivial but the product $S^3 \times S^5$ is not W -trivial by Remark 1.3 (1). We however have the following observation about W -triviality of products.

Proposition 2.6. Let X, Y be two orientable closed smooth manifolds with $\widetilde{KO}(X) = 0$ and $\widetilde{KO}(Y) = 0$. If $X \wedge Y$ is W -trivial, then the product $X \times Y$ is W -trivial.

Proof. We shall check that $w(\alpha) = 1$ for every vector bundle α over $X \times Y$. The cofibre sequence

$$X \vee Y \longrightarrow X \times Y \xrightarrow{p} X \wedge Y$$

gives the exact sequence

$$\widetilde{KO}(X \wedge Y) \xrightarrow{p^*} \widetilde{KO}(X \times Y) \longrightarrow \widetilde{KO}(X \vee Y).$$

The last group is zero and hence p^* is surjective. Given a vector bundle α over $X \times Y$, there exists a vector bundle β over $X \wedge Y$ with $p^*(\alpha) \cong \beta$. Since $X \wedge Y$ is W -trivial, we have $w(\beta) = 1$. This forces $w(\alpha) = w(p^*(\beta)) = 1$. This completes the proof. \square

Corollary 2.7. Let X be an orientable closed smooth manifold with $\widetilde{KO}(X) = 0$. Let $n > 8$ be an integer such that $n \equiv 3, 5, 6, 7 \pmod{8}$. Then $S^n \times X$ is W -trivial.

Proof. It is known that $\widetilde{KO}(S^n) = 0$ for $n \equiv 3, 5, 6, 7 \pmod{8}$ [6]. Since $n > 8$, by [1, Theorem 1, page 223], the wedge $S^n \wedge X = \Sigma^n X$ is W -trivial. Hence by the above proposition $S^n \times X$ is W -trivial. This completes the proof. \square

We now turn to the proof of the main theorems. We begin by proving Theorem 1.4 in dimension 3.

Proposition 2.8. Let X be an orientable closed smooth 3-manifold. Then X is W -trivial if and only if it is a \mathbb{Z}_2 -homology 3-sphere.

Proof. We first assume that X is W -trivial. By Proposition 2.2 (4)

$$H_1(X; \mathbb{Z}) = F$$

is finite and does not contain any element of order 2. It follows that $H^1(X; \mathbb{Z}) \cong H_2(X; \mathbb{Z}) = 0$. Hence X is a \mathbb{Z}_2 -homology 3-sphere. The converse follows from Proposition 2.3 (2). This completes the proof. \square

To prove Theorem 1.4 in dimension 5 we shall make use of a result due to Čadek-Vanžura which we state below for easy reference.

Theorem 2.9. [4, Theorem 1, page 757] Let X be a CW -complex of dimension ≤ 5 and suppose

$$\gamma : [X, BSO(5)] \longrightarrow H^2(X; \mathbb{Z}_2) \oplus H^4(X; \mathbb{Z}_2) \oplus H^4(X; \mathbb{Z})$$

is defined by

$$\gamma(\alpha) = (w_2(\alpha), w_4(\alpha), p_1(\alpha)).$$

Then $\text{image}(\gamma) = \{(a, b, c) : \rho_4(c) = \mathfrak{P}(a) + i_*(b)\}$. Here $i_* : H^*(X; \mathbb{Z}_2) \longrightarrow H^*(X; \mathbb{Z}_4)$ is induced by the coefficient injection $\mathbb{Z}_2 \longrightarrow \mathbb{Z}_4$. \square

Proposition 2.10. Let X be an orientable closed smooth 5-manifold. Then X is W -trivial if and only if it is a \mathbb{Z}_2 -homology 5-sphere.

Proof. Assume that X is W -trivial. We shall show that X is a \mathbb{Z}_2 -homology 5-sphere. We first observe, using Proposition 2.2, that the integral homology groups of X are of the form

$$H_i(X; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0 \\ F & i = 1 \\ F' & i = 2 \\ \mathbb{Z}^t \oplus F'' & i = 3 \\ \mathbb{Z}^{\ell'} & i = 4 \\ \mathbb{Z} & i = 5 \end{cases}$$

where we recall that F, F', F'' are finite abelian groups with F not containing any element of order 2. Since X is orientable we have, by Poincaré duality, that

$$t = 0, \quad \ell' = 0, \quad F = F''.$$

Hence F'' does not contain any element of order 2. Thus the mod-2 cohomology groups of X are of the form

$$H^i(X; \mathbb{Z}_2) = \begin{cases} \mathbb{Z}_2 & i = 0 \\ 0 & i = 1 \\ \mathbb{Z}_2^{s'} & i = 2 \\ \mathbb{Z}_2^{s'} & i = 3 \\ 0 & i = 4 \\ \mathbb{Z}_2 & i = 5. \end{cases}$$

The proof that X is a \mathbb{Z}_2 -homology 5-sphere will be complete if we can show that $s' = 0$. Notice that as $\ell' = 0$ and $F'' = F$ do not contain any element of order 2 we have that

$$H^4(X; \mathbb{Z}_4) = 0.$$

This implies that every tuple (a, b, c) with $a \in H^2(X; \mathbb{Z}_2), b \in H^4(X; \mathbb{Z}_2), c \in H^4(X; \mathbb{Z})$ is in the image of γ . Thus if $s' \neq 0$, then every non zero cohomology class $a \in H^2(X; \mathbb{Z}_2)$ equals $w_2(\alpha)$ for some vector bundle α over X . This contradiction shows that $s' = 0$. This completes the proof that X is a \mathbb{Z}_2 -homology 5-sphere.

The converse follows from Proposition 2.3 (2). This together with Proposition 2.8 completes the proof of Theorem 1.4. \square

We now turn to the proof of Theorem 1.5. We shall need the following theorem.

Theorem 2.11. [5, Proposition 2, page 729] Let X be a connected CW -complex of dimension ≤ 7 and suppose

$$\gamma : [X, BSO(7)] \longrightarrow H^2(X; \mathbb{Z}_2) \oplus H^4(X; \mathbb{Z}_2) \oplus H^4(X; \mathbb{Z})$$

is defined by

$$\gamma(\alpha) = (w_2(\alpha), w_4(\alpha), p_1(\alpha)).$$

Then $\text{image}(\gamma) = \{(a, b, c) : \rho_4(c) = \mathfrak{P}(a) + i_*(b)\}$. Here $i_* : H^*(X; \mathbb{Z}_2) \longrightarrow H^*(X; \mathbb{Z}_4)$ is induced by the coefficient injection $\mathbb{Z}_2 \longrightarrow \mathbb{Z}_4$. \square

Proof of Theorem 1.5. We assume that X is a W -trivial orientable closed smooth 6-manifold. Then we shall show that the integral homology groups of X are of the required form. To begin with, the integral homology groups of X are of the form

$$H_i(X; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0 \\ F & i = 1 \\ F' & i = 2 \\ \mathbb{Z}^t \oplus F'' & i = 3 \\ \mathbb{Z}^{\ell'} \oplus F''' & i = 4 \\ \mathbb{Z}^{\ell''} & i = 5 \\ \mathbb{Z} & i = 6 \end{cases}$$

where we recall that F, F', F'' and F''' are finite abelian groups with F not containing any element of order 2. Since X is orientable, we have

$$\ell'' = 0, \quad F = F''', \quad \ell' = 0, \quad F' = F''.$$

Thus F''' does not contain any element of order 2. The proof will be complete if we can show that $s' = 0$. We assume that $s' > 0$. This leads to a contradiction as follows. First observe that

$$H^4(X; \mathbb{Z}_4) \cong \mathbb{Z}_2^{u'} \oplus \mathbb{Z}_4^{v'}.$$

Identifying the cohomology groups in the long exact sequence

$$\longrightarrow H^4(X; \mathbb{Z}) \longrightarrow H^4(X; \mathbb{Z}) \xrightarrow{\rho_4} H^4(X; \mathbb{Z}_4) \xrightarrow{\beta} H^5(X; \mathbb{Z})$$

associated to the short exact sequence of coefficient groups

$$0 \longrightarrow \mathbb{Z} \xrightarrow{\times 4} \mathbb{Z} \longrightarrow \mathbb{Z}_4 \longrightarrow 0$$

we get the exact sequence

$$\cdots \longrightarrow F' \longrightarrow F' \xrightarrow{\rho_4} \mathbb{Z}_2^{u'} \oplus \mathbb{Z}_4^{v'} \xrightarrow{\beta} F \longrightarrow \cdots.$$

Since F has no elements of order 2, we have that F has no elements of order 4. This implies that the homomorphism β is the zero homomorphism and hence

$$\rho_4 : H^4(X; \mathbb{Z}) \longrightarrow H^4(X; \mathbb{Z}_4)$$

is an epimorphism. Next observe that as $s' \neq 0$ we have

$$H^2(X; \mathbb{Z}_2) \cong \mathbb{Z}_2^{s'} \neq 0.$$

Let $a \in H^2(X; \mathbb{Z}_2)$ with $a \neq 0$. Then $\mathfrak{P}(a) \in H^4(X; \mathbb{Z}_4)$. As ρ_4 is an epimorphism, we choose $c \in H^4(X; \mathbb{Z})$ such that

$$\rho_4(c) = \mathfrak{P}(a).$$

Then, the 3-tuple $(a, 0, c)$ is in the image of γ where γ is the function defined in Theorem 2.11. Thus there exists a vector bundle α over X with $w_2(\alpha) = a \neq 0$. This contradiction implies that $s' = 0$ and completes the proof that the integral homology of X has the required form.

Conversely suppose that the integral homology of X has the form in the theorem. Then the mod-2 cohomology groups satisfy $H^i(X; \mathbb{Z}_2) = 0$ for $i = 1, 2, 4$. This clearly forces X to be W -trivial. This completes the proof of Theorem 1.5. \square

Next, we turn our attention to 7-manifolds. We also recall the following theorem which we shall use.

Theorem 2.12. [5, Theorem 3, page 733] Let X be a CW complex of dimension ≤ 7 and let $P \in H^4(X; \mathbb{Z})$, $W \in H^4(X; \mathbb{Z}_2)$. Then there exists an oriented rank 5 bundle α over X with

$$w_2(\alpha) = 0, \quad w_4(\alpha) = W, \quad p_1(\alpha) = P$$

if and only if there is $U \in H^4(X; \mathbb{Z})$ such that

- (1) $P = 2U$, $\rho_2(U) = W$,
- (2) $Sq^2(W) = 0$, and
- (3) $0 \in \Phi(U)$

where Φ is the secondary cohomology operation from $H^4(X; \mathbb{Z})$ into $H^7(X; \mathbb{Z}_2)$ associated to the relation $Sq^2 \circ Sq^2 \rho_2 = 0$. \square

Before proving Theorem 1.6, we make a few remarks on the secondary cohomology operation Φ . Given a space Y , Φ is defined on the kernel of the homomorphism

$$Sq^2 \circ \rho_2 : H^4(Y; \mathbb{Z}) \longrightarrow H^6(Y; \mathbb{Z}_2)$$

and takes values in $H^7(Y; \mathbb{Z}_2)/Q$ where $Q = Sq^2 H^5(Y; \mathbb{Z}_2)$. If $u \in H^4(Y; \mathbb{Z})$ is in the kernel of $Sq^2 \circ \rho_2$, then $\Phi(u)$ is defined to be the coset

$$\Phi(u) = U^*(x) + Sq^2(H^5(Y; \mathbb{Z}_2))$$

where U is the lift of

$$Sq^2 \circ \rho_2 \circ u : X \longrightarrow K(\mathbb{Z}_2, 6)$$

to E , where E is the pull back of the path fibration over $K(\mathbb{Z}_2, 6)$ to $K(\mathbb{Z}, 4)$ via $Sq^2 \circ \rho_2$. Here $x \in H^7(E; \mathbb{Z}_2) \cong \mathbb{Z}_2$ is a generator. Note that if $H^6(Y; \mathbb{Z}_2) = 0$, then $\Phi(u)$ is defined for all $u \in H^4(X; \mathbb{Z})$.

Proof of Theorem 1.6. Let X be an orientable closed smooth 7-manifold. Assuming that X is W -trivial and using Proposition 2.2, the integral homology groups of X can be readily seen to be of the form

$$H_i(X; \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0 \\ F & i = 1 \\ F' & i = 2 \\ \mathbb{Z}^t \oplus F'' & i = 3 \\ \mathbb{Z}^t \oplus F' & i = 4 \\ F & i = 5 \\ 0 & i = 6 \\ \mathbb{Z} & i = 7 \end{cases}$$

where F, F', F'' are finite abelian groups with F not containing any element of order 2. In particular, we note that $H^6(X; \mathbb{Z}_2) = 0$ and hence the secondary operation Φ is defined on the whole of $H^4(X; \mathbb{Z})$.

To complete the proof of the theorem it is enough to prove that

$$t = s'' = 0.$$

To prove this we first claim that

$$\rho_2 : H^4(X; \mathbb{Z}) \longrightarrow H^4(X; \mathbb{Z}_2)$$

is the zero homomorphism. Assuming the contrary, suppose that there exists $U \in H^4(X; \mathbb{Z})$, $U \neq 0$ with $\rho_2(U) = W \neq 0$. This leads to a contradiction as follows. If we let $P = 2U$ we see that the first condition of Theorem 2.12 is satisfied. Since $H^6(X; \mathbb{Z}_2) = 0$, it follows that $Sq^2(W) = 0$. We may now check that $0 \in \Phi(U)$ as in [9, page 206]. Thus by Theorem 2.12 there exists a vector bundle α over X with

$$w_4(\alpha) = W \neq 0.$$

This contradiction proves the claim. We may now look at the exact sequence

$$H^4(X; \mathbb{Z}) \xrightarrow{\rho_2} H^4(X; \mathbb{Z}_2) \xrightarrow{\beta} H^5(X; \mathbb{Z}) \longrightarrow .$$

Then as $\rho_2 = 0$ we have that

$$\beta : \mathbb{Z}_2^t \oplus \mathbb{Z}_2^{s'} \oplus \mathbb{Z}_2^{s''} \longrightarrow F'$$

is a monomorphism. This implies, as before, that

$$t + s'' = 0$$

and hence $t = s'' = 0$.

Conversely, if the homology of X is of the given form and if additionally, F' contains no elements of order 2, then X is a \mathbb{Z}_2 -homology sphere and hence W -trivial by Proposition 2.3 (2). This completes the proof.

We have not been able to decide whether or not F' can contain any element of order 2. We derive the following necessary conditions for a closed smooth 7-manifold to be W -trivial.

Corollary 2.13. If X is a W -trivial closed smooth 7-manifold, then

- (1) The map

$$Sq^2 : H^2(X; \mathbb{Z}_2) \longrightarrow H^4(X; \mathbb{Z}_2)$$

given by $Sq^2(x) = x \smile x$ is an isomorphism.

- (2) The Pontryagin square

$$\mathfrak{P} : H^2(X; \mathbb{Z}_2) = \mathbb{Z}_2^{s'} \longrightarrow H^4(X; \mathbb{Z}_4) = \mathbb{Z}_2^{u'} \oplus \mathbb{Z}_4^{v'}$$

is a monomorphism.

Proof. By Theorem 1.6

$$H^2(X; \mathbb{Z}_2) \cong \mathbb{Z}_2^{s'} \cong H^4(X; \mathbb{Z}_2).$$

Now if $f(x) = x^2 = 0$ for some non-zero $x \in H^2(X; \mathbb{Z}_2)$, then by [14, Theorem 1, page 176] there exists a vector bundle α with $w_2(\alpha) = x \neq 0$. This contradiction proves (1).

Next, let

$$\rho_{4,2} : H^4(X; \mathbb{Z}_4) \longrightarrow H^4(X; \mathbb{Z}_2)$$

be the homomorphism induced by the coefficient surjection. Then as

$$\rho_{4,2} \circ \mathfrak{P} = Sq^2,$$

(2) follows from (1). This completes the proof. \square

Proposition 2.14. If X, Y are closed smooth manifolds of dimension $n = 3, 5$ or 6 and are W -trivial, then so is their connected sum.

Proof. First note that if X, Y are two orientable closed smooth n -manifolds, then their connected sum $X \# Y$ is also an orientable closed smooth manifold and

$$H_i(X \# Y; R) \cong H_i(X; R) \oplus H_i(Y; R)$$

for $i \neq n, n-1$ and all commutative rings R .

Assume that X, Y are 3-dimensional. Then by Theorem 1.4, X and Y are \mathbb{Z}_2 -homology spheres. Since

$$H_1(X \# Y; \mathbb{Z}_2) \cong H_1(X; \mathbb{Z}_2) \oplus H_1(Y; \mathbb{Z}_2) = 0$$

it follows that $H^1(X \# Y; \mathbb{Z}_2) = 0$. By Poincaré duality we have that $H_2(X \# Y; \mathbb{Z}_2) = 0$. Thus $X \# Y$ is a \mathbb{Z}_2 -homology 3-sphere. Hence by Theorem 1.4, $X \# Y$ is W -trivial.

The proof in the case that X, Y are 5-dimensional is similar and we omit the proof. Finally we assume that X, Y are 6-dimensional. The integral homology groups of X and Y are as given in Theorem 1.5. It can now be verified that the integral homology groups of the connected sum $X \# Y$ is also of the form as given in Theorem 1.5 and hence $X \# Y$ is W -trivial. \square

Acknowledgements. We express our appreciation to the anonymous referee for her/his detailed comments and suggestions which has improved the presentation and corrected several inaccuracies. In particular, we would like to thank the referee for pointing out the reference [13] and suggesting a way to shorten the proof of Proposition 2.10.

3. CONFLICT OF INTEREST STATEMENT

On behalf of all authors, the corresponding author states that there is no conflict of interest.

4. DATA AVAILABILITY STATEMENT

We do not analyse or generate any datasets, because our work proceeds within a theoretical and mathematical approach.

REFERENCES

- [1] Atiyah, M., and Hirzebruch, F., *Bott periodicity and the parallelizability of the spheres*, Proc. Cambridge Philos. Soc., 57 (1961) 223-226.
- [2] Barden, D., *Simply connected 5-manifolds*, Ann. Math., vol 82, no. 3 (19), pp. 365-385.
- [3] Borel, A., *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math., 57 (1953), 115-207.
- [4] Čadek, M., and Vanžura, J., *On the classification of oriented bundles over 5-complexes*, Czechoslovak Math. J., 43 (118) (1993), no. 4, 753-764.
- [5] Čadek, M., and Vanžura, J., *On oriented vector bundles over CW-complexes of dimension 6 and 7*, Comment. Math. Univ. Carolin., 33 4 (1992), 727-736.
- [6] Husemoller D., *Fibre Bundles*, Springer-Verlag, New York 1966.
- [7] Kervaire, Michel, A., *Smooth homology spheres and their fundamental groups*, Trans. Amer. Math. Soc., Vol. 144 (Oct., 1969), pp. 67-72
- [8] Milnor, J., *Some consequences of a theorem of Bott*, Ann. of Math., Vol 68, No. 2 (1958) 444-449.
- [9] Naolekar, A. C., *Euler classes of vector bundles over manifolds*, Math. Slovaca 71 (2021), No. 1, 199-210.
- [10] Naolekar, A. C., Subhash, B., Thakur, Ajay Singh, *On trivialities of Euler classes of oriented vector bundles over manifolds*, HHA, vol. 22(1), 2020, pp. 215-232..
- [11] Ruberman, D., *Null-homotopic embedded spheres of codimension one*, Tight and Taut submanifolds, MSRI Publications, vol. 32, 1997 pp. 229-232
- [12] Tanaka, R., *On trivialities of Stiefel-Whitney classes of vector bundles over iterated suspension spaces*, Homology Homotopy Appl., 12 (2010), no. 1, 357-366.

- [13] Thomas, E., *A generalization of the Pontrjagin square cohomology operation*, Proc. Nat. Acad. Sci. U.S.A., 42 (1956), 266-269.
- [14] Woodward, L. M., *The classification of orientable vector bundles over CW-complexes of low dimensions*, Proc. Royal Soc. Edinburgh Sect. A 92 (1982), no. 3-4, 175-179.

619 RED CEDAR ROAD, WELLS HALL, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI-48824, USA.

Email address: `bhatta94@msu.edu`

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, ROORKEE, HARIDWAR ROAD, 247667, INDIA.

Email address: `bikramju@gmail.com`, `bikramjit.pd@ma.iitr.ac.in`

STAT-MATH UNIT, INDIAN STATISTICAL INSTITUTE, 8TH MILE, MYSORE ROAD, RVCE POST, BANGALORE 560059, INDIA.

Email address: `ani@isibang.ac.in`