

Data-driven fixed-point tuning for truncated realized variations

B. Cooper Boniece*

José E. Figueroa-López[†]

Yuchen Han[‡]

May 20, 2024

Abstract

Many methods for estimating integrated volatility and related functionals of semimartingales in the presence of jumps require specification of tuning parameters for their use in practice. In much of the available theory, tuning parameters are assumed to be deterministic and their values are specified only up to asymptotic constraints. However, in empirical work and in simulation studies, they are typically chosen to be random and data-dependent, with explicit choices often relying entirely on heuristics. In this paper, we consider novel data-driven tuning procedures for the truncated realized variations of a semimartingale with jumps based on a type of random fixed-point iteration. Being effectively automated, our approach alleviates the need for delicate decision-making regarding tuning parameters in practice and can be implemented using information regarding sampling frequency alone. We show our methods can lead to asymptotically efficient estimation of integrated volatility and exhibit superior finite-sample performance compared to popular alternatives in the literature.

1 Introduction

The continuous part of the quadratic variation of an Itô semimartingale, commonly known as the integrated volatility, plays an outsize role in financial econometrics, and its estimation in various settings based on discrete observations has been a major focus in the literature at various points in the past 20+ years. The semimartingale X commonly represents the log-price of a financial asset, and its integrated volatility serves as a measure of the overall uncertainty inherent in the continuous part of X over a given time period.

Among the variety of available methods for integrated volatility estimation, the truncated realized variation (TRV), introduced in [30], was one of the first and remains among the most popular approaches to-date that is *jump-robust*, in the sense that it can still provide reliable estimates of integrated volatility when jumps occur in the process X . Other well known jump-robust methods for estimating integrated volatility include bipower variations and their extensions [5, 7, 11] or those based on empirical characteristic functions [24, 25, 38], among others, giving the practitioner a wide array of choices at their disposal for estimation of integrated volatility in modeling contexts where jumps may be present.

To choose an estimator among this array of options, currently one must first decide between two distinct classes: either asymptotically efficient approaches, like TRV, which require selection

*Department of Mathematics, Drexel University, Philadelphia, PA 19104, USA (cooper.boniece@drexel.edu). Research supported in part by NSF grant DMS-2309570.

[†]Department of Statistics and Data Science, Washington University in St. Louis, St. Louis, MO 63130, USA (figueroa-lopez@wustl.edu). Research supported in part by NSF grant DMS-2015323.

[‡]Washington University in St. Louis, St. Louis, MO 63130, USA

of tuning parameters, or alternatively “tuning-free” estimators but at the unfortunate expense of asymptotic efficiency. From the perspective of minimizing variance, asymptotically efficient approaches are preferable, but their use in practice necessitates the critical additional step of specifying the tuning parameter values themselves. This consequential step can significantly impact estimation performance, but current asymptotic theory does not offer direct guidelines for choosing parameters explicitly, which can be an extremely delicate matter in practice. For instance, even in idealized asymptotic settings, appropriate choices often depend on a priori unknown properties of X and can determine whether or not a given estimator retains even the basic requirement of consistency. In the absence of theoretically supported approaches for specifying explicit values of these parameters, the practical use of tuning-parameter-based methods remains entirely reliant on heuristics. The purpose of the present work is to address this gap.

In the case of TRV, the tuning parameter of importance is called the *threshold*, denoted hereafter as $\varepsilon > 0$, indicating a level above which increments are discarded from the estimation procedure. Concretely, given a discretely observed semimartingale $X = \{X_t\}_{t \geq 0}$ at times $0 = t_0 < t_1 < \dots < t_n = T$, the TRV is defined as

$$\text{TRV}_n(\varepsilon) = \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq \varepsilon\}},$$

where $\Delta_i^n X := X_{t_i} - X_{t_{i-1}}$ is the i^{th} increment of X , often assumed to be observed on a regular sampling grid, so that $t_i - t_{i-1} =: h_n$ for all i . Statistical properties of TRV have been extensively studied when $\varepsilon = \varepsilon(h_n)$ is a *deterministic* function of the time step h_n such that $\varepsilon(h_n) \rightarrow 0$ at specified rates as $h_n \rightarrow 0$. In [31], when either the jump component of the process X is of finite activity or is a pure-jump Lévy process with infinite jump activity, TRV was shown to be consistent whenever

$$\lim_{h_n \rightarrow 0} \varepsilon(h_n) = 0, \quad \text{and} \quad \lim_{h_n \rightarrow 0} \frac{\varepsilon(h_n)}{\sqrt{h_n \log \frac{1}{h_n}}} = \infty. \quad (1.1)$$

A consistency statement for TRV encompassing a broader class of semimartingales was given in [22], but for the more restrictive case of power thresholds, namely, thresholds of the form

$$\varepsilon(h_n) = ch_n^\beta, \quad c > 0, \quad 0 < \beta < 1/2. \quad (1.2)$$

Under finite jump activity, central limit theorems for TRV were established under the threshold constraint (1.1) in [31]; in the infinite-activity case, they were established in [22] for more general semimartingales based on thresholds satisfying (1.2) under the additional assumption the volatility itself is a semimartingale, and also in [10, 32] for general càdlàg volatility processes but for Lévy-type jump behavior, both under additional constraints on ε related to the Blumenthal-Gettoor index of X .

While asymptotic constraints such as (1.1) and (1.2) may be informative for threshold selection, they do not concretely indicate how one should make an explicit choice for ε in a given context. Moreover, even if a particular deterministic choice for ε may lead to good estimation performance under a given model, the same choice of ε under a perturbed version of the same model can lead to dramatically worse estimation performance. To illustrate this point, the left panel of Figure 1, below, shows histograms of the relative estimation errors for TRV using a fixed, deterministically chosen threshold value under two different parameter settings of the same model. While TRV performs satisfactorily with this deterministic threshold value under one of the parameter settings, it performs poorly with the same threshold value under alternate parameter settings, even though the expected quadratic variation of X is the same in both

cases. In contrast, the right panel of Figure 1 displays histograms of relative estimation errors for the approach developed in this paper, where satisfactory performance is maintained across both settings.

Though Monte Carlo studies or empirical insights may help in choosing the value of ε deterministically in a given setting, an arguably more natural approach is to select thresholds through some data-driven procedure, permitting the threshold itself to depend on observed data. Indeed, random, data-driven parameter tuning is often done in numerical studies in the literature – without theoretical support – to illustrate finite-sample behavior of estimators and to improve their numerical performance.




Figure 1: Sampling distributions of the relative estimation error for TRV with deterministic thresholding at a fixed threshold value ε (left panel) versus our automated thresholding procedure¹ (right panel). The average realized quadratic variation is the same in both models. The reference setting is the same as Model 1 as described in Section 4, based on a 1-week time horizon at a sampling frequency of 5 minutes. The perturbed reference setting uses the same parameter settings as the reference setting, but with half overall (average) volatility level and roughly twice the rate of its finite jump activity component (adjusted to match the average quadratic variation of both models).

However, by their very nature, data-driven parameter selection procedures introduce considerable statistical dependencies and associated theoretical challenges that are otherwise absent when parameters are chosen deterministically. Consequently, despite the practicality and potential benefits of data-driven parameter selection, the literature on TRV and related methods employing data-driven tuning procedures has remained relatively scarce. For instance, in the case of finite activity jumps, it was stated without proof in a remark in [35] that consistency holds for time-dependent random thresholds (possibly different for each increment $\Delta_i^n X$) of the form $c_{t_i}\varepsilon$, where $\varepsilon = \varepsilon(h_n)$ satisfies (1.1) and $\{c_t\}_{t \geq 0}$ is a stochastic process that is a.s. bounded above and bounded away from 0. Later, in [18], consistency was rigorously established under finite jump activity for possibly data-dependent time-varying thresholds of the type $\sqrt{2(1+\eta)M_i h_n \log(1/h_n)}$, where $\eta > 0$ and M_i are random variables satisfying $M_i \in [\inf_{s \in [t_{i-1}, t_i]} \sigma_s^2, \sup_{s \in [0, T]} \sigma_s^2]$ a.s. To the authors' knowledge, these statements comprise the totality of asymptotic theory for TRV with data-driven thresholds, and there is currently no theoretical support in the literature for data-driven parameter tuning of TRV outside consistency statements in the finite activity setting.

Moreover, in spite of the considerable focus on asymptotic properties of TRV with the threshold constraints (1.1) and (1.2), recent work [18, 19] has demonstrated that certain *optimal* choices of threshold do not satisfy these asymptotic conditions, leaving a substantive gap in the available asymptotic theory even within the scope of deterministic thresholding. Optimal-type thresholds can lead to substantial gains in finite sample estimation performance, and their explicit expressions can serve as a more direct guideline for threshold selection, making them

¹Specifically our approach as described in (6b) in Section 4, though similar behavior holds in all other cases.

ideal choices for practitioners. However, their direct use, even to first-order approximation, is complicated by the fact that they depend on the volatility itself. For instance, under an idealized constant volatility assumption and general finite jump activity, the MSE-optimal threshold ε_n^* admits the approximation:

$$\varepsilon_n^* \sim \sqrt{2\sigma^2 h_n \log \frac{1}{h_n}}, \quad \text{as } h_n \rightarrow 0, \quad (1.3)$$

where $\sigma > 0$ is the volatility. Under Lévy stable-like infinite jump activity, the MSE-optimal threshold is the same as ε_n^* up to an additional multiplicative constant depending on the Blumenthal-Gettoor index [15]. Though this expression cannot be used directly in practice due to its dependence on knowledge of the volatility, it lends itself naturally to fixed-point iterative procedures, as suggested in [19] and [18], whose asymptotic theory has remained unestablished, until now.

In this work, we consider two classes of iterative procedures for jump-robust estimation of the integrated volatility based on data-driven parameter tuning. Our procedures are designed to turn the otherwise infeasible threshold (1.3) into a feasible one, and will be seen to stem from solutions ξ to random fixed-point equations of the type

$$\xi = \Phi_n \left((\Delta_1^n X) \mathbf{1}_{\{|\Delta_1^n X| \leq \sqrt{r_n \xi}\}}, \dots, (\Delta_n^n X) \mathbf{1}_{\{|\Delta_n^n X| \leq \sqrt{r_n \xi}\}} \right), \quad (1.4)$$

for an appropriate function Φ_n and sequence $r_n \rightarrow 0$. Viewed as random, data-dependent thresholds, our procedures extend the asymptotic theory beyond deterministic thresholding to accommodate automatic, data-driven calibration of TRV, and further extends current asymptotic theory beyond the general rate constraints imposed in (1.1) and (1.2), allowing for time-dependent thresholding, ultimately leading to substantial gains in finite-sample performance and more principled threshold selection procedures. Part of our analysis is based on relating our proposed iterative estimators to oracle-like sequences of estimators; this general approach may be of use for parameter tuning in other jump-robust methods in the literature.

This paper is organized as follows. Section 2 introduces the model, estimation framework and some notation used throughout the paper. Section 3 contains our main results, including instances of uniform and time-varying thresholding, and Section 4 contains some numerical illustrations concerning finite-sample estimation performance. The proofs of the main results and auxiliary lemmas are given in Appendices A and B.

2 Framework and setting

We consider a 1-dimensional Itô semimartingale $X = (X_t)_{t \in \mathbb{R}_+}$ defined on a complete filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$ of the form

$$dX_t = b_t dt + \sigma_t dW_t + \gamma_t dL_t + dJ_t, \quad t \geq 0. \quad (2.1)$$

Above, W is a standard Brownian motion, b, γ, σ are càdlàg adapted, $L = \{L_t\}_{t \geq 0}$ is a pure-jump infinite-activity Lévy process, and $J = \{J_t\}_{t \geq 0}$ is a general pure-jump process with finite jump activity. We refer to Assumption 2.1 for complete conditions on all driving processes and coefficients.

We suppose that on a fixed and finite time interval $[0, T]$, n observations, $X_{t_1}, X_{t_2}, \dots, X_{t_n}$, of the continuous-time process X are available at known times $0 = t_0 < t_1 < \dots < t_n = T$. We assume sampling times are evenly spaced, and denote the time step between observations as $h_n := T/n$. Our estimation target is the integrated volatility (or integrated variance) of X defined as

$$C_T := \int_0^T \sigma_s^2 ds.$$

We consider two classes of estimators of C_T . The first class of estimators we consider are based on an iterative scheme that proceeds as follows:

1. At the start, an initial guess $\widehat{C}_{n,0}$ for the integrated volatility C_T is first put forward. This initial estimate should ideally be free of tuning parameters; for instance, the realized variance (RV) $\widehat{C}_{n,0} = \sum_{i=1}^n (\Delta_i^n X)^2$ or bipower variation $\widehat{C}_{n,0} = \frac{\pi}{2} \sum_{i=1}^{n-1} |\Delta_i^n X| |\Delta_{i+1}^n X|$, among other possibilities. We refer to Theorem 3.2 and Remark 3.1 for further information about the conditions on $\widehat{C}_{n,0}$. This initializes a data-dependent threshold of the form $B_{n,0} = (r_n T^{-1} \widehat{C}_{n,0})^{1/2}$, where the *threshold rate* $r_n = r_n(h_n)$ is some predetermined deterministic function of h_n .
2. Then, an iterative sequence of thresholds $B_{n,j}$ and estimators $\widehat{C}_{n,j}$ is constructed based on the relations

$$\begin{aligned} B_{n,j-1} &:= \sqrt{r_n T^{-1} \widehat{C}_{n,j-1}}, \\ \widehat{C}_{n,j} &:= \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_{n,j-1}\}}, \quad j \geq 1. \end{aligned} \tag{2.2}$$

For each fixed n , the sequence $\{\widehat{C}_{n,j}, j \geq 0\}$ will be shown to always “stabilize” in the sense that the index

$$j_n := \min \{j \geq 0 : \widehat{C}_{n,j} = \widehat{C}_{n,j+\ell} \text{ for all } \ell \geq 0\}, \tag{2.3}$$

always exists. With regards to (1.4), the above scheme can be viewed as a fixed-point iteration for the function $\xi \mapsto \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{(\Delta_i^n X)^2 \leq r_n T^{-1} \xi\}}$.

3. Denoting $B_n := B_{n,j_n}$, we then define the uniform thresholding estimator

$$\widehat{C}_n := \widehat{C}_{n,j_n} = \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_n\}}. \tag{2.4}$$

The second class of estimators we consider are based on time-varying (or local) thresholding, namely estimators $\widehat{C}_n^*$ of the type

$$\widehat{C}_n^* = \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_n^*(i)\}}, \tag{2.5}$$

for appropriate data-driven local thresholds $B_n^*(i)$, $i = 1, \dots, n$ arising from fixed-point equations, whose precise definition is deferred to Section 3.2. The central focus of this work is to study the classes of estimators defined by (2.4) and (2.5). Below we state our main assumptions relating to the model (2.1).

Assumption 2.1.

- (i) σ, γ, b are càdlàg adapted; $\inf_{0 \leq t \leq T} \sigma_t > 0$;
- (ii) L is a Lévy process with characteristics $(0, 0, \nu)$ (see [37]) such that, for some $\alpha \in (0, 2)$ and $K_{\pm} \in [0, \infty)$, $K_+ \vee K_- > 0$,

$$\lim_{x \rightarrow 0^+} x^\alpha \nu((x, \infty)) = K_+, \quad \lim_{x \rightarrow 0^-} x^\alpha \nu((-\infty, x)) = K_-;$$

J is a general finite-activity jump process of the form $J_t = \sum_{i=1}^{N_t'} \xi_i$, where $\{\xi_i\}_{i \geq 1}$ satisfy $\mathbb{P}(\xi_i \neq 0) = 1$, and $\{N_t'\}_{t \geq 0}$ is a non-explosive counting process; W is a Brownian motion independent of L . All processes are adapted.

(iii) There is a sequence τ_n of stopping times increasing to infinity and a positive sequence K_n such that

$$t \leq \tau_n \implies \begin{cases} |b_t| + |\sigma_t| + |\gamma_t| \leq K_n, \\ \mathbb{E}(|\gamma_{t+s} - \gamma_t|^2 | \mathcal{F}_t) \leq K_n s, \quad s > 0. \end{cases} \quad (2.6)$$

Our assumptions, in particular, do not require σ to be a semimartingale, which is important in rough volatility modeling. Note that the condition on γ in (2.6) is satisfied whenever $\{\gamma_t\}_{t \geq 0}$ is an Itô semimartingale with locally bounded characteristics.

We use the following standard notation throughout the paper: for two sequences $a_n, b_n > 0$,

- $a_n \sim b_n$ means that $a_n/b_n \rightarrow 1$;
- $a_n \ll b_n$ means that $a_n = o(b_n)$, i.e., $\lim_{n \rightarrow \infty} a_n/b_n = 0$; $a_n \lesssim b_n$ means $a_n = O(b_n)$, i.e., $\limsup_{n \rightarrow \infty} a_n/b_n < \infty$;
- $a_n \gg b_n$ means that $b_n = o(a_n)$; $a_n \gtrsim b_n$ means $b_n = O(a_n)$;
- $\xrightarrow{P}$ denotes convergence in probability;
- $\xrightarrow{\mathcal{D}}$ denotes convergence in law;
- $\xrightarrow{st}$ denotes stable convergence in law.

3 Main results

3.1 Uniform thresholding

In this section, we study the asymptotic properties of the estimator $\hat{C}_n$ introduced in Section 2. As stated in the introduction, in [18] it was shown that the first-order asymptotic behavior of the MSE-optimal threshold ε_n^* under the idealized assumption of constant volatility takes the form (1.3), which cannot be implemented feasibly in practice as it depends on knowledge of the volatility itself. However, exploiting this relationship is the driving principle behind the iterative algorithm leading to the estimators $\hat{C}_n$. The proposed method can be seen as a natural mechanism to make such a threshold feasible by taking the sequence $r_n = r(h_n) = 2h_n \log(1/h_n)$ in the iterative procedure (2.2). As we will see, our iterative approach will enable us to asymptotically “attain” the infeasible threshold ε_n^* , in principle rendering near-MSE-optimal behavior possible in practice.

In general, it is a nontrivial task to establish asymptotic properties of the $\hat{C}_n$ defined in (2.4), even drawing upon results from the existing literature, which has almost exclusively focused on deterministic uniform thresholding. For instance, in spite of the fact that $\hat{C}_n$ satisfies the random fixed-point equation

$$\hat{C}_n = \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{(\Delta_i^n X)^2 \leq r_n T^{-1} \hat{C}_n\}},$$

such an expression offers little insight into finding closed-form expressions for $\hat{C}_n$.

The central idea in our approach rests on relating the sequence of iterates $\hat{C}_{n,j}$ to an iterative sequence of “oracle-like” estimators $\tilde{C}_{n,j}(y_n)$ that make use of the (unknown) location of jumps of size $y_n > 0$ or larger, where $y_n \rightarrow 0$ at an appropriate rate. More concretely, for each $y \in (0, 1)$, by virtue of the Lévy-Itô decomposition of L , we may reexpress

$$\begin{aligned} b_t dt + \gamma_t dL_t &= \left(b_t + \gamma_t \int_{\{y < |x| \leq 1\}} x \nu(dx) \right) dt + \gamma_t \int_{\{|x| \leq y\}} x \tilde{\mu}(dx, dt) + \gamma_t \int_{\{|x| > y\}} x \mu(dx, dt) \\ &=: db_t(y) + \gamma_t dM_t(y) + \gamma_t dH_t(y), \end{aligned} \quad (3.1)$$

where μ is the jump measure of L with intensity $\nu(dx)dt$, and $\tilde{\mu}(dx, dt) = \mu(dx, dt) - \nu(dx)dt$ is the corresponding compensated jump measure. Above, $H_t(y)$ is a compound Poisson process with finite jump activity satisfying $H_t(y) = \sum_{i=1}^{N_t(y)} \zeta_i(y)$, where $N_t(y)$ is a Poisson process with rate $\lambda_n(y) = \int_{|x|>y} \nu(dx)$, and $\{\zeta_i(y)\}_{i \geq 1}$ are i.i.d. and supported on $(-\infty, y) \cup (y, \infty)$ with distribution $\frac{\mathbf{1}_{\{|x|>y\}} \nu(dx)}{\nu(|x|>y)}$. For each $y > 0$, we first define the random set

$$\mathcal{I}_n(y) = \{i : \Delta_i^n N(y) = 0, \Delta_i^n N' = 0\}, \quad (3.2)$$

which consists of all indices corresponding to intervals where no “large” jumps have occurred. For a sequence $y = y_n \rightarrow 0$, we then define an oracle analog of TRV that eliminates any increments corresponding to time intervals in which “large” jumps of X occur:

$$\mathcal{E}_n(y) := \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{\Delta_i^n N(y)=0, \Delta_i^n N'=0\}} = \sum_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2. \quad (3.3)$$

To connect $\mathcal{E}_n(y)$ with $\hat{C}_n$, we then construct an iterative sequence $\{\tilde{C}_{n,j}(y)\}_{j \geq 1}$, analogous to (2.2), by setting $\tilde{C}_{n,0}(y) := \hat{C}_{n,0}$ (so that the oracle sequence has the same initial value as the original sequence $\hat{C}_{n,j}$) and recursively define, for $j \geq 1$,

$$\begin{aligned} \tilde{B}_{n,j-1}(y) &:= \sqrt{r_n T^{-1} \tilde{C}_{n,j-1}(y)}, \\ \tilde{C}_{n,j}(y) &:= \sum_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq \tilde{B}_{n,j-1}(y)\}}. \end{aligned} \quad (3.4)$$

Though the variables $\tilde{C}_{n,j}(y_n)$ and $\mathcal{E}_n(y)$ are not feasible estimators themselves, their asymptotic behavior in fact completely determines that of $\hat{C}_n$ provided the auxiliary sequence y_n tends to 0 at an appropriate rate. In our arguments, we demonstrate that

$$\tilde{C}_{n,n+1}(y_n) \leq \hat{C}_n \leq \mathcal{E}_n(y_n) + R_n,$$

for an appropriate asymptotically negligible remainder R_n . The above relation allows us to analyze $\hat{C}_n$ in terms of the array of oracle iterates $\{\tilde{C}_{n,j}(y_n)\}_{j \geq 1}$ and the oracle itself $\mathcal{E}_n(y_n)$. We then demonstrate that $\{\tilde{C}_{n,j}(y_n)\}_{j \geq 1}$ are all asymptotically equivalent to the oracle $\mathcal{E}_n(y_n)$ (Proposition A.1); effectively reducing the problem to the analysis of $\mathcal{E}_n(y_n)$, which is considerably simpler.

We now proceed to describe the class of initial estimators we consider in our procedure. Apart from some mild regularity conditions, they are required only to be consistent for C_T when the underlying process is continuous, allowing for a great deal of flexibility in the choice of initialization. More specifically, for a generic process Y , let

$$\hat{C}_{n,0}(Y) = \sum_{i=1}^{n-d+1} F(\Delta_i^n Y, \dots, \Delta_{i+d-1}^n Y), \quad (3.5)$$

where $F : \mathbb{R}^d \rightarrow [0, \infty)$ satisfies, for some $\delta_0 \in (0, 2]$ and for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$ with $\|\mathbf{y}\| \vee \|\mathbf{x}\| \leq 1$,

$$F(\mathbf{x}) \leq K \|\mathbf{x}\|_\infty^2, \quad (3.6)$$

$$|F(\mathbf{x} + \mathbf{y}) - F(\mathbf{x})| \leq K(\|\mathbf{y}\|_\infty \wedge \|\mathbf{x}\|_\infty)^{\delta_0} (\|\mathbf{y}\|_\infty^{2-\delta_0} + \|\mathbf{x}\|_\infty^{2-\delta_0}), \quad (3.7)$$

for some $K < \infty$. Above, $\|\mathbf{x}\|_\infty = \max_{1 \leq i \leq d} |x_i|$. An initial estimate $\hat{C}_{n,0}$ is said to belong to class $\mathcal{C}$ if $\hat{C}_{n,0} = \hat{C}_{n,0}(X)$, where $\hat{C}_{n,0}(\cdot)$ is given by (3.5), and satisfies

$$\hat{C}_{n,0}(\sigma \cdot W) \xrightarrow{P} \int_0^T \sigma_s^2 ds, \quad (3.8)$$

where $(\sigma \cdot W)_t := \int_0^t \sigma_s dW_s$.

Remark 3.1. The class $\mathcal{C}$ includes, for instance, the ordinary realized variance (i.e., $F(x) = x^2$), multipowers of the type $F(x_1, \dots, x_d) \propto \prod_{i=1}^d |x_i|^{r_i}$ with $r_i \geq 0$ and $r_1 + \dots + r_d = 2$, and the nearest-neighbor truncation estimators of [4], among other possibilities.

We now state our first main result.

Theorem 3.2. *Suppose $\hat{C}_{n,0}$ belongs to class $\mathcal{C}$, and that the sequence $r_n \rightarrow 0$ satisfies*

$$r_n \gg h_n \log(1/h_n). \quad (3.9)$$

Then, the following assertions hold:

(i) *If, further $r_n \ll (h \log(1/h))^{\frac{\alpha}{2}}$, then, as $n \rightarrow \infty$,*

$$\hat{C}_n \xrightarrow{P} C_T.$$

(ii) *If $\alpha \in (0, 1)$ and further $r_n \ll h_n^{\frac{\alpha+1}{2}} (\log(1/h))^{\frac{\alpha}{2}}$, then, as $n \rightarrow \infty$,*

$$\frac{1}{\sqrt{h_n}} (\hat{C}_n - C_T) \xrightarrow{st} \mathcal{N}\left(0, 2 \int_0^T \sigma_s^4 ds\right).$$

(iii) *Suppose $\alpha \in (1, 2)$ and $\gamma_t \equiv 1$. Then, as $n \rightarrow \infty$, for any choice of r_n satisfying (3.9),*

$$\frac{1}{\sqrt{h_n}} (\hat{C}_n - C_T) \xrightarrow{P} \infty.$$

Observe that the upper bounds on r_n in (i)–(ii) above depend on the jump activity index α and become more restrictive as α increases. Bearing this in mind, we make the following remarks.

Remark 3.3. A common choice in the literature for both benchmarking estimation performance and applications is $\sqrt{r_n} = 4h_n^{0.49}$ (e.g., [13, 24, 29]), which leads to efficient CLTs across nearly the entire range $\alpha < 1$; Section 4 compares the performance of this threshold choice for TRV against our iterative methods. Though the hypotheses of Theorem 3.2 require $r_n \gg h_n \log(1/h_n)$ in general, under the idealized assumption of constant volatility, our proofs demonstrate that r_n can be chosen of the form $r_n = 2(1 + \eta)h_n \log(1/h_n)$, for any $\eta > 0$ (cf. (1.3)). In general, the common choice $\sqrt{r_n} = 4h_n^{0.49}$ seems to also work reasonably well also for the iterative estimator $\hat{C}_n$; for further discussion of selection of the threshold rate r_n and initializations in practice, see Section 4.

Remark 3.4. When $\alpha \in (1, 2)$, a slowly-decaying bias term renders convergence rates of order $n^{-1/2}$ impossible for TRV itself under deterministic thresholding (c.f. [32]), which is reflected for $\hat{C}_n$ in case (iii) of Theorem 3.2. For these values of α , under deterministic thresholds for TRV, specialized debiasing techniques are required to achieve the optimal $n^{-1/2}$ rate (c.f. [9]). See also [24, 25] for other efficient methods based on empirical characteristic functions (and again deterministic tuning parameters). We leave the extension of these debiasing techniques to data-driven parameter tuning as a topic for future research.

3.2 Time-varying thresholds

For the best possible finite-sample performance, heuristically one should set the threshold ε as *small as possible* – to remove as many jumps as possible – but allow it to remain large enough so that a sufficient number increments remain to ultimately yield efficient estimates of C_T . From this perspective, the asymptotic lower bound on the rate r_n given in the hypotheses of

Theorem 3.2 may appear unsatisfactory, as it precludes rates as fast as the optimal threshold $\varepsilon_n^* \sim \sqrt{2\sigma^2 h_n \log(1/h_n)}$ in the constant volatility case. It is natural to suspect that faster rates may be possible for potential improvement in $\hat{C}_n$. However, the next result shows this is not true, in general.

Proposition 3.5. *For a given semimartingale Y , let $\text{TRV}_n(\varepsilon; Y) = \sum_{i=1}^n (\Delta_i^n Y)^2 \mathbf{1}_{\{|\Delta_i^n Y| \leq \varepsilon\}}$, and let $C_T(Y)$ denote its predictable quadratic variation. For a given $c_0 > 0$, define the threshold*

$$\vartheta_n := \sqrt{c_0 C_T(Y) h_n \log(1/h_n)}.$$

Then, there exists a semimartingale Y' such that $\text{TRV}_n(\vartheta_n; Y') \xrightarrow{P} C_T(Y')$, but

$$\sqrt{n}(\text{TRV}_n(\vartheta_n; Y') - C_T(Y')) \xrightarrow{P} -\infty.$$

Note that the estimator $\hat{C}_n$ defined in (2.4) is a TRV with threshold $\varepsilon_n = (c_0 \hat{C}_n h_n \log(1/h_n))^{1/2}$, which is approximately equal to ϑ_n if $\hat{C}_n$ remains a consistent estimator under this threshold choice. In that case, the above result suggests that $\hat{C}_n$ will not be rate-efficient in general with the threshold rate $r_n = c_0 h_n \log(1/h_n)$ and may remove too many increments even if jumps are completely absent from the process X . In particular, the proof of Proposition 3.5 illustrates that efficiency losses can result from volatility paths that exhibit significant jumps. A natural way to remedy this is to consider localized thresholds that adapt to the volatility level. In this way, thresholds corresponding to periods of high volatility are increased, and conversely, thresholds for periods of low volatility are decreased, so as to prevent efficiency losses that might otherwise occur with uniform thresholding. This is the central motivation behind our second class of estimators, which utilize spot volatility estimates to locally tune the threshold.

To this end, for a given even integer $k_n \leq n$ and $B > 0$, we define

$$\hat{\sigma}_n^2(i; B) := \frac{1}{h_n k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} (\Delta_\ell^n X)^2 \mathbf{1}_{\{|\Delta_\ell^n X| \leq B\}}, \quad \ell = 1, \dots, n,$$

where we set $\Delta_\ell^n X = 0$ if $\ell \leq 0$ or $\ell > n$. The above estimator is a type of kernel-based estimator of the spot volatility $\sigma_{t_i}^2$, as defined in [14, 28], with kernel function $K(x) = \frac{1}{2} \mathbf{1}_{[-1, 1]}$ and bandwidth $b_n = k_n h_n$ (see [23] for the asymptotic theory of the estimator in the case of one-sided uniform kernels $K(x) = \mathbf{1}_{[0, 1]}$ and [16, 21] for general kernels). Our second thresholding scheme for the localized thresholding estimator $\hat{C}_n^*$ then proceeds as follows:

1. First, we begin with some initial local volatility estimates $\hat{c}_{n,0}(i)$, $i = 1, \dots, n$, ideally free of tuning parameters. Natural choices include $\hat{c}_{n,0}(i) = \frac{1}{h_n k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} (\Delta_\ell^n X)^2$ or the analogous local BPV estimator $\hat{c}_{n,0}(i) = \frac{1}{h_n k_n} \frac{\pi}{2} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} |\Delta_\ell^n X| |\Delta_{\ell+1}^n X|$ (see Theorem 3.6 below for precise conditions on $\hat{c}_{n,0}(i)$).
2. Next, for each $i = 1, \dots, n$ and a given deterministic rate sequence $r_n^* \rightarrow 0$, we define a local threshold for the i -th increment of X . With regards to (1.4), each of these local thresholds can be viewed as a fixed-point iteration for one of the n stochastic maps $\xi \mapsto \hat{\sigma}_n^2(i; \sqrt{r_n^* \xi})$, $i = 1, \dots, n$. More specifically, for $j \geq 1$, we iteratively define:

$$\begin{aligned} B_{n,j-1}^*(i) &:= \sqrt{r_n^* \hat{c}_{n,j-1}^*(i)}, \\ \hat{c}_{n,j}^*(i) &:= \hat{\sigma}_n^2(i, B_{n,j-1}^*(i)). \end{aligned} \tag{3.10}$$

We then set

$$B_n^*(i) := B_{n,j_n^*}^*(i), \quad i = 1, \dots, n, \tag{3.11}$$

where $j_n^* := \min \{j \geq 0 : \hat{c}_{n,j}^*(i) = \hat{c}_{n,j+\ell}^*(i) \text{ for all } 1 \leq i \leq n, \ell \geq 0\}$.

3. Finally, we define the localized threshold estimator

$$\hat{C}_n^* := \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_n^*(i)\}}. \quad (3.12)$$

Let us now introduce the class of initial estimates $\mathcal{C}^{\text{spot}}$ for time-varying thresholds, which is essentially a localized analog of the class $\mathcal{C}$ of initial estimates defined in Section 3.1. To this end, for a generic process Y , define

$$\hat{c}_{n,0}(Y; i) = \frac{1}{h_n k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} F(\Delta_\ell^n Y, \dots, \Delta_{\ell+d-1}^n Y), \quad i = 1, \dots, n, \quad (3.13)$$

where $F : \mathbb{R}^d \rightarrow [0, \infty)$, and for convenience we set $F(\Delta_i^n Y, \dots, \Delta_{i+d-1}^n Y) = 0$ if $i \leq 0$ or $i + d - 1 > n$. We say the initializing threshold constants $\hat{c}_{n,0}(i)$ belong to the class $\mathcal{C}^{\text{spot}}$ if $\hat{c}_{n,0}(i) = \hat{c}_{n,0}(X; i)$, $i = 1, \dots, n$, where $\hat{c}_{n,0}(\cdot; i)$ are of the form (3.13) and satisfy

$$\max_{1 \leq i \leq n} \left| \hat{c}_{n,0}(\sigma \cdot W; i) - \frac{1}{h_n k_n} \int_{h_n(i-k_n/2+1)}^{h_n(i+k_n/2)} \sigma_t^2 dt \right| \xrightarrow{P} 0. \quad (3.14)$$

We are now in a position to state our second main result.

Theorem 3.6. *Let r_n^* satisfy*

$$\liminf_{n \rightarrow \infty} \frac{r_n^*}{2h_n \log(1/h_n)} > 1, \quad (3.15)$$

and suppose that $n^a \ll k_n \ll n$ for some $0 < a < 1$. Assume further that the initializing threshold constants $\hat{c}_{n,0}(i)$, $i = 1, \dots, n$, belong to the class $\mathcal{C}^{\text{spot}}$. Then, the following assertions hold:

(i) *If, further, $r_n^* \ll (h \log(1/h))^{\frac{\alpha}{2}}$, then, as $n \rightarrow \infty$,*

$$\hat{C}_n^* \xrightarrow{P} C_T.$$

(ii) *If $\alpha \in (0, 1)$ and further $r_n^* \ll h_n^{\frac{\alpha+1}{2}} (\log(1/h))^{\frac{\alpha}{2}}$, then, as $n \rightarrow \infty$,*

$$\frac{1}{\sqrt{h_n}} (\hat{C}_n^* - C_T) \xrightarrow{st} \mathcal{N}\left(0, 2 \int_0^T \sigma_s^4 ds\right).$$

(iii) *Suppose $\alpha \in (1, 2)$ and $\gamma_t \equiv 1$. Then, for any choice of r_n^* satisfying (3.15), as $n \rightarrow \infty$,*

$$\frac{1}{\sqrt{h_n}} (\hat{C}_n^* - C_T) \xrightarrow{P} \infty.$$

Remark 3.7. Similarly to the case for uniform thresholding, the assumptions on the admissible initial estimates (3.13) are relatively mild and require consistency in a uniform sense only in the continuous case. In particular, a localized version of realized variance, namely $\hat{c}_{n,0}(i, X) = \frac{1}{h_n k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} (\Delta_\ell^n X)^2$, or a localized bipower variation, namely $\hat{c}_{n,0}(i, X) = \frac{1}{h_n k_n} \frac{\pi}{2} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} |\Delta_\ell^n X| |\Delta_{\ell+1}^n X|$ both satisfy these assumptions. Indeed, the validity of (3.14) for the localized realized variance is shown in the proof of Lemma B.5 (see (B.30)), while for the localized bipower variation it follows along the same arguments as the proof of Proposition 3.3 in [36].

Statistical errors for spot volatility estimation are known to be substantially larger by comparison to the $O_P(n^{-1/2})$ -sized errors that occur in estimation of integrated volatility C_T (for instance, optimal choices of k_n in spot volatility estimation lead to errors of order $n^{-1/4}$; see, e.g., [21, 23]). Interestingly enough, the estimator $\hat{C}_n^*$ utilizes the comparatively noisier estimates of spot volatility in an auxiliary manner to lead to potentially improved estimates of C_T .

Remark 3.8. Recall that, to a first-order approximation, the threshold (1.3) is MSE-optimal under the assumption of constant volatility and finite jump activity. Though this is an idealized assumption and not expected to hold in many practical settings, it can serve as a reasonable local approximation at sufficiently high sampling frequencies. This intuition provides further support for our estimator $\hat{C}_n^*$, which roughly incorporates the optimal-type threshold (1.3) in a local manner. Indeed, our simulation study in Section 4 reflects this, showing that the resulting localized estimators exhibit superior finite-sample performance. Furthermore, we conjecture that for TRV based on time-varying thresholds $\varepsilon(i)$, $i = 1 \dots, n$, the threshold choice

$$\varepsilon_n(i) = \sqrt{2\bar{\sigma}_{n,i}^2 h_n \log(1/h_n)},$$

with $\bar{\sigma}_{n,i}^2 = \sup_{t \in [\frac{i-1}{n}, \frac{i}{n})} \sigma_t^2$, is MSE-optimal up to first order approximation under finite jump activity.

Remark 3.9. Feasible CLTs (for construction of confidence intervals) are possible with either $\hat{C}_n$ or $\hat{C}_n^*$. Indeed, minor extensions to our arguments show that under the rates r_n, r_n^* in part (ii) of Theorems 3.2 and 3.6, one has

$$\frac{\sqrt{n}(\hat{C}_n - C_T)}{\sqrt{2 \sum_{i=1}^n (\Delta_i^n X)^4 \mathbf{1}_{\{|\Delta_i^n X| \leq B_n\}}}} \xrightarrow{d} \mathcal{N}(0, 1), \quad \text{and} \quad \frac{\sqrt{n}(\hat{C}_n^* - C_T)}{\sqrt{2 \sum_{i=1}^n (\Delta_i^n X)^4 \mathbf{1}_{\{|\Delta_i^n X| \leq B_n^*(i)\}}}} \xrightarrow{d} \mathcal{N}(0, 1),$$

where the thresholds B_n and $B_n^*(i)$ are the same as for $\hat{C}_n$ and for $\hat{C}_n^*$ as described in Theorems 3.2 and 3.6, respectively.

Remark 3.10. Strictly speaking, the estimators $\hat{C}_n$ and $\hat{C}_n^*$ are not tuning free in the same way as, e.g., bipower variation is: $\hat{C}_n$ depends on r_n and $\hat{C}_n^*$ depends on r_n^* and k_n , which must still be chosen by the practitioner. However, the auxiliary sequences r_n, r_n^* and k_n are relatively data-insensitive and can be chosen based on sampling frequency alone. This stands in contrast with selecting the (full) parameter ε itself, which is highly data-sensitive, as demonstrated in Figure 1. This is further illustrated in the next section.

4 Monte Carlo Study

In this section, we compare the finite-sample performance of $\hat{C}_n$ and $\hat{C}_n^*$ against standard tuning approaches for TRV in the literature based on simulated data from the following stochastic volatility model:

$$X_t = 1 + \int_0^t \sigma_s dW_s + L_t + J_t,$$

$$\sigma_t^2 = \theta + \int_0^t \kappa(\theta - \sigma_s^2) ds + \xi \int_0^t \sigma_s dB_s.$$

Above, W and B are two correlated standard Brownian motions with covariation $d\langle W, B \rangle_t = \rho dt$, L is a CGMY Lévy process independent of W and B , and J is an inhomogeneous compound Poisson process independent of all other processes with intensity $\{\lambda(t)\}_{t \geq 0}$ and jump distribution $\varrho(dx)$.

Based on a 6.5 hour trading day and 252 trading days per year, we consider time horizons of $T \in \{\frac{1}{252}, \frac{5}{252}, \frac{1}{12}\}$, corresponding to 1 day, 1 week, and 1 month, respectively, at the 5-minute ($h_n = (\frac{1}{252})(\frac{1}{6.5})(\frac{5}{60})$) sampling frequency. For illustration, we examine five separate scenarios we now describe. Unless otherwise stated, for ease of comparison the parameters for $\{\sigma_t^2\}_{t \geq 0}$ are set as:

$$\kappa = 5, \quad \xi = 0.3, \quad \theta = (0.2)^2, \quad \rho = -0.5.$$

With these parameter choices, the annualized expected integrated variance is $(1/T)\mathbb{E}C_T = (0.2)^2$, and, in all settings, parameters are chosen so that the expected annualized realized volatility is approximately $\sqrt{(1/T)\mathbb{E}(\text{RV}_n)} \approx 0.275$ (Models 1,2,4,5, below) or 0.3 (Model 3), which are realistic for financial data.

- Model 1 (homogeneous jumps): for the infinite activity component $\{L_t\}_{t \geq 0}$, we choose

$$C_- = 0.148, \quad C_+ = 0.033, \quad G = 3.295, \quad M = 4.685, \quad Y = 0.917.$$

The parameters C_-, C_+, G, M are taken from estimates in [27] for a 1-year interval based on calibration from index options; here $Y = 0.917$ corresponds to the average of the reported estimates of Y^+ and Y^- in their model, namely $Y = \frac{Y^+ + Y^-}{2}$. For $\{J_t\}_{t \geq 0}$, we choose

$$\lambda(t) \equiv 252 \text{ (1 jump per day)}, \quad \varrho(dx) \sim \mathcal{N}(-0.005, 0.01^2).$$

- Model 2 (switching jump intensity): all settings are the same as in Model 1, except $\{J_t\}_{t \geq 0}$ has time-varying intensity:

$$\lambda(t) = (252)\vartheta(t), \quad \vartheta(t) = \begin{cases} 2, & \sigma_t^2 > \theta, \\ 0, & \sigma_t^2 < \theta. \end{cases}$$

- Model 3 (higher jump intensity): all settings are the same as Model 1, except $\{J_t\}_{t \geq 0}$ has a higher jump intensity:

$$\lambda(t) \equiv (1.5)(252) \text{ (1.5 jumps per day)}.$$

- Model 4 (finite jump activity): all settings are the same as Model 1, except we take $L_t \equiv 0$, and adjust $\lambda(t) \equiv (1.15)252$ to match the expected realized variance of Model 1.

- Model 5 (no jumps): All settings are the same as Model 1, except we take $J_t \equiv 0$, $L_t \equiv 0$, and adjust $\theta = (0.275)^2$ to match the expected realized variance of Model 1.

We compare 6 types of estimators based on TRV: two instances of standard approaches, and two instances each of the iterative estimator $\hat{C}_n$ and of the localized iterative estimator $\hat{C}_n^*$ with different types of initializations. Specifically, for $\hat{C}_n$ we use the following estimators as initializations:

$$\text{RV}_n = \sum_{i=1}^n (\Delta_i^n X)^2, \quad \text{BV}_n = \frac{\pi}{2} \sum_{i=2}^n |\Delta_{i-1}^n X| |\Delta_i^n X|.$$

For $\hat{C}_n^*$, we use their localized counterparts, denoted by

$$\hat{\sigma}_n^2(\ell) = \frac{1}{h_n k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} (\Delta_i^n X)^2, \quad \text{BV}_n^{\text{spot}}(\ell) = \frac{1}{h_n k_n} \frac{\pi}{2} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} |\Delta_{i-1}^n X| |\Delta_i^n X|.$$

We consider the following estimation procedures:

- (1) $\text{TRV}_n(\varepsilon_{0,n})$, where $\varepsilon_{0,n} = h_n^{0.49}$;
- (2) $\text{TRV}_n(\varepsilon_{1,n})$, where $\varepsilon_{1,n} = \sqrt{\text{BV}_n r_n}$, with $\sqrt{r_n} = 4h_n^{0.49}$ as used in [13, 24, 29].
- (3) $\hat{C}_n$, with initialization $\hat{C}_{n,0} = \text{RV}_n$ and r_n as in (2);
- (4) $\hat{C}_n$, with initialization $\hat{C}_{n,0} = \text{BV}_n$ and r_n as in item (2);
- (5a,b) $\hat{C}_n^*$, with $r_n^* = 2h_n(\log(1/h_n) - \log \log(1/h_n))$, $k_n = h_n^{-0.5}$ (5a) or $k_n = h_n^{-0.6}$ (5b), and initialization $\hat{c}_{n,0}(i) = \hat{\sigma}_n^2(i)$;
- (6a,b) $\hat{C}_n^*$, with $r_n^* = 2h_n(\log(1/h_n) - \log \log(1/h_n))$, $k_n = h_n^{-0.5}$ (6a) or $k_n = h_n^{-0.6}$ (6b), and initialization $\hat{c}_{n,0}(i) = \text{BV}_n^{\text{spot}}(i)$.

Model 1 (homogeneous jumps)									
	$T = 1/252$ (1 day)			$T = 5/252$ (1 wk.)			$T = 1/12$ (1 mo.)		
	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$
(1) TRV	90.4783	149.1879	2.7355	90.2462	67.2049	8.7914	92.9712	39.0118	31.4622
(2) TRV	5.3009	20.8770	0.3421	4.8734	8.7488	0.8040	4.7463	4.4099	2.2017
(3) $\hat{C}_n$	5.8617	22.3127	0.3665	3.9965	8.4482	0.7505	3.6931	4.2304	1.9256
(4) $\hat{C}_n$	4.7353	20.1997	0.3301	3.9688	8.4253	0.7478	3.6931	4.2304	1.9256
(5a) $\hat{C}_n^*$	3.3682	19.0994	0.3087	2.9690	8.1442	0.6956	2.8222	4.1195	1.7050
(5b) $\hat{C}_n^*$	3.3297	19.0585	0.3079	2.5464	8.0069	0.6740	2.3890	4.0001	1.5881
(6a) $\hat{C}_n^*$	2.8296	18.5914	0.2994	2.7570	8.0723	0.6843	2.6449	4.0711	1.6546
(6b) $\hat{C}_n^*$	2.8294	18.5915	0.2989	2.4985	8.1560	0.6714	2.3621	4.2415	1.5815

Model 2 (switching jump intensity)									
	$T = 1/252$ (1 day)			$T = 5/252$ (1 wk.)			$T = 1/12$ (1 mo.)		
	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$
(1) TRV	98.4132	190.3505	3.5307	89.9458	95.9290	11.3156	83.7461	62.7170	40.2647
(2) TRV	6.4172	23.0589	0.3911	5.1908	10.1623	0.9698	4.8893	6.0500	3.0746
(3) $\hat{C}_n$	6.6465	29.6733	0.5108	3.7809	9.1587	0.8318	3.4781	5.0420	2.3673
(4) $\hat{C}_n$	5.1206	20.8151	0.3475	3.7568	9.1281	0.8278	3.4772	5.0429	2.3673
(5a) $\hat{C}_n^*$	3.6255	19.4813	0.3193	2.8709	8.6546	0.7561	3.0375	4.7020	2.1106
(5b) $\hat{C}_n^*$	3.6256	19.5659	0.3207	2.2625	8.3286	0.7137	2.4621	4.4255	1.8864
(6a) $\hat{C}_n^*$	3.1309	18.7966	0.3062	2.6498	8.5198	0.7385	2.8564	4.6173	2.0377
(6b) $\hat{C}_n^*$	3.1190	18.7985	0.3063	2.2198	8.6419	0.7114	2.4332	4.8536	1.8801

Model 3 (higher jump intensity)									
	$T = 1/252$ (1 day)			$T = 5/252$ (1 wk.)			$T = 1/12$ (1 mo.)		
	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$
(1) TRV	121.7856	167.8171	3.2809	133.4801	81.8533	12.1529	134.3111	49.2704	44.6016
(2) TRV	7.8602	24.2170	0.4053	7.3329	10.1078	0.9921	7.1037	4.8398	2.9336
(3) $\hat{C}_n$	9.1251	36.1741	0.5982	5.2370	9.2686	0.8526	4.8782	4.5093	2.3053
(4) $\hat{C}_n$	6.3337	22.0729	0.3656	5.2050	9.2380	0.8490	4.8782	4.5093	2.3053
(5a) $\hat{C}_n^*$	4.0298	20.3463	0.3307	3.7660	8.7422	0.7627	3.6815	4.2953	1.9582
(5b) $\hat{C}_n^*$	4.0138	20.2726	0.3295	3.0898	8.4432	0.7199	3.0808	4.1801	1.7936
(6a) $\hat{C}_n^*$	3.3382	19.4716	0.3145	3.5166	8.6317	0.7458	3.4983	4.2390	1.8997
(6b) $\hat{C}_n^*$	3.5848	19.5251	0.3140	3.0404	8.6777	0.7159	3.0138	4.5086	1.7595

Table 1: Estimation performance of $\hat{C}_n$, $\hat{C}_n^*$, and standard tuning approaches for TRV in Models 1-3; reported values are based on $m = 5000$ realizations in each model at the 5-minute sampling frequency.

We simulate $m = 5000$ paths for each Model 1-5. Denoting by $\hat{C}$ one of the estimators in (1)-(6), on the j -th realization we compute the estimator value, $\hat{C}_j$, the corresponding true integrated volatility, $C_{T,j}$, and report

- The mean relative error (in %): $100(\frac{1}{m} \sum_{j=1}^m e_j)$, where $e_j = \frac{\hat{C}_j - C_{T,j}}{C_{T,j}}$;
- The standard deviation of the relative error (in %): $100\sqrt{\frac{1}{m} \sum_{j=1}^m (e_j - \bar{e})^2}$;
- $\sqrt{\text{MSE}} = \sqrt{\frac{1}{m} \sum_{j=1}^m (\hat{C}_j - C_{T,j})^2}$.

The results are displayed in Tables 1-2; the smallest bias and MSE for each time horizon are shown in bold.

In general, we see that when jumps are present (Models 1-4), both the iterative estimator $\hat{C}_n$ and localized iterative estimator $\hat{C}_n^*$ can outperform the standard tuning choice (2) for TRV both in terms of relative error and MSE by significant margins, with reductions in bias often by 50% or more by comparison to (2) and reductions in $\sqrt{\text{MSE}}$ as high as 40%. As anticipated, deterministic tuning (1) performs rather poorly by comparison to approaches (2)-(6) on all time horizons, and although the (non-iterative) bipower-tuned TRV in (2) leads to a substantial improvement over (1), it is uniformly outperformed by (4)-(6) on all time horizons considered and also outperformed by (3) except on daily time horizons.

In general, the localized estimators (6a,6b) have the largest relative performance gains compared to standard procedures (1)-(2) over longer time horizons, which is somewhat expected, ranging from 13%-23% reduction in $\sqrt{\text{MSE}}$ at daily horizons to 28%-40% reduction in $\sqrt{\text{MSE}}$ at monthly horizons compared to (2). Also, iterative approaches with jump-robust initializations (4,6a,6b) generally have improved performance compared to those without jump-robust initializations (3,5a,5b). Furthermore, for the localized estimators, the choice $k_n = h_n^{-0.6}$ (5b,6b) tends to lead to improvement relative to the choice $k_n = h_n^{-0.5}$ (5a,6a) over longer time horizons. The best performance in terms of both relative error and MSE is typically achieved by (6b).

Model 4 (finite jump activity)									
	$T = 1/252$ (1 day)			$T = 5/252$ (1 wk.)			$T = 1/12$ (1 mo.)		
	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$
(1) TRV	95.5145	159.3735	2.9314	89.8228	66.1216	8.6889	93.9742	40.3833	31.7663
(2) TRV	4.3921	21.5406	0.3515	3.5605	8.7916	0.7641	3.5693	4.2456	1.9309
(3) $\hat{C}_n$	4.9183	23.2150	0.3783	2.6648	8.4874	0.7164	2.5184	4.1242	1.6911
(4) $\hat{C}_n$	3.5558	20.1041	0.3255	2.6388	8.4716	0.7145	2.5179	4.1244	1.6911
(5a) $\hat{C}_n^*$	1.9505	18.3226	0.2933	1.5714	8.0859	0.6617	1.7152	3.9780	1.5053
(5b) $\hat{C}_n^*$	1.9454	18.2932	0.2928	1.1940	7.9727	0.6475	1.2999	3.8990	1.4207
(6a) $\hat{C}_n^*$	1.5691	18.0673	0.2886	1.3933	8.0169	0.6527	1.5442	3.9375	1.4680
(6b) $\hat{C}_n^*$	1.5623	18.0679	0.2880	1.1613	8.0538	0.6452	1.2761	4.0324	1.4166

Model 5 (no jumps)									
	$T = 1/252$ (1 day)			$T = 5/252$ (1 wk.)			$T = 1/12$ (1 mo.)		
	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$	rel. err (%)	sd(rel. err)	$\sqrt{\text{MSE}} \times 10^4$
(1) TRV	0.0917	15.8450	0.5671	-0.1081	7.0844	1.2695	-0.0536	3.4524	2.6256
(2) TRV	0.0504	15.8488	0.5671	-0.1308	7.0946	1.2717	-0.0936	3.4619	2.6346
(3) $\hat{C}_n$	0.0857	15.8440	0.5670	-0.1274	7.0909	1.2708	-0.0930	3.4617	2.6343
(4) $\hat{C}_n$	0.0632	15.8549	0.5673	-0.1311	7.0941	1.2715	-0.0940	3.4616	2.6344
(5a) $\hat{C}_n^*$	0.0306	15.8651	0.5677	-0.1920	7.1099	1.2744	-0.1748	3.4660	2.6394
(5b) $\hat{C}_n^*$	0.0344	15.8649	0.5677	-0.2343	7.1112	1.2750	-0.2063	3.4639	2.6388
(6a) $\hat{C}_n^*$	-0.0636	15.8999	0.5689	-0.2669	7.1294	1.2782	-0.2492	3.4651	2.6424
(6b) $\hat{C}_n^*$	-0.0668	15.8999	0.5691	-0.2696	7.1293	1.2766	-0.2432	3.4656	2.6425

Table 2: Estimation performance of $\hat{C}_n$, $\hat{C}_n^*$, and standard tuning approaches for TRV in the finite jump activity setting of Model 4 and the jump-free setting of Model 5; reported values are based on $m = 5000$ realizations at the 5-minute sampling frequency.

Comparing performance across Models 1-4, we see that all iterative approaches (3)-(6) are generally more robust against increased levels of jump activity as well as time-varying jump behavior compared to (1)-(2). In both Models 2 and 3, on longer time horizons, the performance advantage of the localized estimators over uniform approaches is typically larger by comparison to the performance advantage they have over uniform approaches in Model 1. Though all estimators (1)-(6) have better overall performance under finite jump activity (Model 4) relative to settings with infinite activity (Models 1-3), the iterative approaches still retain performance advantages over standard choices (1)-(2) even without an infinite activity component in the

model.

Turning to the jump-free case (Model 5), we note that all estimators perform very similarly in terms of both bias and MSE and are typically slightly negatively biased. Over weekly and monthly time horizons, the localized estimators (5-6) incur a very slight increase in bias (appx. 0.15%) compared to uniform thresholding (2), and the deterministic TRV has marginally smaller $\sqrt{\text{MSE}}$ compared to (2)-(6).

We note that although a slight increase in bias occurs in the localized estimators (5)-(6) in the absence of jumps, it is relatively small relative to the potential performance gain one may attain if jumps are present. Since jumps are generically expected in many types of data, for use in practice we recommend the localized estimator with jump-robust initialization and settings of (6b). However, if a simpler implementation is desired, or one wants to avoid the potential marginal additional bias when jumps are absent, method (4) is a reasonable alternative. We remark that in any case, these choices (4,6b) in the presence of jumps can significantly outperform the common choice in the literature (2).

Number of iterations until stabilization ($T = 1/12$)							
	1	2	3	4	5	6	≥ 7
$j_n(\hat{C}_n)$	1.13%	27.21%	54.13%	15.69%	1.65%	0.17%	0.02%
$j_n^*(\hat{C}_n^*)$	1.06%	27.15%	55.78%	14.45%	1.45%	0.10%	0.01%

Table 3: Empirical distribution of j_n and j_n^* at the $T = 1/12$ (1 month) time horizon. Reported values reflect the empirical percentages of the aggregated counts of iterations until stabilization across all computed values of each estimator (for $\hat{C}_n$, across both settings (3) and (4); for $\hat{C}_n^*$, across both (5ab) and (6ab)) and across Models 1-4.

Regarding computational considerations, in Table 3 we report the empirical distribution of the number of iterations required for stabilization for both the localized thresholding and uniform thresholding approaches (i.e., j_n , as in (2.3), and j_n^* as in (3.11)) across all models with jumps (Models 1-4) on 1-month time horizons. Both $\hat{C}_n$ and $\hat{C}_n^*$ stabilize rather quickly, with roughly 98% of all estimates stabilizing in 4 or fewer iterations, and the global and local thresholding approaches take roughly the same number of iterations. Though not included in Table 3, in the jump-free case (Model 5), all observed instances of estimators stabilized in 3 or fewer iterations, with the vast majority taking 1 or 2; also, shorter time horizons typically required fewer iterations to stabilize in all settings.

Unreported simulation studies suggest localized estimators can have further performance gains relative to uniform thresholding approaches when the time horizon is extended or when additional inhomogeneities are incorporated into the model such as volatility jumps. Generally performance improvement of $\hat{C}_n$ and $\hat{C}_n^*$ relative to standard-type TRV tuning (1) and (2) becomes more dramatic as the overall proportion of jump variation increases relative to the quadratic variation of X , or when the activity of either jump component (L or J) is increased, and substantive performance gains are typically observed provided at least one of these components is present. We also remark that at daily horizons, with relatively small sample size ($n = 78$) there is little difference between uniform thresholding (3)-(4) and the localized thresholding (5)-(6), except for the rates r_n and r_n^* ; not included in this study is a detailed examination of the optimal choice of k_n , which could be of future interest, though $k_n = h_n^{-0.6}$ seems to reasonably well in most scenarios.

A Proofs

Throughout the proofs, we often omit the subscript n in h_n and y_n , and K denotes a generic constant that may change from line to line. Without loss of generality, we assume that $T = 1$ (giving $h_n = \frac{1}{n}$), and for notational simplicity when dealing with boundary terms we set $\sigma_t := 0$ for $t \notin [0, 1]$. Also, for any process $(V_i)_{t \in [0, 1]}$, we set $\Delta_i^n V = 0$ whenever $i \leq 0$ or $i > n$.

By a standard localization argument, we may assume without loss of generality that b, σ, σ^{-1} , and γ are bounded above by a nonrandom constant, and that

$$\mathbb{E}(|\gamma_{t+s} - \gamma_t|^2 | \mathcal{F}_t) \leq Ks, \quad s, t \in (0, T].$$

We also collect some useful estimates below that are used throughout the appendix. Based on the decomposition of the process L in (3.1), we have the following as $n \rightarrow \infty$ (and thus as $y_n \rightarrow 0, h_n \rightarrow 0$):

$$\begin{aligned} \lambda_n(y) &\sim Ky^{-\alpha}, \quad \mathbb{P}(\Delta_i^n N(y) \neq 0) \sim Khy^{-\alpha}, \\ |\Delta_i^n b(y)| &\leq Kh(y^{1-\alpha} + 1). \end{aligned} \tag{A.1}$$

In our arguments, we also need oracle analogs of the estimator $\widehat{C}_n^*$ in (3.12). To construct them, we first define

$$\bar{\sigma}_n^2(\ell; B; y) := \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B, \Delta_i^n N(y)=0, \Delta_i^n N'=0\}}, \quad \ell = 1, \dots, n.$$

We then define an auxiliary sequence $\tilde{c}_{n,j}(i; y)$ through a fixed-point iteration for the function $\xi \mapsto \bar{\sigma}_n^2(\ell; \sqrt{r_n^* \xi}; y)$ as follows: we set

$$\tilde{c}_{n,1}(i; y) := \bar{\sigma}_n^2(\ell; B_{n,0}^*(i); y), \quad i = 1, \dots, n,$$

where $B_{n,0}^*(i)$ is defined as (3.10) with the initializing threshold constants $\widehat{c}_{n,0}(i)$, $i = 1, \dots, n$, being *the same* as those for $\widehat{C}_n^*$. Next, we define oracle counterparts of the iterative localized estimates $\widehat{c}_{n,j}(i)$ in (3.10), for every $1 \leq i \leq n$ and $j \geq 2$ by setting

$$\begin{aligned} \tilde{B}_{n,j-1}^*(i; y) &:= \sqrt{r_n^* \tilde{c}_{n,j-1}(i; y)}, \\ \tilde{c}_{n,j}(i; y) &:= \bar{\sigma}_n^2(i; \tilde{B}_{n,j-1}^*(i; y); y). \end{aligned} \tag{A.2}$$

We finally define oracle counterparts of the time-varying threshold iterates $\{\tilde{C}_{n,j}^*, j \geq 1\}$ by setting

$$\tilde{C}_{n,j}^*(y) = \sum_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq \tilde{B}_{n,j}^*(i; y)\}}, \quad j \geq 1, \tag{A.3}$$

where $\mathcal{I}_n(y)$ is given in (3.2). The above oracle sequence plays an analogous role to the oracle sequence $\tilde{C}_{n,j}(y)$ of (3.4), but for the case of time-varying thresholds.

We first establish the following key intermediate result, which shows that the iterative sequences $\tilde{C}_{n,j}(y)$, $\tilde{C}_{n,j}^*(y)$ (defined in (3.4) and (A.3), respectively) are asymptotically equivalent to the oracle version of TRV, $\mathcal{C}_n(y)$, defined in (3.3), provided $y \rightarrow 0$ at an appropriate rate. Below, we use the notation $\bar{\sigma}^2 = \sup_{0 \leq s \leq 1} \sigma_s^2$.

Proposition A.1. *Assume the initial estimates $\widehat{C}_{n,0}$ and $\widehat{c}_{n,0}(i)$ belong to the classes $\mathcal{C}$ and $\mathcal{C}^{spot}$, respectively. Let $y \rightarrow 0$ so that, for some $\delta \in (0, \frac{1}{\alpha} \wedge 1)$,*

$$h^{(\frac{1}{\alpha} \wedge 1) - \delta} \ll y \ll (h \log n)^{\frac{1}{2}}. \tag{A.4}$$

Further suppose that $n^a \ll k_n \ll n$ for some $0 < a < 1$, and

$$\liminf_{n \rightarrow \infty} \frac{r_n}{2h \log n} > \frac{\bar{\sigma}^2}{\int_0^1 \sigma_s^2 ds} \quad \text{and} \quad \liminf_{n \rightarrow \infty} \frac{r_n^*}{2h \log n} > 1. \quad (\text{A.5})$$

Then, with probability tending to 1,

$$\mathcal{E}_n(y) = \tilde{C}_{n,j}(y) = \tilde{C}_{n,j}^*(y), \quad \text{for all } j \geq 1.$$

Proof. Recall the notation $\mathcal{I}_n(y)$ as in (3.2), and let

$$\tilde{\mathbf{c}}_{n,j}(y) = (\tilde{c}_{n,j}(1; y), \dots, \tilde{c}_{n,j}(n; y)).$$

Also, by analogy to $\mathcal{E}_n(y)$, define

$$c_n(i; y) = \frac{n}{k_n} \sum_{\ell=i-k_n/2}^{i+k_n/2} (\Delta_\ell^n X)^2 \mathbf{1}_{\{\Delta_\ell^n N(y)=0, \Delta_\ell^n N'=0\}}, \quad i = 1, \dots, n. \quad (\text{A.6})$$

First we claim it suffices to show that, with probability tending to 1,

$$|\Delta_i^n X|^2 \leq r_n \hat{C}_{n,0}, \quad \text{for all } i \in \mathcal{I}_n(y), \quad (\text{A.7})$$

$$|\Delta_i^n X|^2 \leq r_n \mathcal{E}_n(y), \quad \text{for all } i \in \mathcal{I}_n(y), \quad (\text{A.8})$$

$$|\Delta_i^n X|^2 \leq r_n^* \hat{c}_{n,0}(i), \quad \text{for all } i \in \mathcal{I}_n(y), \quad (\text{A.9})$$

$$|\Delta_i^n X|^2 \leq r_n^* c_n(i; y), \quad \text{for all } i \in \mathcal{I}_n(y). \quad (\text{A.10})$$

Indeed, recalling $B_{n,0} = \sqrt{r_n \hat{C}_{n,0}}$ and the definitions of $\mathcal{E}_n(y)$ and $\tilde{C}_{n,1}(y)$ as in (3.3) and (3.4), respectively, expression (A.7) implies that $\tilde{C}_{n,1}(y) = \mathcal{E}_n(y)$, while (A.8) implies that $\tilde{C}_{n,j}(y) = \tilde{C}_{n,1}(y)$, for any $j \geq 2$. In a similar fashion, (A.9) implies that $\tilde{c}_{n,1}(i; y) = c_n(i; y)$ for all i . From (A.2) this immediately gives $\tilde{B}_{n,1}^*(i; y) = \sqrt{r_n^* c_n(i; y)}$ for all i , which, from (A.3) and using (A.10), implies that $\tilde{C}_{n,1}^*(y) = \mathcal{E}_n(y)$. Continuing, this gives $\tilde{c}_{n,2}(i; y) = c_n(i; y)$ for all i , and thus, $\tilde{B}_{n,2}^*(i; y) = \sqrt{r_n^* c_n(i; y)}$ for all i ; proceeding by induction, we conclude that $\tilde{C}_{n,j}^*(y) = \mathcal{E}_n(y)$ and $\tilde{\mathbf{c}}_{n,j}(y) = (c_n(1; y), \dots, c_n(n; y))$ for all $j \geq 1$.

We first establish (A.7) and (A.8). It suffices to show that for some small (nonrandom) $\eta > 0$, with probability tending to 1,

$$\frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2}{r_n \hat{C}_{n,0}} \leq 1 - \eta, \quad \text{and} \quad \frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2}{r_n \mathcal{E}_n(y)} \leq 1 - \eta. \quad (\text{A.11})$$

Note that, for each $i \in \mathcal{I}_n(y)$, $\Delta_i^n J = 0$, and for $H_t(y)$ as in the decomposition (3.1), we have $\int_{t_{i-1}}^{t_i} \gamma_t H_t(y) dt = 0$. So, denoting $(\sigma \cdot W)_t = \int_0^t \sigma_s dW_s$, if we define

$$\Delta_i^n \chi(y) := \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) + \Delta_i^n b_t(y), \quad (\text{A.12})$$

then we have $\Delta_i^n X - \Delta_i^n (\sigma \cdot W) = \Delta_i^n \chi(y)$ for all $i \in \mathcal{I}_n(y)$. To show (A.11), note that

$$\begin{aligned} \max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2 &\geq \max_{i \in \mathcal{I}_n(y)} \left((\Delta_i^n (\sigma \cdot W))^2 - |\Delta_i^n \chi(y)| (2|\Delta_i^n (\sigma \cdot W)| + |\Delta_i^n \chi(y)|) \right) \\ &\geq \max_{i \in \mathcal{I}_n(y)} (\Delta_i^n (\sigma \cdot W))^2 - \max_{i \in \mathcal{I}_n(y)} \left\{ |\Delta_i^n \chi(y)| (2|\Delta_i^n (\sigma \cdot W)| + |\Delta_i^n \chi(y)|) \right\}, \end{aligned}$$

and clearly,

$$\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2 \leq \max_{i \in \mathcal{I}_n(y)} (\Delta_i^n (\sigma \cdot W))^2 + \max_{i \in \mathcal{I}_n(y)} \left\{ |\Delta_i^n \chi(y)| (2|\Delta_i^n (\sigma \cdot W)| + |\Delta_i^n \chi(y)|) \right\}.$$

We first bound $\max_{i \in \mathcal{I}_n(y)} \{|\Delta_i^n \chi(y)| (2|\Delta_i^n(\sigma \cdot W)| + |\Delta_i^n \chi(y)|)\}$. We have, for some nonrandom $C > 0$,

$$\max_{1 \leq i \leq n} \frac{|\Delta_i^n b_t(y)|}{y} \leq C \left(\frac{h(y^{1-\alpha} + 1)}{y} \right) \rightarrow 0,$$

because in the case when $\alpha < 1$, both $y^{1-\alpha} \rightarrow 0$ and $h/y \ll h^\delta \rightarrow 0$, and when $\alpha > 1$, $hy^{-\alpha} \rightarrow 0$. Moreover, Lemma B.1 gives $\mathbb{P}(|\Delta_i^n M(y)| > k) \leq C(hy^{1-\alpha}k^{-1})^{\frac{k}{2y}}$ for every $k > 0$, and Lemma 2.1.5 in [23] gives

$$\mathbb{E}_{i-1} \left| \int_{t_{i-1}}^{t_i} (\gamma_t - \gamma_{t_{i-1}}) dM_t(y) \right|^2 \leq Ky^{2-\alpha} \mathbb{E}_{i-1} \int_{t_{i-1}}^{t_i} |\gamma_t - \gamma_{t_{i-1}}|^2 dt \leq Ky^{2-\alpha} h^2.$$

Thus, we obtain, for all large enough $\kappa > 0$,

$$\begin{aligned} \mathbb{P} \left(\left| \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right| > \kappa y \right) &\leq \mathbb{P} \left(\left| \int_{t_{i-1}}^{t_i} (\gamma_t - \gamma_{t_{i-1}}) dM_t(y) \right| > \kappa y/2 \right) + \mathbb{P}(|\gamma_{t_i}| |\Delta_i^n M(y)| > \kappa y/2) \\ &\leq K \left(y^{-2} \mathbb{E} \left| \int_{t_{i-1}}^{t_i} (\gamma_t - \gamma_{t_{i-1}}) dM_t(y) \right|^2 + (hy^{1-\alpha}y^{-1})^{\frac{\kappa y}{2y}} \right) \\ &\leq K \left(h^2 y^{-\alpha} + (hy^{-\alpha})^{\frac{\kappa}{2}} \right) \\ &\leq K \left(h^{1+\delta\alpha} + (h^{\delta\alpha})^{\frac{\kappa}{2}} \right) = o(h). \end{aligned}$$

This gives, for all large n , and large enough $\kappa > 0$,

$$\begin{aligned} \mathbb{P} \left(\max_{1 \leq i \leq n} |\Delta_i^n \chi(y)| \geq \kappa y \right) &\leq \sum_{i=1}^n \mathbb{P}(|\Delta_i^n \chi(y)| \geq \kappa y) \\ &\leq \sum_{i=1}^n \left(\mathbb{P} \left(\left| \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right| \geq \kappa y/2 \right) + \mathbb{P} \left(\max_{1 \leq i \leq n} \frac{|\Delta_i^n b_t(y)|}{y} > \kappa/2 \right) \right) \\ &= \sum_{i=1}^n \mathbb{P} \left(\left| \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right| \geq \kappa y/2 \right) = o(1). \end{aligned}$$

In particular,

$$\max_{1 \leq i \leq n} |\Delta_i^n \chi(y)| = O_P(y). \quad (\text{A.13})$$

Thus,

$$\begin{aligned} &h^{-1} \left| \max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2 - \max_{i \in \mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2 \right| \\ &\leq h^{-1} \max_{i \in \mathcal{I}_n(y)} \left\{ |\Delta_i^n \chi(y)| (2|\Delta_i^n(\sigma \cdot W)| + |\Delta_i^n \chi(y)|) \right\} \\ &\leq h^{-1} \max_{1 \leq i \leq n} |\Delta_i^n \chi(y)| \left(2 \max_{1 \leq i \leq n} |\Delta_i^n(\sigma \cdot W)| + \max_{1 \leq i \leq n} |\Delta_i^n \chi(y)| \right) \\ &= h^{-1} O_P(y) \left(O(\sqrt{h \log(1/h)}) + O_P(y) \right) \\ &= O_P(yh^{-1/2}(\log n)^{1/2}) + O_P(h^{-1}y^2) \\ &= o_P(\log n), \end{aligned} \quad (\text{A.14})$$

where $\max_{1 \leq i \leq n} |\Delta_i^n(\sigma \cdot W)| = O(\sqrt{h \log(1/h)})$ holds as a consequence of Lemma B.6, and the last line holds since $y \ll (h \log n)^{1/2}$.

We are now in position to show (A.7)–(A.8). With $\underline{C}_n = \underline{C}_n(X)$ as defined in Proposition B.3 below, we have $\widehat{C}_{n,0} \geq \underline{C}_n$ and $\underline{C}_n \xrightarrow{P} \int_0^1 \sigma_s^2 ds$ as shown in Proposition B.3. Furthermore, condition (A.5) implies, for some small $\delta' > 0$, for large enough n , $\frac{r_n}{2h \log n} \geq (1 + \delta') \left(\frac{\bar{\sigma}^2}{\int_0^1 \sigma_s^2 ds} \right)$. Thus,

$$\begin{aligned} & \frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2}{r_n \widehat{C}_{n,0}} \\ & \leq \left(\frac{2}{2(1 + \delta')} \right) \left(\frac{\int_0^1 \sigma_s^2 ds}{\underline{C}_n} \right) \frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2}{2\bar{\sigma}^2 h \log n} \\ & \leq \left(\frac{1}{1 + \delta'} \right) \left(\frac{\int_0^1 \sigma_s^2 ds}{\underline{C}_n} \right) \left(\frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2}{2\bar{\sigma}^2 h \log n} + \left| \frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2 - \max_{i \in \mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2}{2\bar{\sigma}^2 h \log n} \right| \right) \\ & = \left(\frac{1}{1 + \delta'} \right) \left(1 + o_P(1) \right) \left(\frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2}{2\bar{\sigma}^2 h \log n} + o_P(1) \right). \end{aligned} \quad (\text{A.15})$$

Applying Lemma B.6, we have, for every $\eta' > 0$, $\frac{\max_{i \in \mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2}{2\bar{\sigma}^2 h \log n} < 1 + \eta'$ with probability tending to 1. Thus, taking $\eta' > 0$ small enough, the right-hand side in (A.15) is strictly less than $1 - \eta$ with probability tending to one for some small $\eta > 0$, and we obtain the first statement in (A.11); the statement (A.7) then follows. The second statement in (A.11) is proved along the same lines as in (A.15), replacing both $\widehat{C}_{n,0}$ and $\underline{C}_n$ with $\mathcal{C}_n(y)$ and applying Proposition B.2-(i), giving (A.8).

We now show (A.9)–(A.10). Condition (A.5) implies, for some small $\delta' > 0$, for large enough n , $\frac{r_n}{2h \log n} \geq (1 + \delta')$. Recalling that $B_{n,0}^* = \sqrt{r_n^* \widehat{C}_{n,0}(i)}$, and with $\underline{c}_n(i) = \underline{c}_n(X; i)$ as in Proposition B.4, we have $\widehat{C}_{n,0}(i) \geq \underline{c}_n(i)$ and thus

$$\begin{aligned} & \max_{i \in \mathcal{I}_n(y)} \frac{(\Delta_i^n X)^2}{r_n^* \widehat{C}_{n,0}(i)} \\ & \leq \left(\frac{1}{1 + \delta'} \right) \max_{1 \leq i \leq n} \left(\frac{\sup_{t \in [\frac{i-k_n/2}{n}, \frac{i+k_n/2}{n}] \sigma_t^2}{\underline{c}_n(i)} \right) \max_{i \in \mathcal{I}_n(y)} \frac{(\Delta_i^n X)^2}{2 \left(\sup_{t \in [\frac{i-k_n/2}{n}, \frac{i+k_n/2}{n}] \sigma_t^2} \right) h \log n} \\ & \leq \left(\frac{1}{1 + \delta'} \right) \left(1 + o_P(1) \right) \left(\max_{i \in \mathcal{I}_n(y)} \frac{(\Delta_i^n(\sigma \cdot W))^2}{2 \left(\sup_{t \in [\frac{i-k_n/2}{n}, \frac{i+k_n/2}{n}] \sigma_t^2} \right) h \log n} + o_P(1) \right), \end{aligned} \quad (\text{A.16})$$

where on the last line we applied Proposition B.4. Together with Lemma B.6, this shows, for small enough $\eta > 0$, with probability tending to 1,

$$\max_{i \in \mathcal{I}_n(y)} \frac{(\Delta_i^n X)^2}{r_n^* B_{n,0}^*(i)} \leq 1 - \eta,$$

which in turn implies (A.9) holds with probability tending to 1. The statement (A.10) is shown along the same lines of (A.16), replacing $\widehat{C}_{n,0}(i)$ and $\underline{c}_n(i)$ in (A.16) with $\mathcal{C}_n(i; y)$ and applying Proposition B.4. $\square$

Proof of Theorem 3.2. We begin by laying out some arguments used across all cases (i)–(iii). First note that for each n , the sequence $\widehat{C}_{n,j}$ (and hence, $B_{n,j}$) is either nonincreasing or nondecreasing in j . Indeed, suppose that $\widehat{C}_{n,0} \leq \widehat{C}_{n,1}$. Then, $B_{n,0} \leq B_{n,1}$, giving

$$\widehat{C}_{n,2} = \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_{n,1}\}} \geq \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_{n,0}\}} = \widehat{C}_{n,1}.$$

Then, $B_{n,1} \leq B_{n,2}$ and we can proceed by induction to conclude that $\widehat{C}_{n,j}$ is nondecreasing in j . If $\widehat{C}_{n,0} \geq \widehat{C}_{n,1}$, we can follow the same argument to show that $\widehat{C}_{n,j}$ is nonincreasing in j .

Also, since for each n , the function $B \mapsto \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B\}}$ takes at most $n+1$ possible values, it holds that

$$\widehat{C}_n = \widehat{C}_{n,n+1}, \quad \text{a.s.}$$

For any $y > 0$, we will now show that, for each $j \geq 1$, we have $\widehat{C}_{n,j} \geq \widetilde{C}_{n,j}(y)$. First note the inequality $\widehat{C}_{n,1} \geq \widetilde{C}_{n,1}(y)$ is straightforward by definitions (2.2) and (3.4), which implies $B_{n,1} \geq \widetilde{B}_{n,1}(y)$. Proceeding by induction, suppose that for some $j \geq 1$ we have $\widehat{C}_{n,j} \geq \widetilde{C}_{n,j}(y)$. Then, by definition, $B_{n,j} \geq \widetilde{B}_{n,j}(y)$, implying

$$\begin{aligned} \widehat{C}_{n,j+1} &= \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_{n,j}\}} \\ &\geq \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq \widetilde{B}_{n,j}(y)\}} \\ &\geq \sum_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq \widetilde{B}_{n,j}(y)\}} = \widetilde{C}_{n,j+1}(y). \end{aligned} \quad (\text{A.17})$$

Therefore, for all $j \geq 1$, $\widehat{C}_{n,j} \geq \widetilde{C}_{n,j}(y)$; in particular, $\widehat{C}_n = \widehat{C}_{n,n+1} \geq \widetilde{C}_{n,n+1}(y)$. Next, we decompose $\widehat{C}_n$ as

$$\begin{aligned} \widehat{C}_n &= \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_{n,n}\}} \\ &= \sum_{i=1}^n (\Delta_i^n X)^2 \left(\mathbf{1}_{\{\Delta_i^n N(y)=0, \Delta_i^n N'=0\}} + \mathbf{1}_{\{\Delta_i^n N(y)=0, \Delta_i^n N' \neq 0, |\Delta_i^n X| \leq B_{n,n}\}} \right. \\ &\quad \left. - \mathbf{1}_{\{\Delta_i^n N(y)=0, |\Delta_i^n X| > B_{n,n}, \Delta_i^n N'=0\}} + \mathbf{1}_{\{\Delta_i^n N(y) \neq 0, |\Delta_i^n X| \leq B_{n,n}\}} \right) \\ &\leq \sum_{i=1}^n (\Delta_i^n X)^2 \left(\mathbf{1}_{\{\Delta_i^n N(y)=0, \Delta_i^n N'=0\}} + \mathbf{1}_{\{\Delta_i^n N(y)=0, \Delta_i^n N' \neq 0, |\Delta_i^n X| \leq B_{n,n}\}} \right. \\ &\quad \left. + \mathbf{1}_{\{\Delta_i^n N(y) \neq 0, |\Delta_i^n X| \leq B_{n,n}\}} \right) \\ &= \mathcal{E}_n(y) + R_n, \end{aligned}$$

where $R_n = \sum_{i=1}^n (\Delta_i^n X)^2 \left(\mathbf{1}_{\{\Delta_i^n N(y) \neq 0, |\Delta_i^n X| \leq B_{n,n}\}} + \mathbf{1}_{\{\Delta_i^n N(y)=0, \Delta_i^n N' \neq 0, |\Delta_i^n X| \leq B_{n,n}\}} \right)$. Thus,

$$\widetilde{C}_{n,n+1}(y) \leq \widehat{C}_n \leq \mathcal{E}_n(y) + R_n. \quad (\text{A.18})$$

Further, with the sequence of events $\Omega_n := \{\widetilde{C}_{n,n+1}(y) = \mathcal{E}_n(y)\}$, we may write expression (A.18) as

$$\mathcal{E}_n(y) + \mathbf{1}_{\Omega_n^c} (\widetilde{C}_{n,n+1}(y) - \mathcal{E}_n(y)) \leq \widehat{C}_n \leq \mathcal{E}_n(y) + R_n. \quad (\text{A.19})$$

We now turn to the statement (i), in which case we recall r_n is assumed to satisfy

$$h \log(1/h) \ll r_n \ll (h \log(1/h))^{\frac{\alpha}{2}}. \quad (\text{A.20})$$

Under this assumption, we may choose a sequence $y_n \rightarrow 0$ such that, for some small $\delta > 0$,

$$r_n^{\frac{1}{\alpha}} \vee h^{(\frac{1}{\alpha} \wedge 1) - \delta} \ll y_n \ll (h \log n)^{\frac{1}{2}}, \quad (\text{A.21})$$

under which the hypotheses of Proposition B.2(i) are satisfied, giving $\mathcal{E}_n(y) \xrightarrow{P} C_T$.

Now, since $r_n \gg h \log n$, clearly $r_n \geq 2\left(\frac{\bar{\sigma}^2}{\int_0^1 \sigma_s^2 ds}\right)h \log n$ a.s. for all large n , showing the hypotheses of Proposition A.1 are satisfied, giving $\mathbb{P}(\Omega_n) \rightarrow 1$. Further, by the assumption (A.21), $y^\alpha \gg r_n$, giving

$$\mathbb{E}(N_1(y)r_n) = O(y^{-\alpha}r_n) = o(1).$$

Therefore,

$$0 \leq R_n \leq N_1(y)B_{n,n}^2 + N_1'(y)B_{n,n}^2 = N_1(y)r_n\hat{C}_{n,n-1} + N_1'(y)r_n\hat{C}_{n,n-1} \xrightarrow{P} 0,$$

where above we used that $\hat{C}_{n,n-1} = \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_{n,n-1}\}} \leq \sum_{i=1}^n (\Delta_i^n X)^2 = O_P(1)$. Since $\mathbf{1}_{\Omega_n^c}(\mathcal{E}_n(y) - \tilde{C}_{n,n+1}(y)) = o_P(1)$, expression (A.19) and $R_n = o_P(1)$ give (i).

We now establish (ii), in which case, we recall r_n is assumed to satisfy

$$h \log(1/h) \ll r_n \ll h^{\frac{\alpha+1}{2}} (\log(1/h))^{\frac{\alpha}{2}}. \quad (\text{A.22})$$

Under these constraints, we may again select a sequence $y \rightarrow 0$ satisfying (A.21). By (A.19), we have

$$\begin{aligned} & \sqrt{n}(\mathcal{E}_n(y) - C_T) + \sqrt{n}\mathbf{1}_{\Omega_n^c}(\tilde{C}_{n,n+1}(y) - \mathcal{E}_n(y)) \\ & \leq \sqrt{n}(\hat{C}_n - C_T) \\ & \leq \sqrt{n}(\mathcal{E}_n(y) - C_T) + \sqrt{n}R_n. \end{aligned} \quad (\text{A.23})$$

Note that, for the right side of (A.23), the condition (A.22) gives $r_n \ll n^{-1/2}y^\alpha$, which yields

$$\begin{aligned} \sqrt{n}R_n &= \sqrt{n} \sum_{i=1}^n (\Delta_i^n X)^2 (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0, |\Delta_i^n X| \leq B_{n,n}\}} + \mathbf{1}_{\{\Delta_i^n N(y) = 0, \Delta_i^n N' \neq 0, |\Delta_i^n X| \leq B_{n,n}\}}) \\ &\leq \sqrt{n}N_1(y)r_n\hat{C}_{n,n} + \sqrt{n}N_1'(y)r_n\hat{C}_{n,n} \\ &= O_P(n^{1/2}r_n y^{-\alpha}) + O_P(n^{1/2}r_n) \xrightarrow{P} 0. \end{aligned}$$

Moreover, since $\alpha < 1$, we have $y \ll (h \log n)^{1/2} \ll h^{\frac{1}{2(2-\alpha)}}$, implying the hypotheses of Proposition B.2(ii) are satisfied, and the hypotheses of Proposition A.1 are clearly also satisfied. Statement (ii) then follows from Proposition B.2(ii), Proposition A.1, and the string of inequalities in (A.23), since $\mathbb{P}(\Omega_n^c) \rightarrow 0$, implies

$$\sqrt{n}\mathbf{1}_{\Omega_n^c}(\tilde{C}_{n,n+1}(y) - \mathcal{E}_n(y)) = o_P(1). \quad (\text{A.24})$$

For statement (iii), we now take $y \rightarrow 0$ such that

$$h^{\frac{1}{2}} \ll y \ll (h \log n)^{\frac{1}{2}}.$$

In particular, under such choice of y , from Proposition B.2(iii) we have $\sqrt{n}(\mathcal{E}_n(y) - C_T) \xrightarrow{P} \infty$. Again from (A.19), we have the string of inequalities (A.23), and since we still have $r_n \gg h \log(1/h)$, we may apply Proposition A.1 to again conclude (A.24), and thus the leftmost inequality in (A.23) yields the result. $\square$

Proof of Proposition 3.5. For simplicity we write Y instead of Y' throughout the proof. Let $Y_t = \int_0^t \sigma_s dW_s$, where

$$\sigma_t = a\mathbf{1}_{\{t < \theta\}} + b\mathbf{1}_{\{t \geq \theta\}},$$

so that $C_T(Y) = a^2\theta + b^2(1 - \theta)$. Above, the quantities $\theta \in (0, 1)$, $0 < a < b$ are nonrandom constants chosen so that

$$\delta := \frac{c_0}{2} \cdot \frac{C_T(Y)}{b^2} = \frac{c_0}{2} \left(\frac{a^2}{b^2} \theta + (1 - \theta) \right) \in \left(0, \frac{1}{2} \right).$$

We have

$$\begin{aligned} \text{TRV}_n(Y; \vartheta_n) - C_T(Y) &= \left(\sum_{i=1}^n (\Delta_i^n Y)^2 - C_T(Y) \right) - \sum_{i=1}^n (\Delta_i^n Y)^2 \mathbf{1}_{\{|\Delta_i^n Y| > \vartheta_n\}} \\ &= O_P(n^{-1/2}) - \left(\sum_{i=1}^{\lfloor n\theta \rfloor} + \sum_{i=\lfloor n\theta \rfloor + 2}^n \right) (\Delta_i^n Y)^2 \mathbf{1}_{\{|\Delta_i^n Y| > \vartheta_n\}}. \end{aligned} \quad (\text{A.25})$$

For simplicity let $\mathcal{Y}_i^n = (\Delta_i^n Y)^2 \mathbf{1}_{\{|\Delta_i^n Y| > \vartheta_n\}}$. For $Z \sim \mathcal{N}(0, 1)$, $\phi(x) = (2\pi)^{-1/2} e^{-x^2/2}$, an integration by parts shows

$$\mathbb{E} Z^{2k} \mathbf{1}_{\{|Z| > x\}} \sim 2x^{2k-1} \phi(x), \quad x \rightarrow \infty. \quad (\text{A.26})$$

Hence, for $i > \lfloor n\theta \rfloor + 1$, $\Delta_i^n Y \stackrel{d}{=} b\sqrt{h}Z$, and since $c_0 C_T(Y)/b^2 = 2\delta$,

$$\begin{aligned} \mathbb{E} \mathcal{Y}_i^n &= \mathbb{E} (\Delta_i^n Y)^2 \mathbf{1}_{\{|b\sqrt{h}Z| > \sqrt{c_0 C_T(Y) h \log(1/h)}\}} \\ &= b^2 h \mathbb{E} Z^2 \mathbf{1}_{\{|Z| > \sqrt{2\delta \log n}\}} \\ &\sim 2b^2 h \sqrt{2\delta \log n} \phi\left(\sqrt{2\delta \log n}\right), \end{aligned} \quad (\text{A.27})$$

giving

$$\sum_{i=\lfloor n\theta \rfloor + 2}^n \mathbb{E} \mathcal{Y}_i^n \sim 2b^2(1-\theta) \sqrt{2\delta \log n} \cdot (2\pi)^{-1/2} n^{-\delta} =: K_0 n^{-\delta} \sqrt{\log n}.$$

Analogously, for $i \leq \lfloor n\theta \rfloor$, $\Delta_i^n Y \stackrel{d}{=} a\sqrt{h}Z$, and since $c_0 C_T(Y)/a^2 = 2(b^2/a^2)\delta > 2\delta$, we have $\mathbb{E} \sum_{i=1}^{\lfloor n\theta \rfloor} \mathcal{Y}_i^n = O(n^{-\delta(b^2/a^2)} \sqrt{\log n}) = o(n^{-\delta} \sqrt{\log n})$. This implies

$$\frac{n^\delta}{\sqrt{\log n}} \sum_{i=1}^n \mathbb{E} \mathcal{Y}_i^n \rightarrow K_0. \quad (\text{A.28})$$

On the other hand, using (A.26), we have, for $i > \lfloor n\theta \rfloor + 1$,

$$\mathbb{E} (\mathcal{Y}_i^n)^2 = b^4 h^2 \mathbb{E} Z^4 \mathbf{1}_{\{|Z| > \sqrt{2\delta \log n}\}} = O(n^{-2-\delta} (\log n)^{3/2}).$$

Since (A.27) gives $(\mathbb{E} \mathcal{Y}_i^n)^2 = O(n^{-2-2\delta} \log n)$, we obtain $\text{Var}(\mathcal{Y}_i^n) = O(n^{-2-\delta} (\log n)^{3/2})$ for all $i > \lfloor n\theta \rfloor + 1$. Arguing analogously as for (A.28), for $i \leq \lfloor n\theta \rfloor$ we have $\text{Var}(\mathcal{Y}_i^n) \ll n^{-2-\delta} (\log n)^{3/2}$. Thus,

$$\text{Var}\left(\sum_{i=1}^n \mathcal{Y}_i^n\right) = O(n^{-1-\delta} (\log n)^{3/2}),$$

giving

$$\frac{n^\delta}{\sqrt{\log n}} \sum_{i=1}^n (\mathcal{Y}_i^n - \mathbb{E} \mathcal{Y}_i^n) = o_P(n^{(\delta-1)/2} (\log n)^{1/4}) = o_P(1).$$

Thus, from (A.25), we obtain $\text{TRV}_n(Y; \vartheta_n) \xrightarrow{P} C_T$ and

$$\begin{aligned} n^{1/2} \left(\text{TRV}_n(Y; \vartheta_n) - C_T(Y) \right) &= O_P(1) - n^{1/2} \left(\sum_{i=1}^n \mathbb{E} \mathcal{Y}_i^n + \sum_{i=1}^n (\mathcal{Y}_i^n - \mathbb{E} \mathcal{Y}_i^n) \right) \\ &= O_P(1) - (n^{\frac{1}{2}-\delta} \sqrt{\log n}) \left(\frac{n^\delta}{\sqrt{\log n}} \sum_{i=1}^n \mathbb{E} \mathcal{Y}_i^n + o_P(1) \right) \\ &= O_P(1) - (n^{\frac{1}{2}-\delta} \sqrt{\log n}) (K_0 + o_P(1)) \\ &\xrightarrow{P} -\infty. \end{aligned}$$

□

Proof of Theorem 3.6. The proof is similar to that of Theorem 3.2; we provide details here where there are substantive differences. We first note that for each n and i , $\widehat{c}_{n,j}(i)$ (and, hence, $B_{n,j}^*(i)$) is either nonincreasing or nondecreasing in j . Indeed, suppose that $\widehat{c}_{n,0}(i) \geq \widehat{c}_{n,1}(i)$. Then, $B_{n,0}^*(i) \geq B_{n,1}^*(i)$ and thus,

$$\begin{aligned}\widehat{c}_{n,2}(i) &= \frac{1}{k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} (\Delta_\ell^n X)^2 \mathbf{1}_{\{|\Delta_\ell^n X| \leq B_{n,1}^*(i)\}} \\ &\leq \frac{1}{k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} (\Delta_\ell^n X)^2 \mathbf{1}_{\{|\Delta_\ell^n X| \leq B_{n,0}^*(i)\}} = \widehat{c}_{n,1}(i).\end{aligned}$$

Then, $B_{n,1}^*(i) \geq B_{n,2}^*(i)$ and we can proceed by induction to conclude that $\widehat{c}_{n,j}(i)$ is nonincreasing in j . Analogously, in the case $\widehat{c}_{n,0}(i) \leq \widehat{c}_{n,1}(i)$, both sequences $\widehat{c}_{n,j}(i)$ and $B_{n,j}^*(i)$ are nondecreasing in j . As a consequence, we will have that $B_n^*(i) = B_{n,n+1}^*(i)$ and

$$\widehat{C}_n^* = \sum_{i=1}^n (\Delta_i^n X)^2 \mathbf{1}_{\{|\Delta_i^n X| \leq B_{n,n+1}^*(i)\}}.$$

Next, for any $y > 0$, recalling $\widetilde{C}_{n,j}^*(y)$ as in (A.3), and $\widetilde{c}_{n,j}(i; y)$, $\widetilde{B}_{n,j}^*(i; y)$ as in (A.2), clearly $\widetilde{c}_{n,1}(i; y) \leq \widehat{c}_{n,1}(i)$, $i = 1, \dots, n$, giving $\widetilde{B}_{n,1}^*(i; y) \leq B_{n,1}^*(i)$, $i = 1, \dots, n$. Arguing inductively in an analogous manner to (A.17), we then obtain

$$\widetilde{B}_{n,j}^*(i; y) \leq B_{n,j}^*(i), \quad i = 1, \dots, n, \quad j \geq 1.$$

Arguing the same fashion as in (A.19), we obtain

$$\mathcal{E}_n(y) + \mathbf{1}_{(\Omega_n^*)^c} (\widetilde{C}_{n,n+1}^*(y) - \mathcal{E}_n(y)) \leq \widehat{C}_n^* \leq \mathcal{E}_n(y) + R_n^*, \quad (\text{A.29})$$

where $\Omega_n^* = \{\widetilde{C}_{n,n+1}^*(y) = \mathcal{E}_n(y)\}$ and

$$R_n^* = \sum_{i=1}^n (\Delta_i^n X)^2 (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0, |\Delta_i^n X| \leq B_{n,n}^*(i)\}} + \mathbf{1}_{\{\Delta_i^n N(y) = 0, \Delta_i^n N' \neq 0, |\Delta_i^n X| \leq B_{n,n}^*(i)\}}).$$

Turning now to (i), choose $y \rightarrow 0$ in such a way that (A.21) holds with r_n^* replacing r_n . Applying Proposition A.1, we obtain $\mathbb{P}(\Omega_n^*) \rightarrow 1$. The statement then follows from Proposition B.2(i), the convergence $\mathbf{1}_{\Omega_n^c} (\widetilde{C}_{n,n+1}^*(y) - \mathcal{E}_n(y)) = o_P(1)$, expression (A.29), and

$$\begin{aligned}0 \leq R_n^* &\leq N_1(y) \left(\max_{1 \leq i \leq n} B_n^*(i) \right)^2 + N_1' \left(\max_{1 \leq i \leq n} B_n^*(i) \right)^2 \\ &\leq O_P(y^{-\alpha} r_n^*) + O_P(r_n^*) \xrightarrow{P} 0,\end{aligned}$$

since $\max_{1 \leq i \leq n} (B_n^*(i))^2 / r_n^* \leq \max_{1 \leq i \leq n} \widehat{c}_{n,n+1}(i) \leq \max_{1 \leq i \leq n} \frac{n}{k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} (\Delta_\ell^n X)^2 = O_P(1)$ (see [21]), and $r_n^* \ll y^\alpha$ by assumption. Similarly, for (ii), we again take $y \rightarrow 0$ such that (A.21) is satisfied (with r_n^* replacing r_n). Then, from expression (A.29), we may arrive at (A.23) with $\widetilde{C}_{n,j}^*(y)$ replacing $\widetilde{C}_{n,j}(y)$ and Ω_n^* replacing Ω_n . The statement then follows in view of the convergence $\sqrt{n} \mathbf{1}_{\Omega_n^c} (\widetilde{C}_{n,n+1}^*(y) - \mathcal{E}_n(y)) = o_P(1)$, Proposition B.2(ii), and the estimates

$$\begin{aligned}0 \leq \sqrt{n} R_n^* &\leq N_1(y) \left(\max_{1 \leq i \leq n} B_n^*(i) \right)^2 + N_1' \left(\max_{1 \leq i \leq n} B_n^*(i) \right)^2 \\ &\leq O_P(\sqrt{n} y^{-\alpha} r_n^*) + O_P(\sqrt{n} r_n^*) \xrightarrow{P} 0,\end{aligned}$$

since $r_n^* \ll n^{-1/2} y^\alpha$ by assumption. Statement (iii) is proved analogously to Theorem 3.2(iii). $\square$

B Auxiliary results

Throughout this section, for notational simplicity, we again often omit the subscript n in h_n and y_n .

Lemma B.1. *Let $k > 0$. Then for every fixed y and n , for $M_t(y)$ as in (3.1),*

$$\mathbb{P}(|\Delta_i^n M(y)| > k) \leq 2 \left(\frac{eh\Sigma_y}{ky} \right)^{\frac{k}{2y}},$$

where $\Sigma_y := \int_{|z| \leq y} z^2 \nu(dz) \sim Ky^{2-\alpha}$ for some constant K .

Proof. Note that for each fixed y , $\Delta_i^n M(y)$ is infinitely divisible with triplet $(0, 0, \mathbf{1}_{\{|x| \leq y\}} \nu(dx))$, and in particular $\mathbb{E} e^{u\Delta_i^n M(y)} < \infty$ for all $u \in \mathbb{R}$. Thus, by Lemma 26.4 in [?], we have

$$\mathbb{P}(\Delta_i^n M(y) > k) \leq \exp \left\{ -h \int_0^{k/h} \tau(x) dx \right\},$$

where $\tau : [0, \infty) \rightarrow \mathbb{R}$ is the inverse function of $s : [0, \infty) \rightarrow \mathbb{R}$ defined by

$$s(x) = \int_{|z| \leq y} z(e^{xz} - 1) \nu(dz).$$

Recall $\Sigma_y := \int_{|z| \leq y} z^2 \nu(dz)$, and note that,

$$s(x) \leq \int_{|z| \leq y} xz^2 e^{xz} \nu(dz) \leq x e^{xy} \Sigma_y \leq \frac{e^{2xy} - 1}{y} \Sigma_y.$$

This implies that

$$\tau(z) \geq \frac{1}{2y} \log \left(1 + \frac{y}{\Sigma_y} z \right).$$

So,

$$\begin{aligned} \mathbb{P}(\Delta_i^n M(y) > k) &\leq \exp \left\{ -\frac{h}{2y} \int_0^{k/h} \log \left(1 + \frac{y}{\Sigma_y} z \right) dz \right\} \\ &= \exp \left\{ -\frac{h\Sigma_y}{2y^2} \int_0^{\frac{ky}{h\Sigma_y}} \log(1+s) ds \right\} \\ &\leq \exp \left\{ -\frac{k}{2y} \log \frac{ky}{eh\Sigma_y} \right\} \\ &= \left(\frac{eh\Sigma_y}{ky} \right)^{\frac{k}{2y}}. \end{aligned}$$

The argument for $\mathbb{P}(\Delta_i^n M(y) < -k)$ is analogous. $\square$

For statements ahead, recall $\mathcal{C}_n(y) = \sum_{i \in \mathcal{I}_n(y)} (\Delta_i^n X)^2$, where $\mathcal{I}_n(y) = \{i : \Delta_i^n N(y) = 0, \Delta_i^n N' = 0\}$.

Proposition B.2. *The following statements hold.*

(i) *Suppose $y = y_n \rightarrow 0$ satisfies $(h \log n)^{\frac{1}{\alpha}} \vee h^{1-\delta} \ll y$ for some $\delta \in (0, 1)$. Then,*

$$\mathcal{C}_n(y) \xrightarrow{P} C_T. \tag{B.1}$$

(ii) Suppose $\alpha \in (0, 1)$, and $y \rightarrow 0$ such that $(h^{1/2} \log n)^{\frac{1}{\alpha}} \vee h^{1-\delta} \ll y \lesssim h^{\frac{1}{2(2-\alpha)}}$ for some $\delta \in (0, 1)$. Then,

$$\sqrt{n} \left(\mathcal{E}_n(y) - C_T \right) \xrightarrow{st} \mathcal{N} \left(0, 2 \int_0^1 \sigma_s^4 ds \right).$$

(iii) Suppose $\alpha \in (1, 2)$, and $y \rightarrow 0$. Then,

$$\sqrt{n} \left(\mathcal{E}_n(y) - C_T \right) \xrightarrow{P} \begin{cases} -\infty, & h^{\frac{1}{2(2-\alpha)}} \vee h^{\frac{1}{\alpha}} \ll y \ll h^{\frac{1}{2}}, \\ \infty, & y \gg h^{\frac{1}{2}}. \end{cases} \quad (\text{B.2})$$

Proof. Set

$$\tau_n^2(y) = \sum_{i \in \mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2.$$

Observe that, by definition of $N(y)$,

$$\mathbb{E} N_1(y) \leq K \int_{|x|>y} |x|^{-\alpha-1} dx = K y^{-\alpha}.$$

Since Lemma B.6 gives $\limsup_{n \rightarrow \infty} \max_{i=1, \dots, n} \frac{(\Delta_i^n(\sigma \cdot W))^2}{h \log n} < \infty$, we obtain

$$\begin{aligned} \sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} &\leq \max(\Delta_i^n(\sigma \cdot W))^2 N_1(y) \\ &= O_P(h(\log n) y^{-\alpha}) = o_P(1). \end{aligned}$$

Similarly,

$$\sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N' \neq 0\}} \leq N'_1 \max(\Delta_i^n(\sigma \cdot W))^2 \xrightarrow{P} 0.$$

Thus,

$$\begin{aligned} 0 \leq \sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 - \tau_n^2(y) &= \sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 (1 - \mathbf{1}_{\{\Delta_i^n N(y)=0, \Delta_i^n N'=0\}}) \\ &\leq \sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} + \mathbf{1}_{\{\Delta_i^n N' \neq 0\}}) \xrightarrow{P} 0, \end{aligned}$$

and we obtain

$$\tau_n^2(y) = \sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 + o_P(1) \xrightarrow{P} \int_0^1 \sigma_s^2 ds. \quad (\text{B.3})$$

On the other hand, recalling the notation (A.12), we have

$$\begin{aligned} \mathbb{E} (\Delta_i^n \chi(y))^2 &\leq K \left[\mathbb{E} \left(\int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right)^2 + \mathbb{E} (\Delta_i^n b(y))^2 \right] \\ &\leq K (h y^{2-\alpha} + h^2 (y^{1-\alpha} + 1)^2) = O(h y^{2-\alpha}). \end{aligned} \quad (\text{B.4})$$

Thus,

$$\begin{aligned} |\mathcal{E}_n(y) - \tau_n^2(y)| &= \left| \sum_{i \in \mathcal{I}_n(y)} \left[(\Delta_i^n(\sigma \cdot W) + \Delta_i^n \chi(y))^2 - (\Delta_i^n(\sigma \cdot W))^2 \right] \right| \\ &\leq \sum_{i=1}^n \left((\Delta_i^n \chi(y))^2 + 2 |\Delta_i^n \chi(y) \Delta_i^n(\sigma \cdot W)| \right) \\ &\leq \sum_{i=1}^n \left(O_P(h y^{2-\alpha}) + O_P(h y^{\frac{1-\alpha}{2}}) \right) \\ &= o_P(1), \end{aligned}$$

where the second-to-last line follows from an application of Cauchy-Schwarz. Thus,

$$\mathcal{E}_n(y) - \tau_n^2(y) = o_P(1),$$

which, by (B.3), establishes (B.1).

For (ii), note that

$$\begin{aligned} & \sqrt{n}(\tau_n^2(y) - C_T) \\ &= \sqrt{n} \left(\sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 - C_T \right) - \sqrt{n} \sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} + \mathbf{1}_{\{\Delta_i^n N' \neq 0\}}) \\ &\xrightarrow{st} \mathcal{N} \left(0, 2 \int_0^1 \sigma_s^4 ds \right), \end{aligned} \tag{B.5}$$

where the first term of (B.5) converges to $\mathcal{N}(0, 2 \int_0^1 \sigma_s^4 ds)$ stably in law by the usual CLT for realized variance and the second term of (B.5) converges to 0 in probability because

$$\sqrt{n} \sum_{\Delta_i^n N' \neq 0} (\Delta_i^n(\sigma \cdot W))^2 \leq \sqrt{n} N'_1 \max_{1 \leq i \leq n} (\Delta_i^n(\sigma \cdot W))^2 = \sqrt{n} \cdot O(h \log n) \rightarrow 0,$$

and

$$\sqrt{n} \sum_{\Delta_i^n N(y) \neq 0} (\Delta_i^n(\sigma \cdot W))^2 \leq \sqrt{n} N_1(y) \max_{1 \leq i \leq n} (\Delta_i^n(\sigma \cdot W))^2 = \sqrt{n} \cdot O_p(y^{-\alpha}) \cdot O(h \log n) = o_P(1),$$

where we used that $h^{\frac{1}{2\alpha}}(\log n)^{\frac{1}{\alpha}} \ll y$. On the other hand, recalling the notation (A.12), we have

$$\begin{aligned} \sqrt{n} |\mathcal{E}_n(y) - \tau_n^2(y)| &= \left| \sum_{\mathcal{I}_n(y)} \sqrt{n} \left[(\Delta_i^n(\sigma \cdot W) + \Delta_i^n \chi(y))^2 - (\Delta_i^n(\sigma \cdot W))^2 \right] \right| \\ &\leq 2\sqrt{n} \left| \sum_{\mathcal{I}_n(y)} \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) \right| + \sum_{\mathcal{I}_n(y)} \sqrt{n} (\Delta_i^n \chi(y))^2 \\ &=: T_1 + T_2. \end{aligned}$$

Now, by (B.4), we get

$$\mathbb{E} T_2 \leq K n^{3/2} (h y^{2-\alpha} + h^2 (y^{1-\alpha} + 1)^2) \rightarrow 0,$$

where we used that $y \ll h^{\frac{1}{2(2-\alpha)}}$. For T_1 , first note

$$\begin{aligned} T_1 &\leq 2\sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) \right| + 2\sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} + \mathbf{1}_{\{\Delta_i^n N' \neq 0\}}) \right| \\ &= 2\sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) \right| + o_P(1), \end{aligned}$$

since

$$\begin{aligned} \sqrt{n} \left| \sum_{\Delta_i^n N(y) \neq 0} \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) \right| &\leq \sqrt{n} N_1(y) \max_i |\Delta_i^n(\sigma \cdot W)| \max_i |\Delta_i^n \chi(y)| \\ &= \sqrt{n} O_P(y^{-\alpha}) \cdot O(\sqrt{h \log n}) \cdot O_P(y) \\ &= o_P(y^{1-\alpha} \sqrt{\log n}) = o_P(1), \end{aligned}$$

and similarly

$$\sqrt{n} \left| \sum_{\Delta_i^n N' \neq 0} \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) \right| \leq \sqrt{n} N'_1 \cdot O(\sqrt{h \log n}) \cdot O_P(y) = O_P(y \sqrt{\log n}) \xrightarrow{P} 0.$$

Next, we show that

$$\sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) \right| = o_P(1), \quad (\text{B.6})$$

which will imply $T_1 = o_P(1)$, and consequently that $\sqrt{n}(\mathcal{E}_n(y) - \tau_n^2(y)) = o_P(1)$, and in view of (B.5) will establish the desired convergence. Write

$$\begin{aligned} \sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \Delta_i^n \chi(y) \right| &\leq \sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right| + \sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \Delta_i^n b(y) \right| \\ &=: T_{1,1} + T_{1,2}. \end{aligned}$$

Applying Lemma 2.1.5 in [23], since $y \gg h^{1/\alpha}$, we have, for every $p \geq 1$

$$\mathbb{E} \left| \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right|^{2p} \leq K(hy^{2p-\alpha} + h^p y^{p(2-\alpha)}) \leq Khy^{2p-\alpha}.$$

This gives, for every $p, q > 1$ with $p^{-1} + q^{-1} = 1$,

$$\begin{aligned} \mathbb{E} T_{1,1} &\leq (\mathbb{E} T_{1,1}^2)^{1/2} = \sqrt{n} \left(\sum_{i=1}^n \mathbb{E} \left[(\sigma \cdot W) \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right]^2 \right)^{1/2} \\ &\leq \sqrt{n} \left(\sum_{i=1}^n (\mathbb{E}(\sigma \cdot W)^{2q})^{1/q} \left(\mathbb{E} \left| \int_{t_{i-1}}^{t_i} \gamma_t dM_t(y) \right|^{2p} \right)^{1/p} \right)^{1/2} \\ &\leq K \sqrt{n} \left(\sum_{i=1}^n h \cdot (h^{1/p} y^{2-\alpha/p}) \right)^{1/2} \\ &\leq K(hy^{-\alpha})^{\frac{1}{2p}} h^{-\frac{1}{2}} y \\ &\leq K h^{\frac{1}{4p} - \frac{1}{2}} y, \end{aligned} \quad (\text{B.7})$$

$$\leq K h^{\frac{1}{4p} - \frac{1}{2}} y, \quad (\text{B.8})$$

where for the last inequality we used that $y \gg h^{\frac{1}{2\alpha}}$. Since $\frac{1}{2(2-\alpha)} > \frac{1}{4}$, by taking $p > 1$ close enough to 1, we have $\frac{1}{2} - \frac{1}{4p} = \frac{1}{4} + (\frac{1}{4} - \frac{1}{4p}) < \frac{1}{2(2-\alpha)}$ giving $y \ll h^{\frac{1}{2(2-\alpha)}} \ll h^{\frac{1}{2} - \frac{1}{4p}}$, and thus expression (B.8) tends to 0. For $T_{1,2}$, with $b_0(y) = \int_{|x|>y} x \nu(dx)$, and

$$\begin{aligned} \Delta_i^n b(y) &= h(b_{t_{i-1}} + \gamma_{t_{i-1}} b_0(y)) + \int_{t_{i-1}}^{t_i} (b_s - b_{t_{i-1}} + b_0(y)(\gamma_s - \gamma_{t_{i-1}})) ds \\ &=: hb_{t_{i-1}}(y) + \tilde{b}_i(y), \end{aligned}$$

we have

$$T_{1,2} \leq \sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) (hb_{t_{i-1}}(y)) \right| + \sqrt{n} \left| \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \tilde{b}_i(y) \right|.$$

Since

$$\mathbb{E} \left(\sqrt{n} \sum_{i=1}^n \Delta_i^n(\sigma \cdot W) (hb_{t_{i-1}}(y)) \right)^2 = n \sum_{i=1}^n \mathbb{E} (\Delta_i^n(\sigma \cdot W))^2 h^2 \mathbb{E} b_{i-1}^2(y) \leq Kh$$

and

$$\sqrt{n}\mathbb{E}\left|\sum_{i=1}^n \Delta_i^n(\sigma \cdot W) \Delta_i^n \tilde{b}_i(y)\right| \leq \sqrt{n} \sum_{i=1}^n \sqrt{\mathbb{E}(\Delta_i^n(\sigma \cdot W))^2 \mathbb{E}(\tilde{b}_i(y))^2} = o(1),$$

we get $T_{1,1} + T_{1,2} = o_P(1)$, which implies (B.6) and completes the proof of (ii).

We now turn to (iii). Write

$$\begin{aligned} \sqrt{n}(\mathcal{C}_n(y) - C_T) &= \sqrt{n} \left(\sum_{\mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2 - C_T \right) + \sqrt{n} \sum_{\mathcal{I}_n(y)} (\Delta_i^n M(y))^2 + \sqrt{n} \sum_{\mathcal{I}_n(y)} (\Delta_i^n b(y))^2 \\ &\quad + \sqrt{n} \sum_{\mathcal{I}_n(y)} 2[\Delta_i^n M(y)(\Delta_i^n(\sigma \cdot W) + \Delta_i^n b(y)) + \Delta_i^n b(y)\Delta_i^n(\sigma \cdot W)] \\ &=: \sum_{i=0}^3 I_i. \end{aligned} \tag{B.9}$$

For the term I_0 , the argument for (B.5) shows

$$I_0 = \sqrt{n} \left(\sum_{\mathcal{I}_n(y)} (\Delta_i^n(\sigma \cdot W))^2 - C_T \right) = O_P(1) - \sqrt{n} \sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}}.$$

Let $\beta_i^n(y) = (\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} - \mathbb{E}_{i-1}(\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}}$ (with $\mathbb{E}_{i-1}(\cdot) = \mathbb{E}(\cdot | \mathcal{F}_{(i-1)h})$), and observe, for every $p \geq 1$,

$$\begin{aligned} \mathbb{E}_{i-1}(\Delta_i^n(\sigma \cdot W))^{2p} \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} &= \mathbb{E}_{i-1} \mathbb{E}_{i-1} \left[(\Delta_i^n(\sigma \cdot W))^{2p} \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \middle| \Delta_i^n N(y) \right] \\ &\leq K \mathbb{E}_{i-1} \left(\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \mathbb{E}_{i-1} \left[\left(\int_{t_{i-1}}^{t_i} \sigma_s^2 ds \right)^p \middle| \Delta_i^n N(y) \right] \right) \\ &\leq K h^{p+1} y^{-\alpha}. \end{aligned}$$

Using that $h y^{-\alpha} \ll 1$, we obtain $\mathbb{E}_{i-1} |\beta_i^n(y)|^p \leq K h^{p+1} y^{-\alpha}$, giving

$$\mathbb{E} \left(\sqrt{n} \sum_{i=1}^n \beta_i^n(y) \right)^2 = n \sum_{i=1}^n \mathbb{E}(\beta_i^n(y))^2 = O(h y^{-\alpha}). \tag{B.10}$$

Since σ^2 is bounded away from zero almost surely, we have

$$\begin{aligned} \mathbb{E}_{i-1}(\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} &= \mathbb{E}_{i-1} \left(\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \mathbb{E}_{i-1} \left[\int_{t_{i-1}}^{t_i} \sigma_s^2 ds \middle| \Delta_i^n N(y) \right] \right) \\ &\geq K h \mathbb{P}(\Delta_i^n N(y) \neq 0) \sim K h^2 y^{-\alpha}. \end{aligned}$$

Analogously, we have $\mathbb{E}_{i-1}(\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \leq K' h^2 y^{-\alpha}$, i.e., for suitable $K_0, K_1 > 0$,

$$K_0 h^{1/2} y^{-\alpha} \leq \sqrt{n} \sum_{i=1}^n \mathbb{E}_{i-1}(\Delta_i^n(\sigma \cdot W))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \leq K_1 h^{1/2} y^{-\alpha}.$$

Thus, for all large n , from (B.10),

$$-K_1 h^{1/2} y^{-\alpha} \left(1 + o_P(y^{\alpha/2}) \right) + O_P(1) \leq I_0 \leq O_P(1) - K_0 h^{1/2} y^{-\alpha} \left(1 + o_P(y^{\alpha/2}) \right). \tag{B.11}$$

Showing $I_0 = O_P(1 \vee h^{1/2}y^{-\alpha})$. We now consider I_1 . Let

$$\xi_{j,y} = y^{-(j-\alpha)} \int_{|x| \leq y} x^j \nu(dx) = y^{-(j-\alpha)} h^{-1} \mathbb{E}(\Delta_i^n M(y))^j, \quad j = 2, 4,$$

(e.g., [?, Example 25.1]) which satisfy $\liminf_{n \rightarrow \infty} \xi_{j,y} > 0$ under Assumption 2.1(ii). Observe $\text{Var}(\sum_{i=1}^n (\Delta_i^n M(y))^2) = y^{4-\alpha} \xi_{4,y} - h y^{4-2\alpha} \xi_{2,y}^2 = O(y^{4-\alpha})$, and

$$\begin{aligned} \sum_{i=1}^n \mathbb{E} |(\Delta_i^n M(y))^2 - \mathbb{E}(\Delta_i^n M(y))^2|^4 &\leq nK \left(\mathbb{E}(\Delta_i^n M(y))^8 + [\mathbb{E}(\Delta_i^n M(y))^2]^4 \right) \\ &\leq K y^{8-\alpha} + K' h^3 y^{8-4\alpha} \\ &= o((y^{4-\alpha})^2). \end{aligned}$$

Therefore, the Lyapunov CLT implies that

$$\begin{aligned} \frac{\sum_{i=1}^n (\Delta_i^n M(y))^2 - n \mathbb{E}(\Delta_i^n M(y))^2}{\sqrt{n \text{Var}((\Delta_i^n M(y))^2)}} &= \frac{\sum_{i=1}^n (\Delta_i^n M(y))^2 - y^{2-\alpha} \xi_{2,y}}{\sqrt{y^{4-\alpha} \xi_{4,y} - h y^{4-2\alpha} \xi_{2,y}^2}} \\ &= \frac{\sum_{i=1}^n (\Delta_i^n M(y))^2 - y^{2-\alpha} \xi_{2,y}}{y^{2-\frac{\alpha}{2}} \xi_{4,y}^{1/2} \sqrt{1 + o(1)}} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1). \end{aligned}$$

This gives

$$\begin{aligned} I_1 &= \sqrt{n} \sum_{i=1}^n (\Delta_i^n M(y))^2 \mathbf{1}_{\{\Delta_i^n N(y)=0\}} \\ &= \sqrt{n} \sum_{i=1}^n (\Delta_i^n M(y))^2 - \sqrt{n} \sum_{i=1}^n (\Delta_i^n M(y))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \end{aligned} \quad (\text{B.12})$$

$$= h^{-1/2} \left(y^{2-\alpha} \xi_{2,y} + y^{2-\frac{\alpha}{2}} \xi_{4,y}^{1/2} \sqrt{1 + o(1)} \cdot Y_n \right) + o_P \left(h^{-1/2} y^{2-\alpha} \right), \quad (\text{B.13})$$

where $Y_n \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1)$. The order of the second term in (B.12) is a consequence of

$$\mathbb{E} \left(\frac{\sqrt{n} \sum_{i=1}^n (\Delta_i^n M(y))^2 \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}}}{h^{-1/2} y^{2-\alpha}} \right) = y^{\alpha-2} \mathbb{E}(N_1(y)) \mathbb{E}((\Delta_i^n M(y))^2) = O(h y^{-\alpha}) = o(1),$$

since $h^{1/\alpha} \ll y$. Moreover, (B.13) implies $I_1/(h^{-1/2} y^{2-\alpha} \xi_{2,y}) \xrightarrow{P} 1$. Next, for I_2 , observe

$$\sqrt{n} \sum_{\mathcal{I}_n(y)} (\Delta_i^n b(y))^2 \leq \sqrt{n} \sum_{i=1}^n (\Delta_i^n b(y))^2 = K n^{3/2} h^2 y^{2-2\alpha} = o(h^{1/2} y^{-\alpha}),$$

showing $I_2 = o_P(I_0)$. We now show $I_3 = o_P(|I_0| + I_1)$. For the first term in I_3 , (B.7) shows, for any $p > 1$,

$$\sqrt{n} \sum_{\Delta_i^n N(y)=0} |(\Delta_i^n(\sigma \cdot W))(\Delta_i^n M(y))| = O_P((h y^{-\alpha})^{\frac{1}{2p}}),$$

which is smaller than $|I_0|$ since $h^{\frac{1}{p}(\frac{1}{2}-\frac{\alpha}{4})} \ll h^{1/2} y^{-\alpha} \iff y^\alpha \ll h^{\frac{1}{2}-\frac{1}{p}(\frac{1}{2}-\frac{\alpha}{4})} \iff y \ll h^{\frac{1}{2\alpha}}$. Also, noting in this case $b_t(y)$ is deterministic, we have

$$\begin{aligned} \mathbb{E} \left(\sqrt{n} \sum_{\Delta_i^n N(y)=0} |(\Delta_i^n b(y))(\Delta_i^n M(y))| \right) &\leq \sqrt{n} \sqrt{\sum_{i=1}^n (\Delta_i^n b(y))^2} \sqrt{\sum_{i=1}^n \mathbb{E}(\Delta_i^n M(y))^2} \\ &= O(y^{2-3\alpha/2}) = o_P(I_1), \end{aligned}$$

since $y^{2-3\alpha/2} \ll h^{-1/2}y^{2-\alpha}$ under $y \gg h^{\frac{1}{\alpha}}$. Similarly,

$$\begin{aligned} & \mathbb{E} \left(\sqrt{n} \sum_{\Delta_i^n N(y)=0} |(\Delta_i^n(\sigma \cdot W))(\Delta_i^n b(y))| \right) \\ & \leq 2\sqrt{n} \max_{1 \leq i \leq n} |\Delta_i^n b(y)| \sqrt{\mathbb{E} \left(\sum_{i=1}^n (\Delta_i^n(\sigma \cdot W))^2 \right)} \leq Kh^{1/2}y^{1-\alpha} = o(h^{-1/2}y^{-\alpha}), \end{aligned}$$

which shows $I_3 = o_P(|I_0| + I_1)$, and thus the limit (B.9) is determined by $I_0 + I_1$. Turning to the first case in (B.2), under $h^{\frac{1}{2(2-\alpha)}} \vee h^{\frac{1}{\alpha}} \ll y \ll h^{\frac{1}{2}}$, we have $I_1 = o_P(|I_0|)$ (since $y \ll h^{1/2}$ gives $h^{-1/2}y^{2-\alpha} \ll h^{1/2}y^{-\alpha}$), and thus from (B.11) we have $I_0 \xrightarrow{P} -\infty$, giving $\sqrt{n}(\mathcal{C}_n(y) - C_T) \xrightarrow{P} -\infty$. For the second case in (B.2), under $y \gg h^{1/2}$ we have instead $|I_0| = o_P(I_1)$, and from (B.13) we have $I_1 \xrightarrow{P} \infty$, giving $\sqrt{n}(\mathcal{C}_n(y) - C_T) \xrightarrow{P} \infty$. $\square$

Below, in Propositions B.3 and B.4, we establish the convergence in probability of the sequences $\underline{C}_n(X)$ and $\underline{c}_n(X; i)$ that are used in the proof of Proposition A.1. Recall that for notational convenience we set $F(\Delta_i^n Y, \dots, \Delta_{i+d-1}^n Y) = 0$ when $i + d - 1 > n$ or $i \leq 0$.

Proposition B.3. *Define the set*

$$\mathcal{I}_n^{(d)}(y) = \{i = 1, \dots, n - d + 1 : \Delta_{i+\ell}^n N(y) = 0, \Delta_{i+\ell}^n N' = 0, \quad \ell = 0, \dots, d-1\}. \quad (\text{B.14})$$

Let $\widehat{C}_{n,0} = \widehat{C}_{n,0}(X)$ belong to the class $\mathcal{C}$ with $\widehat{C}_{n,0}(\cdot)$ defined as in (3.5) and suppose (3.8) holds. With δ_0 as in (3.7), let $y = y_n \rightarrow 0$ in such a way that

$$(h \log n)^{\frac{1}{\alpha}} \ll y \ll h^{\frac{1}{2}} (\log n)^{\frac{1}{2} - \frac{1}{\delta_0}}, \quad (\text{B.15})$$

and for F as in (3.5), define

$$\underline{C}_n(Y) = \sum_{i \in \mathcal{I}_n^{(d)}(y)} F(\Delta_i^n Y, \dots, \Delta_{i+d-1}^n Y),$$

for a generic process Y . Then, $\underline{C}_n(X) \leq \widehat{C}_{n,0}$, and

$$\underline{C}_n(X) \xrightarrow{P} \int_0^1 \sigma_s^2 ds. \quad (\text{B.16})$$

Proof. Observe $\underline{C}_n(X) \leq \widehat{C}_{n,0}$ is immediate, so we need only prove (B.16). Clearly

$$\begin{aligned} & \sum_{i=1}^n F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) - \underline{C}_n(\sigma \cdot W; y) \\ &= \sum_{i=1}^n F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) \left(\sum_{\ell=0}^{d-1} (\mathbf{1}_{\{\Delta_{i+\ell}^n N(y) \neq 0\}} + \mathbf{1}_{\{\Delta_{i+\ell}^n N' \neq 0\}}) \right) \\ &=: T_1 + T_2. \end{aligned}$$

Note that, using (A.1), for each i ,

$$\mathbb{P}(\Delta_i^n N(y) \neq 0) \leq Khy^{-\alpha} = o\left(\frac{1}{\log n}\right),$$

since $h(\log n)y^{-\alpha} = o(1)$ by (A.4). In particular,

$$\mathbb{E} \sum_{i=1}^n \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \leq Ky^{-\alpha}.$$

Using (3.6), this gives

$$\begin{aligned}
T_1 &\leq K \max(\Delta_i^n(\sigma \cdot W))^2 \sum_{i=1}^n \sum_{\ell=0}^{d-1} \mathbf{1}_{\{\Delta_{i+\ell}^n N(y) \neq 0\}} \\
&\leq K \max(\Delta_i^n(\sigma \cdot W))^2 \sum_{i=1}^n \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \\
&= K \frac{\max(\Delta_i^n(\sigma \cdot W))^2}{h \log n} O_P(hy^{-\alpha} \log n) \\
&= O_P(1) \cdot o_P(1).
\end{aligned}$$

Similarly, again using (3.6),

$$T_2 = \sum_{i=1}^n F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) \sum_{\ell=0}^{d-1} \mathbf{1}_{\{\Delta_{i+\ell}^n N' \neq 0\}} \leq K N'_1 \max(\Delta_i^n(\sigma \cdot W))^2 = o_P(1).$$

Thus,

$$\sum_{i=1}^n F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) - \underline{C}_n(\sigma \cdot W) = o_P(1). \quad (\text{B.17})$$

Using assumption (3.8) together with (B.17), we obtain

$$\underline{C}_n(\sigma \cdot W) \xrightarrow{P} \int_0^1 \sigma_s^2 ds. \quad (\text{B.18})$$

On the other hand, using the notation (A.12), we have $\Delta_i^n X = \Delta_i^n(\sigma \cdot W) + \Delta_i^n \chi(y)$. Furthermore, (A.13) gives $\max_{i \in \mathcal{I}_n^{(d)}(y)} |\Delta_i^n \chi(y)| \leq \max_{i \in \mathcal{I}_n(y)} |\Delta_i^n \chi(y)| = O_P(y) = o_P(1)$ and Lemma B.6 gives $\max_{i \in \mathcal{I}_n^{(d)}(y)} |\Delta_i^n(\sigma \cdot W)| \leq \max_{1 \leq i \leq n} |\Delta_i^n(\sigma \cdot W)| = O_P(\sqrt{h \log n}) = o_P(1)$, so that

$$\max_{i \in \mathcal{I}_n(y)} |\Delta_i^n(\sigma \cdot W)| \vee \max_{i \in \mathcal{I}_n^{(d)}(y)} |\Delta_i^n \chi(y)| \leq 1$$

with probability tending to 1. Thus, we may use condition (3.7) to obtain

$$\begin{aligned}
&|\underline{C}_n(X) - \underline{C}_n(\sigma \cdot W)| \\
&= \left| \sum_{i \in \mathcal{I}_n^{(d)}(y)} \left[F(\Delta_i^n X, \dots, \Delta_{i+d-1}^n X) - F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) \right] \right| \\
&\leq nK \left(\max_{i \in \mathcal{I}_n(y)} |\Delta_i^n \chi(y)| \right)^{\delta_0} \left(\left(\max_{i \in \mathcal{I}_n(y)} |\Delta_i^n \chi(y)| \right)^{2-\delta_0} + \left(\max_{i \in \mathcal{I}_n(y)} |\Delta_i^n(\sigma \cdot W)| \right)^{2-\delta_0} \right) + o_P(1) \\
&= n O_P(y^{\delta_0}) \cdot \left(O_P(y^{2-\delta_0}) + O_P\left((h \log n)^{\frac{2-\delta_0}{2}}\right) \right) + o_P(1) \\
&= O_P(ny^2) + o_P\left(O_P(y^{\delta_0} n^{\frac{\delta_0}{2}} (\log n)^{1-\frac{\delta_0}{2}}\right) + o_P(1) \\
&= o_P(1),
\end{aligned} \quad (\text{B.19})$$

where the last line follows from the condition (B.15). Thus,

$$\underline{C}_n(y; X) - \underline{C}_n(y; \sigma \cdot W) = o_P(1),$$

which by (B.18) establishes (B.16). $\square$

Proposition B.4. Let $\widehat{c}_{n,0} = \widehat{c}_n(X; i)$, $i = 1, \dots, n$ belong to class $\mathcal{C}^{spot}$ with $\widehat{c}_n(\cdot; i)$ of the form (3.13). For each $y > 0$, with $\mathcal{I}_n^{(k)}(y)$ as in (B.14), define

$$\underline{c}_n(X; i) = \frac{n}{k_n} \sum_{\ell=i-k_n/2+1}^{i+k_n/2} F(\Delta_\ell^n X, \dots, \Delta_{\ell+d-1}^n X) \mathbf{1}_{\{\Delta_{\ell+m}^n N(y)=0, \Delta_{\ell+m}^n N'=0, m=0, \dots, d-1\}}.$$

Then, $\underline{c}_n(i) \leq \widehat{c}_{n,0}(i)$, and if $y = y_n \rightarrow 0$ as in (B.15), and $(\log n)^3 \ll k_n \ll n$,

$$\max_{1 \leq i \leq n} \frac{\sup_{t \in [\frac{i-k_n/2}{n}, \frac{i+k_n/2}{n})} \sigma_t^2}{\underline{c}_{n,0}(X, i)} \xrightarrow{P} 1. \quad (\text{B.20})$$

In particular, if for some $0 < a < 1$, $n^a \ll k_n \ll n$, then the variables $\underline{c}_n(i; y)$ defined in (A.6) satisfy

$$\max_{1 \leq i \leq n} \frac{\sup_{t \in [\frac{i-k_n/2}{n}, \frac{i+k_n/2}{n})} \sigma_t^2}{\underline{c}_n(i; y)} \xrightarrow{P} 1. \quad (\text{B.21})$$

Proof. Since F is nonnegative, $\underline{c}_n(i) \leq \widehat{c}_{n,0}(i)$ is immediate, so it remains to show (B.20). Consider

$$\underline{c}_{n,0}(\sigma W; \ell) = \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} F(\Delta_i^n(\sigma W), \dots, \Delta_{i+d-1}^n(\sigma W)) \mathbf{1}_{\{\Delta_{i+m}^n N(y)=0, \Delta_{i+m}^n N'=0, 0 \leq m \leq d-1\}}$$

We first show $\max_{1 \leq \ell \leq n} |\underline{c}_{n,0}(\sigma W; \ell) - \sup_{t \in [\frac{\ell-k_n/2}{n}, \frac{\ell+k_n/2}{n})} \sigma_t^2| \xrightarrow{P} 0$. Note

$$\begin{aligned} 0 &\leq \max_{1 \leq \ell \leq n} \left(\widehat{c}_{n,0}(\sigma W; \ell) - \underline{c}_{n,0}(\sigma W; \ell) \right) \\ &\leq \max_{1 \leq \ell \leq n} \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} F(\Delta_i^n(\sigma W), \dots, \Delta_{i+d-1}^n(\sigma W)) \left(\sum_{m=0}^{d-1} (\mathbf{1}_{\{\Delta_{i+m}^n N(y) \neq 0\}} + \mathbf{1}_{\{\Delta_{i+m}^n N' \neq 0\}}) \right). \end{aligned} \quad (\text{B.22})$$

Using (3.6), we have

$$\begin{aligned} &\max_{1 \leq \ell \leq n} \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} F(\Delta_i^n(\sigma W), \dots, \Delta_{i+d-1}^n(\sigma W)) \sum_{m=0}^{d-1} \mathbf{1}_{\{\Delta_{i+m}^n N(y) \neq 0\}} \\ &\leq K \max(\Delta_i^n(\sigma W))^2 \max_{1 \leq \ell \leq n} \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} \sum_{m=0}^{d-1} \mathbf{1}_{\{\Delta_{i+m}^n N(y) \neq 0\}} \\ &= K \frac{\max(\Delta_i^n(\sigma W))^2}{h \log n} \max_{1 \leq \ell \leq n} \frac{\log n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \\ &= O_P(1) \cdot \max_{1 \leq \ell \leq n} \frac{\log n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} \mathbf{1}_{\{\Delta_{i+m}^n N(y) \neq 0\}}. \end{aligned} \quad (\text{B.23})$$

Recalling from (A.1) that $p_n(y) := \mathbb{P}(\Delta_i^n N(y) \neq 0) \sim K h y^{-\alpha} = o(1/(\log n))$ due to (B.15), we have

$$\begin{aligned} &\max_{1 \leq \ell \leq n} \frac{\log n}{k_n} \sum_{i=(\ell-k_n/2+1)}^{(\ell+k_n/2)} \mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} \\ &= o_P(1) + \max_{0 \leq \ell \leq n} \frac{\log n}{k_n} \left| \sum_{i=(\ell-k_n/2+1) \vee 1}^{(\ell+k_n/2) \wedge n} (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} - p_n(y)) \right|. \end{aligned}$$

Hoeffding's inequality together with a union bound give

$$\begin{aligned}
& \mathbb{P} \left(\max_{1 \leq \ell \leq n} \frac{\log n}{k_n} \left| \sum_{i=(\ell-k_n/2+1) \vee 1}^{(\ell+k_n/2) \wedge n} (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} - p_n(y)) \right| \geq \delta \right) \\
& \leq n \mathbb{P} \left(\left| \sum_{i=(\ell-k_n/2+1) \vee 1}^{(\ell+k_n/2) \wedge n} (\mathbf{1}_{\{\Delta_i^n N(y) \neq 0\}} - p_n(y)) \right| \geq \frac{\delta k_n}{\log n} \right) \\
& \leq 2n \exp(-2\delta^2 k_n / \log^2(n)) \\
& \leq 2n \exp(-3 \log n) = o(1),
\end{aligned}$$

since for large enough n , $2\delta^2 k_n / \log^2(n) \gg 3 \log n$ due to the assumption $k_n \gg (\log n)^3$. Hence from (B.23), we obtain

$$\max_{1 \leq \ell \leq n} \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) \sum_{m=0}^{d-1} \mathbf{1}_{\{\Delta_{i+m}^n N(y) \neq 0\}} = o_P(1). \quad (\text{B.24})$$

Turning to the second term in (B.22), since $\sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} \sum_{m=0}^{d-1} \mathbf{1}_{\{\Delta_{i+m}^n N' \neq 0\}} \leq dN'_1$, we have,

$$\begin{aligned}
& \max_{1 \leq \ell \leq n} \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) \sum_{m=0}^{d-1} \mathbf{1}_{\{\Delta_{i+m}^n N' \neq 0\}} \\
& \leq K \max_{1 \leq i \leq n} (\Delta_i^n(\sigma \cdot W))^2 \frac{n}{k_n} \cdot dN'_1 \\
& = O_P(h \log n) \cdot \frac{n}{k_n} \cdot O_P(1) = o_P(1). \quad (\text{B.25})
\end{aligned}$$

Putting together (B.24) and (B.25), by (B.22), we obtain

$$\max_{1 \leq \ell \leq n} (\hat{c}_{n,0}(\sigma \cdot W; \ell) - \underline{c}_{n,0}(\sigma \cdot W; \ell)) = o_P(1).$$

Thus, from (3.14), we obtain

$$\begin{aligned}
& \max_{1 \leq \ell \leq n} \left| \underline{c}_{n,0}(\sigma \cdot W; \ell) - \sup_{t \in [\frac{\ell-k_n/2}{n}, \frac{\ell+k_n/2}{n})} \sigma_t^2 \right| \quad (\text{B.26}) \\
& = o_P(1) + \max_{1 \leq i \leq n} \left| \hat{c}_{n,0}(\sigma \cdot W; i) - \frac{n}{k_n} \int_{\frac{i-k_n/2}{n}}^{\frac{i+k_n/2}{n}} \sigma_t^2 dt \right| \\
& \quad + \max_{1 \leq i \leq n} \left| \frac{n}{k_n} \int_{\frac{i-k_n/2}{n}}^{\frac{i+k_n/2}{n}} \sigma_t^2 dt - \sup_{t \in [\frac{i-k_n/2}{n}, \frac{i+k_n/2}{n})} \sigma_t^2 \right| \\
& = o_P(1),
\end{aligned}$$

where the last term on the second line tends to 0 due to (B.28). On the other hand, we have

$$\begin{aligned}
& \max_{1 \leq \ell \leq n} |\underline{c}_{n,0}(X; \ell) - \underline{c}_{n,0}(\sigma \cdot W; \ell)| \\
& = \left| \max_{1 \leq \ell \leq n} \frac{n}{k_n} \sum_{i=\ell-k_n/2+1}^{\ell+k_n/2} \left[F(\Delta_i^n X, \dots, \Delta_{i+d-1}^n X) - F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) \right] \mathbf{1}_{\{i \in \mathcal{I}_n^{(k)}(y)\}} \right| \\
& \leq n \max_{i \in \mathcal{I}_n^{(d)}} (y) \left| F(\Delta_i^n X, \dots, \Delta_{i+d-1}^n X) - F(\Delta_i^n(\sigma \cdot W), \dots, \Delta_{i+d-1}^n(\sigma \cdot W)) \right| \\
& = o_P(1),
\end{aligned}$$

where the last line follows by repeating the argument leading to (B.19). Thus, using (B.26),

$$\begin{aligned} & \max_{1 \leq \ell \leq n} \left| \mathcal{C}_{n,0}(X; \ell) - \sup_{t \in [\frac{\ell - k_n/2}{n}, \frac{\ell + k_n/2}{n})} \sigma_t^2 \right| \\ & \leq \max_{1 \leq \ell \leq n} \left| \mathcal{C}_{n,0}(X; \ell) - \mathcal{C}_{n,0}(\sigma \cdot W; \ell) \right| + \max_{1 \leq \ell \leq n} \left| \mathcal{C}_{n,0}(\sigma \cdot W; \ell) - \sup_{t \in [\frac{\ell - k_n/2}{n}, \frac{\ell + k_n/2}{n})} \sigma_t^2 \right| = o_P(1), \end{aligned}$$

which establishes (B.20). For the statement (B.21), since $n^a \ll k_n \ll n$, Lemma B.5 implies the initialization $\hat{\mathcal{C}}_{n,0}$ as in (3.13) with $F(x) = x^2$ belongs to class $\mathcal{C}^{\text{spot}}$. Hence, (B.21) follows from (B.20). $\square$

Lemma B.5. *Suppose for some $0 < a < 1$, $n^a \ll k_n \ll n$. Then,*

$$\max_{\frac{k_n}{2} < \ell \leq n - \frac{k_n}{2}} \left| \frac{n}{k_n} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} (\Delta_i^n(\sigma \cdot W))^2 - \sup_{t \in [\frac{\ell - k_n/2}{n}, \frac{\ell + k_n/2}{n})} \sigma_t^2 \right| = o_P(1). \quad (\text{B.27})$$

Proof. Throughout we set $\underline{t}_\ell = \frac{\ell - k_n/2}{n}$ and $\bar{t}_\ell = \frac{\ell + k_n/2}{n}$. Since σ is càdlàg, we claim

$$\max_{\frac{k_n}{n} < \ell \leq n - \frac{k_n}{2}} \left(\sup_{t \in [\underline{t}_\ell, \bar{t}_\ell)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds \right) \rightarrow 0 \quad \text{a.s.} \quad (\text{B.28})$$

Indeed, to show (B.28), for each ω and any $\delta > 0$, we can find a $\nu = \nu(\omega, \delta)$ and $0 = s_0 < s_1(\omega) < \dots < s_\nu(\omega) = 1$ such that

$$\sup_{s, t \in [s_{i-1}, s_i)} |\sigma_s^2 - \sigma_t^2| < \delta, \quad i = 1, \dots, \nu \quad (\text{B.29})$$

(see, e.g., p.122 of [8]). So, take $n_0(\omega)$ large enough so that $n \geq n_0(\omega)$ gives

$$\frac{k_n}{n} < \min \frac{s_i - s_{i-1}}{2}.$$

Then for each ℓ , there is at most one s_j with $\underline{t}_\ell \leq s_j < \bar{t}_\ell$. For any such ℓ , we have

$$\begin{aligned} & \sup_{t \in [\underline{t}_\ell, \bar{t}_\ell)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds \\ & = \max \left\{ \sup_{t \in [\underline{t}_\ell, s_j)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds, \sup_{t \in [s_j, \bar{t}_\ell)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds \right\} \\ & \leq \max \left\{ \sup_{t \in [\underline{t}_\ell, s_j)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 \mathbf{1}_{\{s \in [\underline{t}_\ell, s_j)\}} ds, \sup_{t \in [s_j, \bar{t}_\ell)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 \mathbf{1}_{\{s \in [s_j, \bar{t}_\ell)\}} ds \right\} \\ & = \max \left\{ \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \left(\sup_{t \in [\underline{t}_\ell, s_j)} \sigma_t^2 - \sigma_s^2 \mathbf{1}_{\{s \in [\underline{t}_\ell, s_j)\}} \right) ds, \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \left(\sup_{t \in [s_j, \bar{t}_\ell)} \sigma_t^2 - \sigma_s^2 \mathbf{1}_{\{s \in [s_j, \bar{t}_\ell)\}} \right) ds \right\} \\ & \leq \max \left\{ \sup_{t, s \in [\underline{t}_\ell, s_j)} |\sigma_t^2 - \sigma_s^2|, \sup_{t \in [s_j, \bar{t}_\ell)} |\sigma_t^2 - \sigma_s^2| \right\} < \delta. \end{aligned}$$

For each remaining ℓ , we have $[\underline{t}_\ell, \bar{t}_\ell] \subseteq [s_{j-1}, s_j)$ for some j , giving, for each such ℓ ,

$$0 \leq \sup_{t \in [\underline{t}_\ell, \bar{t}_\ell)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds \leq \sup_{t, s \in [s_{j-1}, s_j)} |\sigma_t^2 - \sigma_s^2| < \delta.$$

Thus, $\max_{\frac{k_n}{n} < \ell \leq n - \frac{k_n}{2}} \left(\sup_{t \in [\underline{t}_\ell, \bar{t}_\ell)} \sigma_t^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds \right) < \delta$, for all $n \geq n_0(\omega)$, which by arbitrariness of δ , gives (B.28).

So, we need only to show

$$\max_{\frac{k_n}{n} < \ell \leq n - \frac{k_n}{2}} \left| \frac{n}{k_n} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} (\Delta_i^n(\sigma \cdot W))^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds \right| = o(1). \quad (\text{B.30})$$

With $Y_t = (\sigma \cdot W)_t$, Itô's formula gives

$$(\Delta_i^n(\sigma \cdot W))^2 = 2 \int_{t_{i-1}}^{t_i} (Y_s - Y_{t_{i-1}}) dY_s + \int_{t_{i-1}}^{t_i} \sigma_s^2 ds. \quad (\text{B.31})$$

Using (B.31), we have

$$\begin{aligned} \max_{\frac{k_n}{n} < \ell \leq n - \frac{k_n}{2}} \left| \frac{n}{k_n} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} (\Delta_i^n(\sigma \cdot W))^2 - \frac{n}{k_n} \int_{\underline{t}_\ell}^{\bar{t}_\ell} \sigma_s^2 ds \right| \\ \leq 2 \max_{\frac{k_n}{n} < \ell \leq n - \frac{k_n}{2}} \left| \frac{n}{k_n} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} \int_{t_{i-1}}^{t_i} (Y_s - Y_{t_{i-1}}) dY_s \right|. \end{aligned}$$

With $g_\ell(s) = \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} (Y_s - Y_{t_{i-1}}) \mathbf{1}_{(t_{i-1}, t_i]}(s)$, since $[Y, Y]_t = \int_0^t \sigma_s^2 ds$, we obtain, for any $p \geq 2$

$$\begin{aligned} \mathbb{E} \left| \frac{n}{k_n} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} \int_{t_{i-1}}^{t_i} (Y_s - Y_{t_{i-1}}) dY_s \right|^p &= \left(\frac{n}{k_n} \right)^p \mathbb{E} \left| \int_0^1 g_\ell(s) dY_s \right|^p \\ &\leq K \left(\frac{n}{k_n} \right)^p \mathbb{E} \left(\int_0^1 g_\ell^2(s) \sigma_s^2 ds \right)^{p/2} \\ &\leq K \left(\frac{n}{k_n} \right)^p \left(\sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} \int_{t_{i-1}}^{t_i} \mathbb{E} |Y_s - Y_{t_{i-1}}|^2 ds \right)^{p/2} \\ &\leq K \left(\frac{n}{k_n} \right)^p k_n^{p/2-1} h^{p/2-1} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} \int_{t_{i-1}}^{t_i} \mathbb{E} |Y_s - Y_{t_{i-1}}|^p ds \\ &\leq K k_n^{-p/2}, \end{aligned}$$

which holds uniformly in ℓ . Thus,

$$\begin{aligned} \mathbb{P} \left(2 \max_{\frac{k_n}{n} < \ell \leq n - \frac{k_n}{2}} \left| \frac{n}{k_n} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} \int_{t_{i-1}}^{t_i} (Y_s - Y_{t_{i-1}}) dY_s \right| > \delta \right) \\ \leq \sum_{\ell = \frac{k_n}{2} + 1}^{n - \frac{k_n}{2}} \mathbb{P} \left(2 \frac{n}{k_n} \left| \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} \int_{t_{i-1}}^{t_i} (Y_s - Y_{t_{i-1}}) dY_s \right| > \delta \right) \\ \leq (\delta/2)^{-p} \sum_{\ell = \frac{k_n}{2} + 1}^{n - \frac{k_n}{2}} \mathbb{E} \left| \frac{n}{k_n} \sum_{i=\ell - \frac{k_n}{2}}^{\ell + \frac{k_n}{2}} \int_{t_{i-1}}^{t_i} (Y_s - Y_{t_{i-1}}) dY_s \right|^p \\ \leq K n k_n^{-p/2}. \end{aligned}$$

By taking $p > 2/a$ we get (B.27). $\square$

Lemma B.6.

$$\limsup_{h \rightarrow 0} \sup_{0 \leq t \leq 1-h} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{2(\sup_{u \in [t, t+h]} \sigma_u^2) h \log(1/h)}} \leq 1, \quad \text{a.s.}$$

In particular, with $\bar{\sigma}_{i,n}^2 = \sup_{t \in [t_{i-1}, t_i]} \sigma_t^2$,

$$\limsup_{n \rightarrow \infty} \max_{i=1, \dots, n} \frac{|\Delta_i^n(\sigma \cdot W)|}{\sqrt{2\bar{\sigma}_{i,n}^2 h_n \log(1/h_n)}} \leq 1, \quad \text{a.s.}$$

Proof. Define

$$D(h) = \sup_{0 \leq t \leq 1-h} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\sup_{u \in [t, t+h]} \sigma_u^2}}.$$

It suffices to show that for every small $\eta > 0$,

$$\limsup_{h \rightarrow 0} D(h) \leq 1 + 3\eta.$$

So, for any $m < 1/h$, define

$$D(m, h) = \max_{j=1, \dots, m} \sup_{t \in [\frac{(j-1)}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\sup_{u \in I_{j,m,h}} \sigma_u^2}},$$

where $I_{j,m,h} = [\frac{(j-1)}{m}, (\frac{j}{m} + h) \wedge 1)$. We first establish for appropriate subsequence $h_\ell \rightarrow 0$, and $m = m(h_\ell) \rightarrow \infty$,

$$\limsup_{\ell \rightarrow \infty} \frac{D(h_\ell, m(h_\ell))}{\sqrt{2h_\ell \log(1/h_\ell)}} \leq 1 + \eta, \quad \text{a.s.}$$

So, let $0 < \eta < 1$ be given, and let $\theta = \frac{(1+\eta)^2}{(1+\eta/2)^2} > 1$. With $C_t = \int_0^t \sigma_s^2 ds$, the Dambis-Dubins-Schwarz Theorem (e.g., Theorem 4.6 in [26]) implies that a.s.,

$$(\sigma \cdot W)_t = B_{C_t}, \quad t \geq 0,$$

where B_t a Brownian motion. Also, since σ is bounded above and below, we may take a $J, J_0 > 0$ such that $\theta^J \geq \bar{\sigma}^2$ and $\theta^{-J_0} < \inf \sigma_t^2$ a.s. For any $-J_0 \leq \ell \leq J$, on the event $\{\sup_{u \in I_{j,m,h}} \sigma_u^2 \in$

$(\theta^{\ell-1}, \theta^\ell]$ observe $C_t \leq u, v \leq C_{t+h}$ implies $|u - v| \leq h\theta^\ell$ for each $t \in [\frac{(j-1)}{m}, \frac{j}{m})$. This gives

$$\begin{aligned}
& \mathbb{P} \left(\sup_{t \in [\frac{(j-1)}{m}, \frac{j}{m})} \frac{|B_{C_{t+h}} - B_{C_t}|}{\sqrt{2(\sup_{u \in I_{j,m,h}} \sigma_u^2) h \log(1/h)}} > 1 + \eta, \sup_{u \in I_{j,m,h}} \sigma_u^2 \in (\theta^{\ell-1}, \theta^\ell] \right) \\
& \leq \mathbb{P} \left(\sup_{t \in [\frac{(j-1)}{m}, \frac{j}{m})} \frac{\sup_{C_t \leq u, v \leq C_{t+h}} |B_u - B_v|}{\sqrt{2(\sup_{u \in I_{j,m,h}} \sigma_u^2) h \log(1/h)}} > 1 + \eta, \sup_{u \in I_{j,m,h}} \sigma_u^2 \in (\theta^{\ell-1}, \theta^\ell] \right) \\
& \leq \mathbb{P} \left(\sup_{|u-v| \leq h\theta^\ell} \sup_{0 \leq u, v \leq \theta^J} \frac{|B_u - B_v|}{\sqrt{2\theta^{\ell-1} h \log(1/h)}} > 1 + \eta \right) \\
& = \mathbb{P} \left(\sup_{|u-v| \leq h} \sup_{0 \leq s, t \leq \theta^J / \theta^\ell} \frac{\sqrt{\theta^\ell} |B_s - B_t|}{\sqrt{2\theta^{\ell-1} h \log(1/h)}} > 1 + \eta \right) \\
& \leq \mathbb{P} \left(\sup_{|s-t| \leq h} \sup_{0 \leq s, t \leq \theta^{J-\ell}} \frac{|B_s - B_t|}{\sqrt{2h \log(1/h)}} > (1 + \eta)/\sqrt{\theta} \right) \\
& \leq \mathbb{P} \left(\sup_{|s-t| \leq h} \sup_{0 \leq s, t \leq \theta^{J+J_0}} \frac{|B_s - B_t|}{\sqrt{2h \log(1/h)}} > (1 + \eta)/\sqrt{\theta} \right) \\
& = \mathbb{P} \left(\sup_{|s-t| \leq h} \sup_{0 \leq s, t \leq \theta^{J+J_0}} \frac{|B_s - B_t|}{\sqrt{2h \log(1/h)}} > 1 + \eta/2 \right), \tag{B.32}
\end{aligned}$$

where on the 4th line we used self-similarity of B_t . Summing over ℓ and applying a union bound we obtain

$$\mathbb{P} \left(\frac{D(h, m)}{\sqrt{2h \log(1/h)}} > 1 + \eta \right) \leq Km \mathbb{P} \left(\sup_{|s-t| \leq h} \sup_{0 \leq s, t \leq \theta^{J+J_0}} \frac{|B_t - B_s|}{\sqrt{2h \log(1/h)}} > 1 + \eta/2 \right). \tag{B.33}$$

Now, recall (c.f. Lemma 1.1.1 in [12]), that for every $M, \delta > 0$, there exists a $K_0 = K_0(\delta, M)$ such that for every positive $v > 0$,

$$\mathbb{P} \left(\sup_{0 \leq s, t \leq M, |s-t| \leq h} |B_t - B_s| \geq v\sqrt{h} \right) \leq \frac{K_0}{h} \exp \left\{ \frac{-v^2}{2 + \delta} \right\}.$$

So, taking $\delta > 0$ small enough so that $\frac{2(1+\eta/2)^2}{2+\delta} = 1 + \kappa$ for some $0 < \kappa < 1$, and taking $M = \theta^{J+J_0}$, and $v = (1 + \eta/2)\sqrt{2 \log(1/h)}$, we get

$$\begin{aligned}
\mathbb{P} \left(\sup_{|s-t| \leq h, 0 \leq s, t \leq \theta^{J+J_0}} \frac{|B_t - B_s|}{\sqrt{2h \log(1/h)}} > 1 + \eta/2 \right) & \leq \frac{K_0}{h} \exp \left\{ \frac{-2(1 + \eta/2)^2 \log(1/h)}{2 + \delta} \right\} \\
& \leq K_0 h^\kappa \tag{B.34}
\end{aligned}$$

Thus, with $m = m(h) = \lfloor h^{-\kappa/2} \rfloor$, combining (B.33) and (B.34) we get

$$\mathbb{P} \left(\frac{D(h, m(h))}{\sqrt{2h \log(1/h)}} > 1 + \eta \right) \leq Km h^\kappa \leq K h^{\kappa/2}. \tag{B.35}$$

Now, for some integer $r_0 > 2/\kappa$ consider the subsequence h_ℓ defined by

$$h_\ell = \ell^{-r_0}.$$

Using (B.35) we obtain

$$\sum_{\ell=2}^{\infty} \mathbb{P} \left(\frac{D(h_\ell, m(h_\ell))}{\sqrt{2h_\ell \log(1/h_\ell)}} > 1 + \eta \right) \leq \sum_{\ell=2}^{\infty} \ell^{-(r_0 \kappa/2)} < \infty.$$

Thus, the Borel-Cantelli lemma gives

$$\limsup_{\ell \rightarrow \infty} \frac{D(h_\ell, m(h_\ell))}{\sqrt{2h_\ell \log(1/h_\ell)}} \leq 1 + \eta, \quad \text{a.s.}$$

This implies for a.s. ω we can find a large $\ell_0(\omega)$ such that

$$\frac{D(h_\ell, m(h_\ell))}{\sqrt{2h_\ell \log(1/h_\ell)}} < 1 + 2\eta, \quad \ell \geq \ell_0(\omega). \quad (\text{B.36})$$

We now proceed to establish, for some $\ell_1(\omega) < \infty$, a.s.,

$$\frac{D(h_\ell)}{\sqrt{2h_\ell \log(1/h_\ell)}} \leq \left(\frac{1 + 3\eta}{1 + 2\eta} \right) \frac{D(h_\ell, m(h_\ell))}{\sqrt{2h_\ell \log(1/h_\ell)}}, \quad \ell \geq \ell_1(\omega). \quad (\text{B.37})$$

Recall (c.f. (B.29)) we may take $\nu = \nu(\omega)$ large enough so that for all $u_1 \in [s_0, s_1), \dots, u_\nu \in [s_{\nu-1}, s_\nu)$,

$$\max_{i=1, \dots, \nu} \sup_{t \in [s_{i-1}, s_i)} \sigma_t^2 - \sigma_{u_i}^2 \leq \delta' \theta^{-J_0},$$

where $\delta' > 0$ is chosen small enough so that $\sqrt{1 - \delta'} = \frac{1+2\eta}{1+3\eta}$. This gives, for any $u \in [s_{i-1}, s_i)$,

$$\begin{aligned} \sigma_u^2 &= \sup_{t \in [s_{i-1}, s_i)} \sigma_t^2 - \left(\sup_{t \in [s_{i-1}, s_i)} \sigma_t^2 - \sigma_u^2 \right) \\ &\geq \sup_{t \in [s_{i-1}, s_i)} \sigma_t^2 - \delta' \theta^{-J_0} \\ &\geq (1 - \delta') \sup_{t \in [s_{i-1}, s_i)} \sigma_t^2, \end{aligned}$$

since $\sup_{t \in I_i} \sigma_t^2 \geq \inf_t \sigma_t^2 \geq \theta^{-J_0}$. So, taking $m_0(\omega)$ large enough so that

$$\frac{2}{m_0(\omega)} \leq \min_{i=1 \dots \nu} s_i - s_{i-1},$$

for every $m \geq m_0(\omega)$ there is at most one s_i in each interval $I_{j,m,h} \subseteq [\frac{j-1}{m}, \frac{j+1}{m})$. In the case that for some i , $I_{j,m,h} \subseteq [s_{i-1}, s_i)$, we have

$$\sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\sup_{u \in [t, t+h)} \sigma_u^2}} \leq \sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{(1 - \delta') \sup_{u \in I_{j,m,h}} \sigma_u^2}}.$$

In the other case, i.e. for some i , $s_i \in I_{j,m,h}$ (i.e., $s_{i-1} < \frac{j-1}{m} < s_i < (\frac{j}{m} + h) \wedge 1 < s_{i+1}$), we have:

$$\begin{aligned} &\sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\sup_{u \in [t, t+h)} \sigma_u^2}} \\ &= \sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\max\{\sup_{u \in [t, (t+h) \wedge s_i)} \sigma_u^2, \sup_{u \in [(t+h) \wedge s_i, t+h)} \sigma_u^2\}}} \\ &\leq \sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\max\{(1 - \delta') \sup_{u \in [\frac{j-1}{m}, s_i)} \sigma_u^2, (1 - \delta') \sup_{u \in [s_i, (\frac{j}{m} + h) \wedge 1)} \sigma_u^2\}}} \\ &= \sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{(1 - \delta') \sup_{u \in I_{j,m,h}} \sigma_u^2}} \end{aligned}$$

Hence, for all $m \geq m_0(\omega)$, and all $h < \frac{1}{m}$,

$$\begin{aligned}
D(h) &= \sup_{0 \leq t \leq 1-h} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\sup_{u \in [t, t+h]} \sigma_u^2}} \\
&= \max_{j=1, \dots, m} \sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\sup_{u \in [t, t+h]} \sigma_u^2}} \\
&\leq \frac{1}{\sqrt{1-\delta'}} \max_{j=1, \dots, m} \sup_{t \in [\frac{j-1}{m}, \frac{j}{m} \wedge (1-h))} \frac{|\int_t^{t+h} \sigma_u dW_u|}{\sqrt{\sup_{u \in [\frac{j-1}{m}, \frac{j}{m}] \sigma_u^2}} \\
&= \frac{1+3\eta}{1+2\eta} D(h, m).
\end{aligned}$$

So, if we choose $\ell_1(\omega) \geq \ell_0(\omega)$ large enough so that $m(h_\ell) > m_0(\omega)$ for all $\ell \geq \ell_1(\omega)$, expression (B.37) holds. From (B.36) we then have

$$\frac{D(h_\ell)}{\sqrt{2h_\ell \log(1/h_\ell)}} \leq \left(\frac{1+3\eta}{1+2\eta} \right) \frac{D(h_\ell, m(h_\ell))}{\sqrt{2h_\ell \log(1/h_\ell)}} < 1+3\eta, \quad \ell \geq \ell_1(\omega),$$

i.e.,

$$\limsup_{\ell \rightarrow \infty} \frac{D(h_\ell)}{\sqrt{2h_\ell \log(1/h_\ell)}} \leq 1+3\eta, \quad \text{a.s.}$$

Now, for any h with $h_{\ell+1} \leq h < h_\ell$, we have

$$\frac{D(h)}{2h \log(1/h)} \leq \frac{D(h_\ell)}{\sqrt{2h_{\ell+1} \log(1/h_{\ell+1})}} = \frac{D(h_\ell)}{\sqrt{2h_\ell \log(1/h_\ell)}} \frac{\sqrt{2h_\ell \log(1/h_\ell)}}{\sqrt{2h_{\ell+1} \log(1/h_{\ell+1})}}$$

which gives

$$\limsup_{h \rightarrow 0} \frac{D(h)}{2h \log(1/h)} \leq \limsup_{\ell \rightarrow \infty} \frac{D(h_\ell)}{\sqrt{2h_\ell \log(1/h_\ell)}} \limsup_{\ell \rightarrow \infty} \frac{\sqrt{2h_\ell \log(1/h_\ell)}}{\sqrt{2h_{\ell+1} \log(1/h_{\ell+1})}} \leq 1+3\eta, \quad \text{a.s.},$$

which completes the proof. $\square$

References

- [1] Y. Aït-Sahalia and J. Jacod. Estimating the degree of activity of jumps in high frequency data. *Annals of Statistics*, 37(5A):2202–2244, 2009.
- [2] Y. Aït-Sahalia and J. Jacod. Testing for jumps in a discretely observed process. *The Annals of Statistics*, 37(1):184–222, 2009.
- [3] Y. Aït-Sahalia and J. Jacod. Testing whether jumps have finite or infinite activity. *The Annals of Statistics*, 39(3), 2011.
- [4] T. G. Andersen, D. Dobrev, and E. Schaumburg. Jump-robust volatility estimation using nearest neighbor truncation. *Journal of Econometrics*, 169(1):75–93, 2012.
- [5] O. E. Barndorff-Nielsen. Power and bipower variation with stochastic volatility and jumps. *Journal of Financial Econometrics*, 2(1):1–37, 2004.
- [6] O. E. Barndorff-Nielsen and N. Shephard. Realized power variation and stochastic volatility models. *Bernoulli*, 9(2), 2003.

- [7] O. E. Barndorff-Nielsen, N. Shephard, and M. Winkel. Limit theorems for multipower variation in the presence of jumps. *Stochastic Processes and their Applications*, 116(5):796–806, 2006.
- [8] P. Billingsley. *Convergence of Probability Measures*. Wiley, 2nd edition, 1999.
- [9] B. C. Boniece, J. E. Figueroa-López, and Y. Han. Efficient integrated volatility estimation in the presence of infinite variation jumps via debiased truncated realized variations, 2022. arXiv preprint, arXiv:2209.10128.
- [10] R. Cont and C. Mancini. Nonparametric tests for pathwise properties of semimartingales. *Bernoulli*, 17(2):781–813, 2011.
- [11] F. Corsi, D. Pirino, and R. Renò. Threshold bipower variation and the impact of jumps on volatility forecasting. *Journal of Econometrics*, 159(2):276–288, 2010.
- [12] M. Csörgö and P. Révész. *Strong Approximations in Probability and Statistics*. Probability and Mathematical Statistics. Acad. Press, 1981.
- [13] Y. Ding, Y. Li, G. Liu, and X. Zheng. Stock co-jump networks. *Journal of Econometrics*, page 105420, 2023.
- [14] J. Fan and Y. Wang. Spot volatility estimation for high-frequency data. *Statistics and Its Interface*, 1(2):279–288, 2008.
- [15] J. E. Figueroa-López, R. Gong, and Y. Han. Estimation of tempered stable Lévy models of infinite variation. *Methodology and Computing in Applied Probability*, 24(2):713–747, June 2022.
- [16] J. E. Figueroa-López and C. Li. Optimal kernel estimation of spot volatility of stochastic differential equations. *Stochastic Processes and their Applications*, 130(8):4693–4720, 2020.
- [17] J. E. Figueroa-López, C. Li, and J. Nisen. Optimal iterative threshold-kernel estimation of jump diffusion processes. *Statistical Inference for Stochastic Processes*, 23(3):517–552, 2020.
- [18] J. E. Figueroa-López and C. Mancini. Optimum thresholding using mean and conditional mean squared error. *Journal of Econometrics*, 208(1):179–210, 2019.
- [19] J. E. Figueroa-López and J. Nisen. Optimally thresholded realized power variations for Lévy jump diffusion models. *Stochastic Processes and their Applications*, 123(7):2648–2677, 2013.
- [20] J. E. Figueroa-López and J. Nisen. Second-order properties of thresholded realized power variations of FJA additive processes. *Statistical Inference for Stochastic Processes*, 22(3):431–474, 2019.
- [21] J. E. Figueroa-López and B. Wu. Kernel estimation of spot volatility with microstructure noise using pre-averaging. *Econometric Theory*, pages 1–50, 2022.
- [22] J. Jacod. Asymptotic properties of realized power variations and related functionals of semimartingales. *Stochastic Processes and their Applications*, 118(4):517–559, 2008.
- [23] J. Jacod and P. Protter. *Discretization of Processes*. Springer, 2011.
- [24] J. Jacod and V. Todorov. Efficient estimation of integrated volatility in presence of infinite variation jumps. *The Annals of Statistics*, 42(3):1029–1069, 2014.
- [25] J. Jacod and V. Todorov. Limit theorems for integrated local empirical characteristic exponents from noisy high-frequency data with application to volatility and jump activity estimation. *The Annals of Applied Probability*, 28(1), 2018.
- [26] I. Karatzas and S. E. Shreve. *Brownian Motion and Stochastic Calculus*. Springer, 1998.

- [27] R. Kawai. On sequential calibration for an asset price model with piecewise Lévy processes. *IAENG International Journal of Applied Mathematics*, 40(4):1–8, 2010.
- [28] D. Kristensen. Nonparametric filtering of the realized spot volatility: a kernel-based approach. *Econometric Theory*, 26(1):60–93, Feb. 2010.
- [29] J. Li, V. Todorov, and G. Tauchen. Jump regressions. *Econometrica*, 85(1):173–195, 2017.
- [30] C. Mancini. Disentangling the jumps of the diffusion in a geometric jumping Brownian motion. *Gornale dell’Istituto Italiano degli Attuari*, 64:19–47, 2001.
- [31] C. Mancini. Non-parametric threshold estimation for models with stochastic diffusion coefficient and jumps. *Scandinavian Journal of Statistics*, 36(2):270–296, 2009.
- [32] C. Mancini. The speed of convergence of the threshold estimator of integrated variance. *Stochastic Processes and their Applications*, 121(4):845–855, 2011.
- [33] C. Mancini. Truncated realized covariance when prices have infinite variation jumps. *Stochastic Processes and their Applications*, 127(6):1998–2035, 2017.
- [34] C. Mancini and F. Gobbi. Identifying the Brownian covariation from the co-jumps given discrete observations. *Econometric Theory*, 28(2):249–273, 2012.
- [35] C. Mancini and R. Renò. Threshold estimation of Markov models with jumps and interest rate modeling. *Journal of Econometrics*, 160(1):77–92, 2011.
- [36] C. Palmes and J. H. C. Woerner. A mathematical analysis of the Gumbel test for jumps in stochastic volatility models. *Stochastic Analysis and Applications*, 34(5):852–881, 2016.
- [37] K.-i. Sato. *Lévy Processes and Infinitely Divisible Distributions*. Cambridge University Press, 1999.
- [38] V. Todorov and G. Tauchen. The realized Laplace transform of volatility. *Econometrica*, 80(3):1105–1127, 2012.