

ON THE SUBADDITIVITY OF GENERALIZED KODAIRA DIMENSIONS

BOJIE HE AND XIANGYU ZHOU

ABSTRACT. The goals of this paper are of two aspects. Firstly, we introduce the notion of generalized numerical Kodaira dimension with multiplier ideal sheaf and establish the subadditivity inequalities in terms of this notion, which can be used to give an analytic proof of O. Fujino's result on the subadditivity of the log Kodaira dimensions. Secondly, motivated by Zhou-Zhu's subadditivity of generalized Kodaira dimensions, we adopt another definition of generalized Kodaira dimension with multiplier ideal sheaf and show they are equal by using Okounkov bodies. As one application, we show that the superadditivity part in Zhou-Zhu's setting also holds true. As another application, we give an alternative proof of Zhou-Zhu's subadditivity formula, in the case when the singular metric h_L has analytic singularities, by using generalized Iitaka fibrations.

1. INTRODUCTION AND MAIN RESULTS

1.1. Introduction and background. Let X and Y be two compact connected complex manifolds of dimension n and m ($n \geq m$). In the analytic setting, we call $f : X \rightarrow Y$ an **(analytic) fiber space**, if f is a surjective holomorphic mapping with connected fibers. Moreover, when X and Y are supposed to be projective manifolds (resp. Kähler manifolds), the fiber space $f : X \rightarrow Y$ is usually called an **algebraic fiber space** (resp. **Kähler fiber space**).

Let

$$\kappa(X) := \max\{\nu \in \mathbb{Z} \mid \limsup_{k \rightarrow \infty} \frac{h^0(X, \mathcal{O}_X(kK_X))}{k^\nu} > 0\} \in \{-\infty, 0, 1, \dots, n\}$$

be the Kodaira dimension of X , where K_X is the canonical divisor. When $\kappa(X) = \dim X = n$, X is said to be of general type.

Conjecture 1.1 (Conjecture $C_{n,m}$ or Iitaka conjecture). Let $f : X \rightarrow Y$ be an algebraic fiber space between two projective manifolds X and Y . Let F denote the general fiber of f , then

$$(1.1) \quad \kappa(X) \geq \kappa(F) + \kappa(Y).$$

Note that when X is not assumed to be Kähler, inequality (1.1) does not always hold in general (cf. Remark 15.3 of [32]).

In [33], E. Viehweg proposed a stronger version of Iitaka conjecture:

2020 *Mathematics Subject Classification.* 32J25, 14C30, 32L10, 32U05, 14M25, 14F18.

Key words and phrases. multiplier ideal sheaf, generalized Kodaira-Iitaka dimension, Iitaka fibration, Okounkov body, fibre spaces.

The second author is partially supported by the National Natural Science Foundation of China (grant no. 11688101 and no. 12288201).

Conjecture 1.2 (Conjecture $C_{n,m}^+$ or generalized Iitaka conjecture). Let $f : X \rightarrow Y$ be an algebraic fiber space between two projective manifolds X and Y . Let F denote the general fiber of f , then

$$(1.2) \quad \kappa(X) \geq \kappa(F) + \max\{\text{Var}(f), \kappa(Y)\}.$$

Here $\text{Var}(f)$, roughly speaking, stands for the dimension of the birational equivalence class of all the general fibers F .

Let L (resp. D) be a \mathbb{Q} -line bundle (resp. \mathbb{Q} -Cartier divisor) over a compact (connected) complex manifold X , its **Iitaka L (resp. D)-dimension** $\kappa(X, L)$ (resp. $\kappa(X, D)$) is usually defined as

$$\begin{aligned} \kappa(X, L) &:= \max\{\nu \in \mathbb{Z} \bigcup \{-\infty\}; \limsup_{k \rightarrow \infty} \frac{h^0(X, kk_0 L)}{k^\nu} > 0\} \in \{-\infty, 0, \dots, \dim X\} \\ (\text{resp. } \kappa(X, D) &:= \max\{\nu \in \mathbb{Z} \bigcup \{-\infty\}; \limsup_{k \rightarrow \infty} \frac{h^0(X, \mathcal{O}_X(kk_0 D))}{k^\nu} > 0\} \\ &\in \{-\infty, 0, \dots, \dim X\}), \end{aligned}$$

where k_0 is the smallest integer such that $k_0 L$ (resp. $\mathcal{O}_X(k_0 D)$) is a line bundle. One may see in section 4 for some basic and further explanations of this notion.

Assume that X is projective and let A be an ample line bundle on X . By perturbing all sections of $kk_0 L$ with A , Nakayama's **numerical dimension** of D or $L = \mathcal{O}(D)$ (cf. Definition 2.7 in [15], Chapter V in [27] or Definition 2.4.8 in [12]) is defined as

$$\kappa_\sigma(X, L) := \max_{E: \text{a divisor of } X} \{\kappa_\sigma(X, L; E)\} = \max_{m \in \mathbb{N}} \{\kappa_\sigma(X, L; mA)\},$$

where $\kappa_\sigma(X, L; E)$ is defined to be

$$\max\{\nu \in \mathbb{Z} \cup \{-\infty\}; \limsup_{k \rightarrow \infty} \frac{h^0(X, kk_0 L + E)}{k^\nu} > 0\}.$$

There are some basic properties about κ and κ_σ . For example, $\kappa(X, L) \geq 0$ if and only if $m_0 L$ is effective for some $m_0 \in \mathbb{N}$, $\kappa_\sigma(X, L) \geq 0$ if and only if L is pseudoeffective. Some other properties are e.g. $\kappa(X, L) \leq \kappa_\sigma(X, L)$, $\kappa_\sigma(X, L)$ equals the numerical dimension $\nu(X, L)$ when L is nef, etc. To see other various useful characterizations of κ and κ_σ , one may refer to [27] or [12].

In case when $L = K_X$ is the canonical bundle, $\kappa(X, K_X)$ (resp. $\kappa_\sigma(X, K_X)$) is called the Kodaira dimension (resp. numerical Kodaira dimension) of X .

When L is moreover assumed to be pseudoeffective, i.e. which can be equipped with a singular metric h_L whose curvature current is semi-positive, one may similarly define the **generalized Kodaira dimension** (cf. Definition 1.1 in [40]) $\kappa(X, K_X + L, h_L)$ as

$$(1.3) \quad \max\{\nu \in \mathbb{Z} \bigcup \{-\infty\}; \limsup_{k \rightarrow \infty} \frac{h^0(X, kk_0(K_X + L) \otimes \mathcal{I}_{kk_0}(h_L))}{k^\nu} > 0\}$$

and the **generalized numerical Kodaira dimension** $\kappa_\sigma(X, K_X + L, h_L)$ as

$$\max_{E: \text{a divisor of } X} \{\kappa_\sigma(X, K_X + L, h_L; E)\} = \max_{k \in \mathbb{N}} \{\kappa_\sigma(X, K_X + L, h_L; kA)\},$$

where $\kappa_\sigma(X, K_X + L, h_L; E)$ is defined to be

$$(1.4) \quad \max\{\nu \in \mathbb{Z} \bigcup \{-\infty\}; \limsup_{k \rightarrow \infty} \frac{h^0(X, (kk_0(K_X + L) + E) \otimes \mathcal{I}_{kk_0}(h_L))}{k^\nu} > 0\}.$$

Here the kk_0 -multiplier ideal sheaves $\mathcal{I}_{kk_0}(h_L)$ are precisely defined as in (2.1).

The definition of generalized Kodaira dimensions with multiplier ideal sheaves is motivated by the study of the graded subalgebra $S = \bigoplus_{k \in \mathbb{N}} S_k \subset R(X, K_X + L)$ with $S_k = H^0(X, kk_0(K_X + L) \otimes \mathcal{I}_{kk_0}(h_L))$ (cf. (4.6)), according to the very basic relationship $\mathcal{I}_k(h_L)\mathcal{I}_l(h_L) \subset \mathcal{I}_{k+l}(h_L)$ (which is due to Hölder inequality) for $k, l \in \mathbb{N}$.

1.2. Subadditivity of generalized numerical Kodaira dimension. The following result obtained by O. Fujino is mainly motivated by the study of the subadditivity of the logarithmic Kodaira dimension of algebraic varieties, which is denoted by Conjecture $\overline{C}_{n,m}$ (see its precise form in e.g. Conjecture 1.2.2 in [13]). Note that Conjecture $C_{n,m}$ is a special case of Conjecture $\overline{C}_{n,m}$.

Theorem 1.1. (Theorem 1.3 in [15] or Theorem 2.1 in [16]) *Let $f : X \rightarrow Y$ be an algebraic fiber space between two projective manifolds X and Y . Let D_X (resp. D_Y) be a simple normal crossing divisor (see Definition 9.1.7 in [24]) on X (resp. Y). Assume that $f^*D_Y \subset D_X$, then*

$$(1.5) \quad \kappa_\sigma(X, K_X + D_X) \geq \kappa_\sigma(F, K_F + D_F) + \kappa(Y, K_Y + D_Y)$$

and

$$(1.6) \quad \kappa_\sigma(X, K_X + D_X) \geq \kappa(F, K_F + D_F) + \kappa_\sigma(Y, K_Y + D_Y),$$

where F is a sufficiently general fiber of $f : X \rightarrow Y$.

Thus Conjecture $\overline{C}_{n,m}$ follows immediately from the generalized abundance conjecture (cf. Conjecture 2.10 in [15]), which implies that

$$\kappa(X, K_X + D_X) = \kappa_\sigma(X, K_X + D_X)$$

whenever D_X is a simple normal crossing divisor on the projective manifold X .

Our first goal will be to generalize the above theorem with multiplier ideal sheaves involved. To this end, let us recall that the key ingredient of the original proof of Theorem 1.1 gives a generically globally generated proposition of direct image sheaves (cf. Corollary 3.7 in [15] or Theorem 3.35 of Chapter V in [27]). His proof of this property is based on Nakayama's theory of ω -sheaves, which is closely related to Viehweg's covering trick and weak positivity ([33], [34]).

To convert this important (general) global generation property into a purely analytic statement (e.g. see Lemma 2.4 and Lemma 2.5), we present the following theorem, which is an application of Ohsawa-Takegoshi L^2 extension theorem on weakly pseudoconvex Kähler manifolds (cf. Theorem 2.3 and Remark 2.3). One might also compare it with Siu's uniform global generation formula on pseudoeffective line bundles (cf. Chapter 6.E in [6]).

Theorem 1.2. *Let $f : X \rightarrow Y$ be an algebraic fiber space between two projective manifolds X and Y , $L \rightarrow X$ a line bundle equipped with a singular hermitian metric h_L whose curvature current is semi-positive. Then there exist two ample line bundles $G \rightarrow Y$ and $H \rightarrow X$ such that*

$$(1.7) \quad \mathcal{F}_1 := \mathcal{O}_Y(G) \otimes f_*\mathcal{O}_X((kK_{X/Y} + kL) \otimes \mathcal{I}_k(h_L))$$

is sufficiently generically globally generated and

$$(1.8) \quad \mathcal{F}_2 := \mathcal{O}_Y(G) \otimes f_*\mathcal{O}_X(kK_{X/Y} + kL + H)$$

is generically globally generated for any integer $k \geq 1$.

Here by saying a coherent sheaf \mathcal{F} is “*generically globally generated*” (resp. “*sufficiently generically globally generated*”), we mean \mathcal{F} globally generates its stalks on a Zariski open dense subset (resp. on a subset whose complement has measure zero).

Theorem 1.2 will be obtained through two extension statements: Lemma 2.4 with multiplier ideal sheaves and Lemma 2.5 without multiplier ideal sheaves. As an immediate application of Theorem 1.2 (or more precisely, Lemma 2.4), we obtain an analogue of Theorem 1.1 for the subadditivity of generalized numerical Kodaira dimensions.

Theorem 1.3 (Main Theorem 1). *Let $f : X \rightarrow Y$ be an algebraic fiber space between two projective manifolds X and Y . Let L be a \mathbb{Q} -line bundle on X equipped with a singular metric h_L whose curvature current is semi-positive. Then inequality*

$$(1.9) \quad \kappa_\sigma(X, K_X + L, h_L) \geq \kappa_\sigma(F, K_F + L_F, h_L|_F) + \kappa(Y, K_Y)$$

and

$$(1.10) \quad \kappa_\sigma(X, K_X + L, h_L) \geq \kappa(F, K_F + L_F, h_L|_F) + \kappa_\sigma(Y, K_Y)$$

holds, where F is a sufficiently general fiber of $f : X \rightarrow Y$.

By “*sufficiently general*”, we mean this inequality holds for $F = X_y$ when y varies outside a measure zero subset of Y . In terms of (1.10), we will see from its proof that the left hand side κ_σ can be optimized by a smaller number, which is defined to be the generalized numerical Kodaira dimension along the horizontal direction, i.e.

$$(1.11) \quad \kappa_{\sigma, f, \text{hor}} := \max_{E: \text{a divisor of } Y} \{ \kappa_\sigma(X, K_X + L, h_L; f^*E) \} \leq \kappa_\sigma.$$

Let us remark that $\kappa_\sigma(F, K_F + L|_F, h_L|_F)$ is invariant for sufficiently general F . Indeed, for any $k, l \in \mathbb{N}$, there exists Zariski open dense subsets $Y_{k,l} \subset Y$ such that $h^0(X_y, k k_0(X_y + L|_{X_y}) \otimes l A \otimes \mathcal{I}_{k k_0}(h_L)|_{X_y})$ is invariant when $y \in Y_{k,l}$ (e.g. Theorem 10.7 in [3]). As $\mathcal{I}_{k k_0}(h_L)|_{X_y} = \mathcal{I}_{k k_0}(h_L|_{X_y})$ holds for $y \in \Sigma_k$, where $Y \setminus \Sigma_k$ is a subset with zero measure, it follows that $\kappa_\sigma(F, K_F + L_F, h_L|_F)$ is constant on $\bigcap_{k,l=1}^\infty (Y_{k,l} \cap \Sigma_k)$. Note also by almost the same argument, $\kappa(F, K_F + L|_F, h_L|_F)$ does not vary for almost all F . One can refer to Remark 1.1 in [40] or Section 2.1 for further explanations of this result.

At the end of the proof of Theorem 1.3, we will provide an analytic approach towards Theorem 1.1 (cf. Remark 3.2), which can be respectively viewed as an application of Theorem 1.2 (or more precisely, Lemma 2.5).

1.3. Subadditivity of generalized Kodaira dimension. The second goal of this paper is to give another interpretation of the subadditivity of generalized Kodaira dimension, which is mainly motivated by a recent result of Zhou and Zhu.

Theorem 1.4. (Theorem 1.1 in [40]) *Let $f : X \rightarrow Y$ be an analytic fiber space between a compact Kähler manifold X and a connected compact complex manifold Y . Let $L \rightarrow X$ be a \mathbb{Q} -line bundle equipped with a singular metric h_L with semi-positive curvature current. Assume that Y has general type, then inequality*

$$(1.12) \quad \kappa(X, K_X + L, h_L) \geq \kappa(F, K_F + L|_F, h_L|_F) + \dim Y$$

holds, where F denotes the sufficiently general fiber of f .

Remark 1.1. As a slight improvement to Theorem 1.4, we can show that (1.12) is indeed an equality without any singularity restrictions on h_L (cf. Proposition 5.1).

Note that in [40] $\kappa(X, K_X + L, h_L)$ is defined (cf. 4.19) to be the largest order $m \in \mathbb{N} \cup \{-\infty\}$, where some subsequence of $\{h^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L))\}_{k=1}^\infty$ grows as $O(k^m)$, $k \rightarrow \infty$. Essentially, Zhou and Zhu's proof of Theorem 1.4 is mainly based on two ingredients: Demailly's L^2 extension theory on weakly pseudoconvex manifolds ([8]) and their results on positivity of relative k -Bergman metrics for Kähler fibrations ([39]).

However, we will try **NOT** to adopt the definition of generalized Kodaira dimension along this direction. Instead, we define $\tilde{\kappa}(X, K_X + L, h_L)$ (cf. (4.18)) to be the maximal dimension of the image of the meromorphic Kodaira map corresponding to the linear system

$$(1.13) \quad |H^0(X, kk_0(K_X + L) \otimes \mathcal{I}_{kk_0}(h_L))|$$

(cf. 4.18) as k varies among all integers. We will show that both $\kappa(X, K_X + L, h_L)$ and $\tilde{\kappa}(X, K_X + L, h_L)$ coincide with the dimension of an appropriate Newton-Okounkov body (cf. Proposition 4.14), even when X is possibly non-algebraic, which will be crucial for the improvement of Theorem 1.4 as mentioned in Remark 1.1. Note in the usual setting (i.e. set (L, h_L) to be the trivial line bundle equipped with the complex Euclidean metric), the fact that $\kappa(X, K_X) = \tilde{\kappa}(X, K_X)$ has already been proved in [26]; the fact that both $\kappa(X, K_X)$ and $\tilde{\kappa}(X, K_X)$ coincide with an appropriate Newton-Okounkov body in case X is algebraic has been proved in (Theorem 3.3 and 3.4 of) [22].

Before turning to the second main results, let us talk about the singularities of h_L . Recall that h_L is said to be of **analytic singularities**, if the local weight function of h_L has the form $c \cdot \log(|F_1|^2 + \cdots + |F_N|^2) + O(1)$, where the F_j are all local holomorphic functions and $c > 0$.

Now let us first fix the notation $S_k = H^0(X, kk_0(K_X + L) \otimes \mathcal{I}_{kk_0}(h_L))$. Then the second main result of this paper, which guarantees the existence of generalized Itaka fibration when h_L has analytic singularities, can be stated as follows.

Theorem 1.5 (=Theorem 4.15, **Main Theorem 2**). *Let X be a compact complex manifold and $L \rightarrow X$ be a pseudoeffective line bundle equipped with a singular metric h_L whose curvature current is semi-positive. Assume that h_L has analytic singularities and $\kappa(X, K_X + L, h_L) \geq 0$. Let $f^b : X^b \rightarrow Y^b$ be a representative of the Kodaira meromorphic mapping*

$$(1.14) \quad f(= \Phi_{|S_k|}) : X \dashrightarrow \text{Im } \Phi_{|S_k|} =: Y$$

associated to S_k for any $k \in N(X, K_X + L, h_L)$ sufficiently large. Then $f^b : X^b \rightarrow Y^b$ is an analytic fiber space, $\kappa(X, K_X + L, h_L) = \dim Y^b$ and $\kappa(F, K_F + L_F, h_L|_F) = 0$, where F is the very general (which means, $F = X_y$ when y varies outside a countable union of analytic subsets of Y) fiber of f^b .

Remark 1.2. Theorem 4.15 generalizes a classical result on Itaka fibration (cf. (i) of Theorem 1.11 in [26]) with multiplier ideal sheaves involved.

The key to the proof of Theorem 1.5 is using the fact that $\kappa(X, K_X + L, h_L) = \tilde{\kappa}(X, K_X + L, h_L)$, where $\tilde{\kappa}(X, K_X + L, h_L)$ is defined by (1.13). As an immediate application of Theorem 1.5, we are able to obtain a distinct proof of Theorem 1.4 in the case when h_L has analytic singularities.

Corollary 1.6 (a special case of Theorem 1.4). *Let $f : X \rightarrow Y$ be an analytic fiber space between a compact Kähler manifold X and a projective manifold Y . Let $L \rightarrow X$ be a \mathbb{Q} -line bundle equipped with a singular metric h_L with semi-positive*

curvature current. Assume moreover that h_L has analytic singularities and Y has general type, then

$$(1.15) \quad \kappa(X, K_X + L, h_L) = \kappa(F, K_F + L|_F, h_L|_F) + \dim Y$$

holds, where F denotes the sufficiently general fiber of f .

Remark 1.3. In the case when (X, Δ) is a compact Kähler klt pair with a \mathbb{Q} -effective divisor Δ , Y is of general type and set $(L, h_L) = (\mathcal{O}_X(\Delta), h_\Delta)$, it follows immediately from Corollary 1.6 that

$$(1.16) \quad \kappa(X, K_X + L) = \kappa(F, K_F + L_F) + \dim Y.$$

Such addition formula (1.16) has also been proved in [4] and [27] when (X, Δ) is a projective log canonical pair (see also Theorem 1.7 in [14]).

The proof of Corollary 1.6 will be different from that of Theorem 1.4. It is based on a quantitative analysis of the classical Kodaira meromorphic map (cf. Theorem 1.5) and some recent developments of analytic tools, which mainly consists of L^2 extension theory and the positivity of relative k -Bergman metrics.

This paper is organized as follows. In section 2, we give some preliminaries for the proofs of main theorems. The most important results will be the extension results: Lemma 2.4 and Lemma 2.5. In section 3, we will complete the proofs of Theorem 1.2 and Theorem 1.3. We also provide an analytic approach to Theorem 1.1. In section 4, we first recall some basic results about Iitaka D -dimension and Okounkov body associated to graded algebra of almost integral type. Then we discuss their applications to generalized Kodaira dimensions. Consequently, we give the proof of Theorem 1.5 in the end. In section 5, we give another proof of Corollary 1.6 in the special case when h_L has analytic singularities, which will be different from that of [40].

2. PRELIMINARIES

2.1. Relative k -Bergman metrics for Kähler fibrations. Let $f : X \rightarrow Y$ be a surjective holomorphic map between an n -dimensional compact Kähler manifold X and an m -dimensional compact connected complex manifold Y . Let L be a pseudoeffective line bundle over X equipped with a (semi-)positively curved singular hermitian metric $h_L = e^{-\varphi_L}$, where $\varphi_L \in L^1_{\text{loc}}$ is a plurisubharmonic local weight function. For $k \in \mathbb{N}$, the stalk of the k -multiplier ideal sheaf $\mathcal{I}_k(h_L) \subset \mathcal{O}_X$ at $x \in X$ is defined as follows:

$$(2.1) \quad \mathcal{I}_k(h_L)_x := \{F \in \mathcal{O}_{X,x}; |F|^{2/k} e^{-\varphi_L} \in L^1_{\text{loc}} \text{ near } x\}.$$

When $k = 1$, It is well-known from Nadel's theorem that $\mathcal{I}(h_L) = \mathcal{I}_1(h_L)$ is a coherent analytic sheaf. For general k , the coherence of the k -multiplier ideal sheaf follows from the strong openness property of the multiplier ideal sheaves ([18]) and Demailly's equisingular approximation theorem ([9]) for plurisubharmonic functions (cf. [31], [5]).

Let $K_{X/Y} := K_X - f^*K_Y$ be the relative canonical bundle. We set

$$Y_0 := \{y \in Y \mid \text{rank } df(x) = m, \forall x \in f^{-1}(y)\}$$

to be the regular values of f , $Y_h := \{y \in Y_0 : h_L|_{X_y} \not\equiv +\infty\}$ and

$$(2.2) \quad Y_{k,\text{ext}} := \{y \in Y_0 \mid \text{rank } f_*(kK_{X/Y} + kL) = h^0(X_y, kK_{X_y} + kL|_{X_y})\}.$$

and respectively $X_{k,\text{ext}} := f^{-1}(Y_{k,\text{ext}})$. It is clear that both Y_0 and $Y_{k,\text{ext}}$ are Zariski open dense subsets of Y . Assume that

$$H^0(X_y, (kK_{X_y} + kL|_{X_y}) \otimes \mathcal{I}_k(h_L)) \neq 0$$

holds for some $y \in Y_{k,\text{ext}} \cap Y_h$. Now we can assign the k -Bergman metric, denoted by $B_{k,y}^{-1}$, where

$$B_{k,y}(x) := \sup\{u(x) \otimes \bar{u}(x); u \in H^0(X_y, kK_{X_y} + kL|_{X_y}) \text{ and } \int_{X_y} |u|^{2/k} h_L \leq 1\},$$

to each fiber X_y for $y \in Y_{k,\text{ext}}$. Gluing them together we may endow $(kK_{X/Y} + kL)|_{X_{k,\text{ext}}}$ with a metric denoted by $B_{k,X/Y}^{-1}$, which is known as the relative k -Bergman metric. It has been proved that

Theorem 2.1. (Theorem 1.5 in [39]) $B_{k,X/Y}^{-1}$ has semi-positive curvature current on $X_{k,\text{ext}}$. Furthermore, it can be extended across $X \setminus X_{k,\text{ext}}$ as a new metric with semi-positive curvature current.

Remark 2.1. When f is projective, this result was initially obtained in [30] (see also [1]). When $k = 1$ and f is a proper fibration between a Kähler manifold X and a complex manifold Y (X not necessarily compact), this result was obtained in [37] by using the so-called Guan-Zhou method in [19]. For general $k \geq 2$, a crucial step in [39] to establish Theorem 2.1 is overcoming the extension difficulties which were pointed out in Remark 4.2.4 in [30].

Now consider the set $\tilde{Y}_{k,h_L,\text{ext}} := \{y \in Y_0; h^0(X_y, (kK_{X_y} + kL|_{X_y}) \otimes \mathcal{I}_k(h_L)|_{X_y}) = \text{rank } f_*(kK_{X/Y} + kL) \otimes \mathcal{I}_k(h_L)\}$ and respectively

$$Y_{k,h_L,\text{ext}} := \{y \in Y_{k,\text{ext}} \cap \tilde{Y}_{k,h_L,\text{ext}}; \mathcal{I}_k(h_L)|_{X_y} = \mathcal{I}_k(h_L|_{X_y})\}.$$

In [10], the authors has obtained another new proof of the positivity of the relative k -Bergman metric on $f^{-1}(\tilde{Y}_{k,h_L,\text{ext}})$. According to Fubini's theorem and the basic extension property of k -multiplier ideal sheaves (cf. Theorem 2.4 in [40] or Proposition 0.2 in [2]), one may find that $\mathcal{I}_k(h_L)|_{X_y} = \mathcal{I}_k(h_L|_{X_y})$ for almost all $y \in Y$ and $Y \setminus Y_{k,h_L,\text{ext}}$ is of measure zero. As a consequence, for any $y \in \bigcap_{k=1}^{\infty} Y_{k,h_L,\text{ext}}$, $h^0(X_y, (kK_{X_y} + kL|_{X_y}) \otimes \mathcal{I}_k(h_L|_{X_y}))$ is constant for any $k \in \mathbb{N}$, which explains why $\kappa(F, K_F + L|_F, h_L|_F)$ does not vary for almost all F in Theorem 1.4.

The following proposition will be useful for clarifying the relationship between Nadel's multiplier ideal sheaves and k -multiplier ideal sheaves:

Proposition 2.2. Let L be a line bundle, $k \in \mathbb{N}$ and $B_{k,X/Y}^{-1}$ be the natural extension of the relative Bergman metric on $k(K_{X/Y} + L)$ according to Theorem 2.1, then

- a) $\Gamma(X_y, k(K_{X_y} + L|_{X_y}) \otimes \mathcal{I}_k(h_L|_{X_y})) \subset \Gamma(X_y, k(K_{X_y} + L|_{X_y}) \otimes \mathcal{I}((B_{k,X/Y}^{-\frac{k-1}{k}} \cdot h_L)|_{X_y}))$ for general X_y ;
- b) Let G be a line bundle on Y . Then $\Gamma(X, k(K_{X/Y} + L) \otimes f^*G \otimes \mathcal{I}(B_{k,X/Y}^{-\frac{k-1}{k}} \cdot h_L)) \subset \Gamma(X, k(K_{X/Y} + L) \otimes f^*G \otimes \mathcal{I}_k(h_L))$.

Proof. a) Note $B_{k,X/Y}^{-1}|_{X_y}$ for $y \in Y_{k,\text{ext}}$ is nothing but the k -Bergman kernel metric on X_y , therefore, for any $s \in \Gamma(X_y, k(K_{X_y} + L|_{X_y}) \otimes \mathcal{I}_k(h_L|_{X_y}))$ we get

$$(2.3) \quad \int_{X_y} |s|^2 B_{k,X/Y}^{-\frac{k-1}{k}} \cdot h_L \leq \int_{X_y} |s|^2 \cdot \left(\frac{\|s\|_k}{|s|}\right)^{2\frac{k-1}{k}} \cdot h_L = \|s\|_k^2,$$

where

$$(2.4) \quad \|s\|_k := \left(\int_{X_y} |s|^{2/k} h_L \right)^{k/2}.$$

Our claim follows by (2.3).

- b) We first denote that $\tilde{h} := B_{k,X/Y}^{-\frac{k-1}{k}} \cdot h_L$. For any $\tilde{s} \in \Gamma(X, k(K_{X/Y} + L) \otimes f^*G \otimes \mathcal{I}_k(h_L))$, and any smooth metric h_{K_Y} (resp. h_G) on K_Y (resp. G), we have

$$\begin{aligned} \int_X |\tilde{s}|^{\frac{2}{k}} h_L \cdot f^*(h_G^{-1} h_{K_Y}^{-1}) &= \int_X \frac{|\tilde{s}|^{\frac{2}{k}} f^*(h_G h_{K_Y}^{-1})^{\frac{1}{k}}}{B_{k,X/Y}^{\frac{k-1}{k^2}}} \cdot \frac{B_{k,X/Y}^{\frac{k-1}{k^2}}}{(f^* h_{K_Y})^{\frac{k-1}{k}}} \cdot h_L \\ &\stackrel{\text{(H\"older inequality)}}{\leq} \left(\int_X \frac{|\tilde{s}|^2 h_L f^*(h_G h_{K_Y}^{-1})}{B_{k,X/Y}^{\frac{k-1}{k}}} \right)^{\frac{1}{k}} \left(\int_X \frac{B_{k,X/Y}^{\frac{1}{k}} h_L}{f^*(h_{K_Y})} \right)^{\frac{k-1}{k}} \\ &= \left(\int_X |\tilde{s}|^2 \tilde{h} f^*(h_G h_{K_Y}^{-1}) \right)^{\frac{1}{k}} \left(\int_X \frac{B_{k,X/Y}^{\frac{1}{k}} h_L}{f^*(h_{K_Y})} \right)^{\frac{k-1}{k}} \\ (2.5) \quad &\stackrel{\text{Fubini}}{=} \left(\int_X |\tilde{s}|^2 \tilde{h} f^*(h_G h_{K_Y}^{-1}) \right)^{\frac{1}{k}} \left(\int_{y \in Y} \frac{1}{h_{K_Y}} \int_{X_y} B_{k,y}^{\frac{1}{k}} h_L \right)^{\frac{k-1}{k}}. \end{aligned}$$

To analyze the last integral, one just needs to use the fact that

$$(2.6) \quad \int_{X_y} B_{k,y}^{\frac{1}{k}} h_L \leq \text{rank } f_*((kK_{X/Y} + L) \otimes \mathcal{I}_k(h_L)) < \infty$$

holds for $y \in Y_{k,\text{ext}}$ (cf. Lemma 2.7 in [40]). Finally we verify our claim from inequality (2.5). \square

Remark 2.2. When L is a pseudoeffective \mathbb{Q} -line bundle which can be equipped with a (semi-)positively curved singular metric h_L , Proposition 2.2 is still true for any $k \in \mathbb{N}$ sufficiently divisible (so that kL is a line bundle).

2.2. L^2 Extension theory on weakly pseudoconvex manifolds. We will adopt the following Ohsawa-Takegoshi L^2 extension theorem on weakly pseudoconvex K\"ahler manifolds:

Theorem 2.3. (Theorem 2.8 and Remark 2.9 (b) in [7], or Theorem 2.1 in [40]) *Let (X, ω) be a weakly pseudoconvex K\"ahler manifold and ψ be a quasi-psh function on X with neat analytic singularities, i.e. ψ can be locally written as:*

$$\psi = c \log(|F_1|^2 + \cdots + |F_N|^2) + u,$$

where each F_j is holomorphic, $c \in \mathbb{R}_{\geq 0}$ and u is smooth. Let S be the zero variety $V(\mathcal{I}(\psi))$. Assume that ψ has log canonical singularities, i.e.

$$\mathcal{I}((1 - \varepsilon)\psi)|_S = \mathcal{O}_X|_S$$

holds for every $\varepsilon > 0$. Denote by S^0 the regular set of S . Let P be a holomorphic line bundle over X equipped with a singular hermitian metric h_P . Assume that there is an $\alpha > 0$ such that

$$(2.7) \quad \sqrt{-1}\Theta_{h_P} + \xi \sqrt{-1}\partial\bar{\partial}\psi \geq 0$$

holds for all $\xi \in [1, 1 + \alpha]$. Then for every

$$f \in \Gamma(S^0, (K_X + P)|_{S^0})$$

and

$$\int_{S^0} |f|_{h_\omega \otimes h_P}^2 dV_{X,\omega}[\psi] < \infty,$$

there exists an $F \in \Gamma(X, K_X + P)$ such that $F|_S = f$ and

$$(2.8) \quad \int_X \frac{\rho(\alpha\psi)}{e^\psi} |F|_{h_\omega \otimes h_P}^2 dV_{X,\omega} \leq \frac{34}{\alpha} \int_{S^0} |f|_{h_\omega \otimes h_P}^2 dV_{X,\omega}[\psi],$$

where $\rho : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $\rho(t) = e^{-\frac{t}{2}}$ when $t > 0$, $\rho(t) = \frac{1}{1+t^2}$ when $t \leq 0$.

Remark 2.3. Here the Ohsawa measure $dV_{X,\omega}[\psi]$ on S^0 is defined by

$$(2.9) \quad \int_{S^0} g dV_{X,\omega}[\psi] := \limsup_{t \rightarrow -\infty} \int_{\{x: t < \psi < t+1\}} \tilde{g} e^{-\psi} dV_{X,\omega},$$

where \tilde{g} is any smooth extension of a non-negative continuous function $g \in \mathcal{C}(S^0)$ with compact support. Originally, the measure $dV_{X,\omega}[\psi]$ was claimed to be well-defined on S^0 by Proposition 4.5 in [7].

However, as the latest paper [23] suggests, this claim was made under the implicit “unique lc place” assumption, which says that each irreducible component of the non-klt locus S is dominated by a unique divisor with discrepancy equals to -1 . It can be checked easily that when ψ is locally of the form $\psi = k \cdot \log(|z_1|^2 + \dots + |z_k|^2) + \mathcal{C}^\infty$ (which is also the setting for later use), where $(z_1 = \dots = z_k = 0)$ are local defining equations for S (thus S is non-singular and k is the codimension of S), the so-called “unique lc place” assumption is automatically satisfied and the Ohsawa measure $dV_{X,\omega}[\psi]$ has smooth positive density with respect to the measure $dV_{S,\omega|_S}$ (cf. (2.6) in [7]). In fact, this special log canonical singularity assumption on ψ is the setting where the Ohsawa measure first appeared (cf. [28]).

One can also refer to [38] for an optimal version of the above theorem.

Lemma 2.4. *Let $f : X \rightarrow Y$ be a surjective holomorphic mapping between a compact Kähler manifold X and a compact complex manifold Y . Let $L \rightarrow X$ be a pseudoeffective \mathbb{Q} -line bundle equipped with a singular metric h_L whose curvature current is semi-positive. Let k_0 be the least integer so that $k_0 L$ is a line bundle and let G be an ample line bundle over Y . Let $\psi \leq 0$ be a quasi-psh function on Y , which is smooth outside a very general point $y \in \bigcap_{k=1}^\infty Y_{kk_0, \text{ext}}$ and assume that*

$$\psi(z) - \log |z - y|^{2m}$$

is smooth near y (such ψ always exists! cf. (3.12)). Moreover, assume that there exists a hermitian metric h_{G-K_Y} on $G - K_Y$ such that:

$$(2.10) \quad \sqrt{-1} \Theta_{h_{G-K_Y}}(G - K_Y) + \xi \sqrt{-1} \partial \bar{\partial} \psi \geq 0$$

for all $\xi \in [1, 1 + \alpha]$ and $\alpha > 0$. Then for any integers k and

$$s \in \Gamma(X_y, kk_0(K_{X_y} + L|_{X_y}) \otimes \mathcal{I}_{kk_0}(h_L|_{X_y}))$$

and $y \in \bigcap_{k=1}^\infty Y_{kk_0, \text{ext}}$, there exists a section

$$\tilde{s} \in \Gamma(X, (kk_0(K_{X/Y} + L) + f^*G) \otimes \mathcal{I}_{kk_0}(h_L))$$

such that $\tilde{s}|_{X_y} = s$.

Proof. This is proved by a similar argument of Theorem 1.2 in [40]. Let

$$(2.11) \quad P := (kk_0 - 1)(K_{X/Y} + L) + L + f^*(G - K_Y)$$

be a line bundle endowed with a singular hermitian metric:

$$(2.12) \quad h_P := B_{kk_0, X/Y}^{-(kk_0-1)/kk_0} \cdot f^* h_{G-K_Y} \cdot h_L.$$

According to (2.10), (2.12) and Theorem 2.1, we may compute its curvature by:

$$(2.13) \quad \begin{aligned} \sqrt{-1}\Theta_{h_P} + \xi\sqrt{-1}\partial\bar{\partial}f^*\psi &= (kk_0 - 1)\sqrt{-1}\Theta_{B_{kk_0, X/Y}^{-1/kk_0}}(K_{X/Y} + L) + \sqrt{-1}\Theta_{h_L}(L) + \\ &f^*(\xi\sqrt{-1}\partial\bar{\partial}\psi + \sqrt{-1}\Theta_{h_{G-K_Y}}(G - K_Y)) \geq 0. \end{aligned}$$

Note $V(\mathcal{I}(f^*\psi)) = X_y$ and $f^*\psi$ has log canonical singularities (which also satisfies the “unique lc place” assumption), applying Theorem 2.3 and Remark 2.3 we see that

$$\begin{aligned} \Gamma(X, (K_X + P) \otimes \mathcal{I}(h_P)) &= \Gamma(X, (kk_0(K_{X/Y} + L) + f^*G) \otimes \mathcal{I}(B_{kk_0, X/Y}^{-\frac{kk_0-1}{kk_0}} \cdot h_L)) \rightarrow \\ \Gamma(X_y, (K_X + P)|_{X_y} \otimes \mathcal{I}(h_P|_{X_y})) &= \Gamma(X_y, kk_0(K_{X_y} + L|_{X_y}) \otimes \mathcal{I}(B_{kk_0, X/Y}^{-\frac{kk_0-1}{kk_0}}|_{X_y} \cdot h_L|_{X_y})) \end{aligned}$$

is surjective for every $y \in Y_0$. Combining relation *a*) and *b*) in Proposition 2.2 (see also Remark 2.2) we immediately get the desired extension section \tilde{s} when $y \in \bigcap_{k=1}^{\infty} Y_{kk_0, \text{ext}}$. \square

Remark 2.4. One may refer to Theorem 2.11 in [11] for a qualitative formulation of this result under the assumption that f is projective. Our improvement consists of two aspects. One is that it gives a more precise version of a quantitative estimate of the restriction map, the other is that we only assume $f : X \rightarrow Y$ to be a Kähler fiber space with the help of Theorem 2.1. Compared with Theorem 1.2 in [40], we do not assume the positivity of the canonical bundle K_Y to control the negative part of the closed $(1, 1)$ -current $f^*\sqrt{-1}\partial\bar{\partial}\psi$.

We can also formulate a slight modified version of Lemma 2.4. The main difference is that the inequality does not involve any multiplier ideal sheaves.

Lemma 2.5. *Let $f : X \rightarrow Y$ be an algebraic fiber space between two projective manifolds X and Y . Let $k \in \mathbb{N}$. Let $L = \mathcal{O}_X(D)$ be a line bundle over X , where D is a simple normal crossing divisor on X . Let G be an ample line bundle over Y . Let H be an ample line bundle over X . Let y be a very general point in Y and $\psi \leq 0$ be a quasi-psh function on Y which is smooth outside y . Assume that*

$$\psi(z) - \log|z - y|^{2m}$$

is smooth near y . Moreover, assume that there exists a hermitian metric h_{G-K_Y} on $G - K_Y$ such that:

$$(2.14) \quad \sqrt{-1}\Theta_{h_{G-K_Y}}(G - K_Y) + \xi\sqrt{-1}\partial\bar{\partial}\psi \geq 0$$

for all $\xi \in [1, 1 + \alpha]$ and $\alpha > 0$. Then for any $k \in \mathbb{N}$ and

$$s \in \Gamma(X_y, k(K_{X_y} + L|_{X_y}) + H|_{X_y}),$$

there exists a section

$$\tilde{s} \in \Gamma(X, (k(K_{X/Y} + L) + H + f^*(G)))$$

such that $\tilde{s}|_{X_y} = s$.

Proof. Since D is a simple normal crossing divisor, for each $k \in \mathbb{N}$, the big \mathbb{Q} -line bundle $L_k := L + \frac{1}{k}H$ can be equipped with a singular metric h_k with analytic singularities, whose curvature current is semi-positive and multiplier ideal sheaf $\mathcal{I}(h_k)$ is trivial. Thus $\mathcal{I}(h_k|_{X_y})$ is also trivial for general $y \in Y$. Hence $\|s\|_k^{2/k} = \int_{X_y} |s|^{2/k} h_k < \infty$. Let $B_{k,X/Y,L_k}^{-1}$ denote the (extended) relative k -Bergman kernel metric on $k(K_{X/Y} + L_k)$, with respect to the algebraic fiber space $f : X \rightarrow Y$ and the metric h_k on L_k . Fix the integer $k \in \mathbb{N}$. Then for general $y \in Y$ and $s \in \Gamma(X_y, k(K_{X_y} + L|_{X_y}) + H|_{X_y}) = \Gamma(X_y, k(K_{X_y} + L_k|_{X_y}))$, the following inequality

$$(2.15) \quad (B_{k,X/Y,L_k}^{-1})|_{X_y} = B_{k,y,h_k}^{-1} \leq \left| \frac{s}{\|s\|_k} \right|^{-\frac{2}{k}}$$

holds, where B_{k,y,h_k} stands for the k -Bergman kernel of $k(K_{X_y} + L_k|_{X_y})$ with respect to the metric $h_k|_{X_y}$. Let

$$(2.16) \quad P := (k-1)(K_{X/Y} + L_k) + L_k + f^*(G - K_Y)$$

be a line bundle over X . Now we define a metric on P as

$$(2.17) \quad h_P := B_{k,X/Y,L_k}^{-(k-1)/k} \cdot h_k \cdot f^* h_{G-K_Y}$$

whose curvature current is semi-positive, according to (2.14) and Theorem 2.1. Then for general $y \in Y$,

$$(2.18) \quad C_y \cdot \int_{X_y} |s|^2 h_P = \int_{X_y} |s|^2 B_{k,X/Y,L_k}^{-(k-1)/k} \cdot h_k \leq \|s\|_k^{\frac{2(k-1)}{k}} \cdot \int_{X_y} |s|^{\frac{2}{k}} \cdot h_k = \|s\|_k^2 < \infty$$

holds thanks to (2.15), where $C_y > 0$ is a constant depending only on y . Therefore, applying Theorem 2.3 and Remark 2.3 again we get a section

$$\tilde{s} \in \Gamma(X, K_X + P) = \Gamma(X, (k(K_{X/Y} + L) + H + f^*(G)))$$

such that $\tilde{s}|_{X_y} = s$. As k varies among all integers, the desired extension property holds for very general $y \in Y$. \square

3. PROOF OF THEOREM 1.2 AND THEOREM 1.3

Before giving the proof, let us give some simple but useful lemmas.

Lemma 3.1. *Let X be a compact complex manifold and (L, h_L) be a pseudoeffective line bundle over X . Let $a \in \mathbb{N}$ and $\kappa_{\sigma,a,A}(X, K_X + L, h_L)$ be the following number*

$$(3.1) \quad \max\{\nu \in \mathbb{Z}; \limsup_{k \rightarrow \infty} \frac{h^0(ka(K_X + L) \otimes A \otimes \mathcal{I}_{ak}(h_L))}{k^\nu} > 0\}$$

and $\kappa_{\sigma,a} := \max_{l \in \mathbb{N}} \{\kappa_{\sigma,a,lA}\}$. Then $\kappa_{\sigma,a} = \kappa_\sigma$.

Proof. It suffices to prove $\kappa_{\sigma,a} \geq \kappa_\sigma$. By the definition of κ_σ there exists l_0 and a sequence $\{k_\nu\}_{\nu=1}^\infty$ such that $N_\nu := h^0(k_\nu(K_X + L) \otimes l_0 A \otimes \mathcal{I}_{k_\nu}(h_L)) \geq C \cdot k_\nu^{\kappa_\sigma}$, then we obtain

$$(3.2) \quad h^0(ak_\nu(K_X + L) \otimes al_0 A \otimes \mathcal{I}_{ak_\nu}(h_L)) \geq N_\nu \geq C \cdot k_\nu^{\kappa_\sigma}$$

for any $\nu \in \mathbb{N}$. Therefore, $\kappa_{\sigma,a} \geq \kappa_{\sigma,a,al_0 A} \geq \kappa_\sigma$. \square

Lemma 3.2. *Let $f : X \rightarrow Y$ be an algebraic fiber space between two projective manifolds X and Y . Let D_Y be an effective divisor on Y and \mathcal{E} be a coherent sheaf*

on X . Assume $\mathcal{F} = f_*\mathcal{E}$ is a sufficiently generically globally generated coherent sheaf on Y . Then

$$(3.3) \quad h^0(X, \mathcal{E} \otimes f^*\mathcal{O}_Y(D_Y)) \geq h^0(Y, \mathcal{O}_Y(D_Y)) \cdot \text{rk } \mathcal{F},$$

where $\text{rk } \mathcal{F}$ is the rank of \mathcal{F} .

Proof. Let us assume $q = h^0(Y, \mathcal{O}_Y(D_Y)) \geq 1, r = \text{rk } \mathcal{F} \geq 1$ without loss of generality. According to projection formula, $H^0(X, \mathcal{E} \otimes f^*\mathcal{O}_Y(D_Y)) = H^0(Y, \mathcal{O}_Y(D_Y) \otimes \mathcal{F})$. It follows from (generalized) Grauert's direct image and semi-continuity theorem (cf. Theorem 10.6 and Theorem 10.7 in [3]) that $\mathcal{F}|_{Y_0}$ is locally free for some analytic Zariski open dense subset Y_0 contained in the regular values of f and $\mathcal{F}_y \otimes k(y) = H^0(X_y, \mathcal{E}|_{X_y})$, $\text{rk } \mathcal{F} = h^0(X_y, \mathcal{E}|_{X_y})$ for any $y \in Y_0$. Since $\mathcal{F} = f_*\mathcal{E}$ is sufficiently generically globally generated, one may pick $y_0 \in Y_0$ and r linearly independent sections $\sigma_1, \dots, \sigma_r \in H^0(Y, \mathcal{F})$ such that $\sigma_{1,y_0}, \dots, \sigma_{r,y_0}$ generates the stalk \mathcal{F}_{y_0} . Note \mathcal{F} is coherent, hence there is a small open neighborhood $\Omega \subset Y_0$ of y_0 , such that $\sigma_{1,y}, \dots, \sigma_{r,y}$ generates the stalk \mathcal{F}_y for any $y \in \Omega$. In particular, $\sigma_1|_{X_y}, \dots, \sigma_r|_{X_y}$ form a basis of $H^0(X_y, \mathcal{E}|_{X_y})$ for $y \in \Omega$.

Let ξ_1, \dots, ξ_q be a basis of $H^0(Y, \mathcal{O}_Y(D_Y))$. We claim that $\{\xi_i \sigma_j\}_{i=1, \dots, q, j=1, \dots, r}$ are linearly independent in $H^0(Y, \mathcal{O}_Y(D_Y) \otimes \mathcal{F}) = H^0(X, \mathcal{E} \otimes f^*\mathcal{O}_Y(D_Y))$ and $h^0(X, \mathcal{E} \otimes f^*\mathcal{O}_Y(D_Y)) \geq qr$.

Actually, consider the following equations for $t_{ij} \in \mathbb{C}$:

$$(3.4) \quad \sum_{i=1, \dots, q, j=1, \dots, r} t_{ij} \xi_i \sigma_j = 0,$$

We will show $t_{ij} = 0$. Take q general points $y_1, \dots, y_q \in \Omega$ which will be determined later. Assume Ω also trivializes $\mathcal{O}_Y(D_Y)$ with a holomorphic frame e and $\xi_i|_{\Omega} = \tilde{\xi}_i(y) \cdot e$ with $\tilde{\xi}_i \in \mathcal{O}(\Omega)$. When restricting (3.4) to X_{y_1}, \dots, X_{y_q} , we find q equations:

$$(3.5) \quad \sum_{j=1}^r \left(\sum_{i=1}^q t_{ij} \tilde{\xi}_i(y_k) \cdot f^*e \right) \sigma_j|_{X_{y_k}} = 0, k = 1, \dots, q$$

Since $\sigma_1|_{X_{y_1}}, \dots, \sigma_r|_{X_{y_k}}$ are linearly independent,

$$(3.6) \quad \sum_{i=1}^q t_{ij} \tilde{\xi}_i(y_k) = 0, j = 1, \dots, r, k = 1, \dots, q.$$

Fix any j now and consider the $q \times q$ matrix $[\tilde{\xi}_i(y_k)]$, if $[\tilde{\xi}_i(y_k)]$ is non-degenerate, then $t_{ij} = 0$ for all $i = 1, \dots, q$. To this end, we will pick $y_k \in \Omega$ as follows.

We will argue step by step. First choose y_1 such that $\tilde{\xi}_1(y_1) \neq 0$, this is possible since $\xi_1 \neq 0$. Consider the analytic subset $A_2 := \{y \in \Omega; \xi_1(y_1)\tilde{\xi}_2(y) - \tilde{\xi}_2(y_1)\tilde{\xi}_1(y) = 0\}$ of Ω . It can be shown that $A_2 \neq \Omega$ since otherwise $\xi_1(y_1)\xi_2 - \xi_2(y_1)\xi_1 = 0$, which contradicts the fact that ξ_1, ξ_2 are linearly independent. Therefore we pick $y_2 \in \Omega \setminus A_2$. Iterating this process $q-1$ times, we finally get an analytic subset

$$(3.7) \quad A_q := \{y \in \Omega; \sum_{i=1}^q (-1)^{q-i} \alpha_i \tilde{\xi}_i(y) = 0\}$$

with $\alpha_i = \det(\tilde{\xi}_s(y_t))_{s=1, \dots, \hat{i}, \dots, q, t=1, \dots, q-1}$ and $\alpha_q \neq 0$. Therefore, A_q is proper since ξ_1, \dots, ξ_q are linearly independent. One may pick $y_q \in \Omega \setminus A_q$ and $[\tilde{\xi}_i(y_k)]_{i,k=1, \dots, q}$ is non-degenerate.

In summary, $t_{ij} = 0$ and $h^0(X, \mathcal{E} \otimes f^*\mathcal{O}_Y(D_Y)) \geq qr$. \square

Now we are in position to give the proof of Theorem 1.3.

Proof. (Proof of Theorem 1.3) Let us first verify that for any semi-positive line bundle $H \rightarrow X$ and any $k \in \mathbb{N}$, some divisible $k_0 \in \mathbb{N}$ (such that $k_0 L$ is a line bundle) and some ample enough line bundle $G \rightarrow Y$,

$$(3.8) \quad \mathcal{F} := \mathcal{O}_Y(G) \otimes f_* \mathcal{O}_X((kk_0 K_{X/Y} + kk_0 L) \otimes H \otimes \mathcal{I}_{kk_0}(h_L))$$

is sufficiently generically globally generated. H is to be determined at the end of the proof.

Actually, It suffices to show that the restriction morphisms

$$(3.9) \quad \Gamma(X, (kk_0 K_{X/Y} + kk_0 L + f^* G + H) \otimes \mathcal{I}_{kk_0}(h_L)) \rightarrow \Gamma(X_y, kk_0 (K_{X_y} + L|_{X_y}) \otimes \mathcal{O}(H|_{X_y}) \otimes \mathcal{I}_{kk_0}(h_L|_{X_y}))$$

are surjective for sufficiently general y . The reason why (3.9) implies (3.8) is that

$$(3.10) \quad \mathcal{F}_y = \Gamma(X_y, kk_0 (K_{X_y} + L|_{X_y}) \otimes \mathcal{O}(H|_{X_y}) \otimes \mathcal{I}_{kk_0}(h_L|_{X_y}) \otimes \mathcal{O}_{Y,y})$$

for y belonging to a Zariski dense open subset of Y thanks to the Grauert's direct image and semi-continuity theorem (cf. Theorem 10.6 and Theorem 10.7 in [3]), and

$$(3.11) \quad \mathcal{I}_{kk_0}(h_L)|_{X_y} = \mathcal{I}_{kk_0}(h_L|_{X_y})$$

holds for sufficiently general $y \in Y$.

The function ψ can be constructed as in [40] as follows. Take a smooth concave function $\theta : \mathbb{R} \rightarrow (-\infty, 0]$ such that $\theta(t) = t$ when $t \in (-\infty, -1]$, $\theta(t) = 0$ when $t \in [1, \infty)$ and $0 \leq \theta' \leq 1$, $\theta'' \geq -C_1$ for all $t \in \mathbb{R}$. Let $z = (z_1, \dots, z_m)$ be the coordinate functions on a coordinate chart U centered at y . Let ψ be a global quasi-plurisubharmonic function on Y defined by

$$(3.12) \quad \psi = \theta(\log |z - y|^{2m}).$$

In fact, $\psi \leq 0$ and $i\partial\bar{\partial}\psi$ is only supported on $U' := U \cap \{z; -e \leq |z - y| \leq e\}$ and

$$(3.13) \quad i\partial\bar{\partial}\psi|_{U'} \geq \theta''(\log |z - y|^{2m}) \cdot (i\partial\bar{\partial} \log |z - y|^{2m})|_{U'} \geq -C_2 \omega|_{U'}$$

for some Kähler metric ω and $C_2 > 0$. By compactness, for any $y \in Y$, we find a uniform constant C_3 independent of y , so that there exists a quasi-plurisubharmonic function ψ such that $\psi \sim 2m \log |z - y|$ near y and $i\partial\bar{\partial}\psi \geq -C_3 \omega$.

Now applying Lemma 2.4 (replace L in Lemma 2.4 by $L + (kk_0)^{-1}H$ equipped with the metric $h_H^{1/kk_0} \cdot h_L$, where h_H is semi-positive) we know that (3.9) holds true for some G ample enough, which finishes our claim of (3.8).

Setting h^0 as the complex dimension of the cohomology group H^0 and using projection formula, one may observe that

$$(3.14) \quad \begin{aligned} h^0(X, kk_0(K_X + L) \otimes H \otimes f^* G \otimes \mathcal{I}_{kk_0}(h_L)) &\geq \text{rk } \mathcal{F} \cdot h^0(Y, kk_0 K_Y) \\ &= h^0(F, kk_0(K_F + L_F) \otimes H|_F \otimes \mathcal{I}_{kk_0}(h_L|_F)) \cdot h^0(Y, kk_0 K_Y) \end{aligned}$$

holds for sufficiently general fiber F thanks to Lemma 3.2.

In order to prove (1.9), let us take a positive integer k_1 and a positive integer C_4 such that

$$(3.15) \quad h^0(Y, kk_1 K_Y) \geq C_4 k^{\kappa(Y, K_Y)}$$

holds for every integers $k \gg 0$. Then we obtain

$$h^0(X, kk_2(K_X + L) \otimes H \otimes f^* G \otimes \mathcal{I}_{kk_2}(h_L)) \geq$$

$$(3.16) \quad C_5 k^{\kappa(Y, K_Y)} \cdot h^0(F, kk_2(K_F + L_F) \otimes H|_F \otimes \mathcal{I}_{kk_2}(h_L|_F))$$

holds for $k_2 = k_1 k_0$ and every positive enough integer k according to (3.14) and (3.15), where $C_5 = C_4 k_0^{\kappa(Y, K_Y)}$. If we choose H to be sufficiently ample then

$$(3.17) \quad h^0(F, k_\nu k_2(K_F + L_F) \otimes H|_F \otimes \mathcal{I}_{k_\nu k_2}(h_L|_F)) \geq C' k_\nu^{\kappa_\sigma(F, K_F + L_F, h_L|_F)} = C' k_\nu^{\kappa_\sigma(F, K_F + L_F, h_L|_F)}$$

holds true with $C' > 0$, by Lemma 3.1 and passing to some subsequence $\{k_\nu\}_{\nu=1}^\infty$. Finally, inequality (1.9) follows immediately by the fact that for any $\nu \gg 0$

$$(3.18) \quad h^0(X, k_\nu k_2(K_X + L) \otimes H \otimes f^*G \otimes \mathcal{I}_{k_\nu k_2}(h_L)) \geq C_5 C' k_\nu^{\kappa_\sigma(F, K_F + L_F, h_L|_F) + \kappa(Y, K_Y)},$$

which is due to (3.16) and (3.17).

It remains to prove (1.10). For this purpose, by setting $D_Y = kk_0 K_Y + (N-1)G$ and $H = \mathcal{O}_X$ in Lemma 3.2, it follows that

$$(3.19) \quad h^0(X, kk_0(K_X + L) \otimes f^*(NG) \otimes \mathcal{I}_{kk_0}(h_L)) \geq \text{rk } \mathcal{F} \cdot h^0(Y, kk_0 K_Y + (N-1)G) = h^0(F, kk_0(K_F + L_F) \otimes \mathcal{I}_{kk_0}(h_L|_F)) \cdot h^0(Y, kk_0 K_Y + (N-1)G)$$

holds for sufficiently general fiber F , where N is any positive integer (actually, we see from its proof that (3.14) still holds if we replace G by NG for any positive integers N). Let us take another positive integer \tilde{k}_1 so that

$$(3.20) \quad h^0(F, k\tilde{k}_1(K_F + L_F) \otimes \mathcal{I}_{k\tilde{k}_1}(h_L|_F)) \geq \tilde{C}_4 k^{\kappa(F, K_F + L_F, h_L|_F)}$$

holds for every integers $k \gg 0$ from the definition of $\kappa(F, K_F + L_F, h_L|_F)$. Therefore, thanks to (3.19) and (3.20), we obtain that

$$(3.21) \quad h^0(X, k\tilde{k}_2(K_X + L) \otimes f^*(NG) \otimes \mathcal{I}_{k\tilde{k}_2}(h_L)) \geq \tilde{C}_5 k^{\kappa(F, K_F + L_F, h_L|_F)} \cdot h^0(Y, k\tilde{k}_2 K_Y + (N-1)G)$$

holds for $\tilde{k}_2 = \tilde{k}_1 k_0$ and every $k \gg 0$, where $\tilde{C}_5 = \tilde{C}_4 k_0^{\kappa(F, K_F + L_F, h_L|_F)}$. Now choose N sufficiently large so that after extracting a subsequence $\{k_\nu\}_{\nu=1}^\infty$,

$$(3.22) \quad h^0(Y, k_\nu \tilde{k}_2 K_Y + (N-1)G) \geq \tilde{C}' k_\nu^{\kappa_\sigma(Y, K_Y)}$$

holds for every ν with $\tilde{C}' > 0$. In conclusion, we finally get that

$$(3.23) \quad h^0(X, k_\nu \tilde{k}_2(K_X + L) \otimes f^*G \otimes \mathcal{I}_{k_\nu \tilde{k}_2}(h_L)) \geq \tilde{C}_5 \tilde{C}' k_\nu^{\kappa_\sigma(F, K_F + L_F, h_L|_F) + \kappa_\sigma(Y, K_Y)}$$

for any $\nu \gg 0$ by (3.21) and (3.22), which confirms (1.10). \square

Remark 3.1. From the proof of (1.10), we may observe that the left hand side in (1.10) can be replaced by a more optimal number $\kappa_{\sigma, f, \text{hor}}$ defined as in (1.11).

Remark 3.2 (Analytic proof of Theorem 1.1). By the same token as above, we can also give an analytic method towards Theorem 1.1. For this purpose, we first claim that for any $k \in \mathbb{N}$

$$(3.24) \quad \mathcal{O}_Y(G) \otimes f_* \mathcal{O}_X(kK_{X/Y} + k(D_X - f^*D_Y) + H)$$

is generically globally generated for some ample line bundle G over Y and any ample line bundle H over X .

Indeed, the projection formula yields that

$$\Gamma(Y, \mathcal{O}_Y(G) \otimes f_* \mathcal{O}_X(kK_{X/Y} + k(D_X - f^*D_Y) + H)) =$$

$$(3.25) \quad \Gamma(X, k(K_{X/Y} + D_X - f^*D_Y) + H + f^*G)$$

holds. Since $D_X \supset f^*D_Y$, $D_X - f^*D_Y$ has simple normal crossings. Hence we can also construct ψ as in (3.12) and choose G and H as in Lemma 2.5. Then the general stalk of the locally free sheaf (3.24) (whose restriction to the complement of some $\text{codim} \geq 2$ analytic subset in Y is locally free) becomes

$$(3.26) \quad \Gamma(X_y, (k(K_{X_y} + D_X|_{X_y}) + H|_{X_y}) \otimes \mathcal{O}_{Y,y})$$

and our claim is an immediate consequence of Lemma 2.5 by setting $L = \mathcal{O}_X(D_X - f^*D_Y)$.

The rest of the proof follows by Lemma 3.2 and the last half (i.e., computing and estimating h^0) of the proof of Theorem 1.3. The only difference is that in order to prove (1.6), though the use of ample H in (3.24) is unavoidable, we can still apply inequality

$$(3.27) \quad h^0(F, k\tilde{k}_1(K_F + D_F) + H_F) \geq h^0(F, k\tilde{k}_1(K_F + D_F)) \geq \tilde{C}k^{\kappa(F, K_F + D_F)}$$

to substitute (3.20).

Remark 3.3 (about the proof of Theorem 1.2). Among the first half (i.e. the reduction to the surjective statement and the construction of ψ) of the proof of Theorem 1.3 and Remark 3.2 as above, we have seen that Theorem 1.2 can be obtained via Lemma 2.4 and Lemma 2.5 immediately.

4. IITAKA D -DIMENSIONS AND THEIR GENERALIZATIONS

The goal of this part is to recall three equivalent definitions of Iitaka D -dimensions and generalize these results to generalized Kodaira dimensions. The main references for this section are [26] and [32].

4.1. Iitaka D -dimension. Let us first recall some facts in commutative algebra.

Let $R = \bigoplus_{k \in \mathbb{N}} R_k$ be a graded \mathbb{C} -algebra and an integral domain (or briefly, \mathbb{C} -domain). Suppose that $R_k = 0$ for all $k < 0$ and $R_0 = \mathbb{C}$. Let $Q(R)$ be the quotient field of R . Let $R^{(m)} := \bigoplus_{k \in \mathbb{N}} R_{km}$ be the graded \mathbb{C} -subdomain for some $m \in \mathbb{N}$. Let $R^\#$ denote the multiplicative subset of all nonzero homogeneous elements.

The quotient ring of R , denoted by $R^{\#-1}R$, is also a graded \mathbb{C} -domain. Its degree 0 part $(R^{\#-1}R)_0$ is also a field which will be denoted by $Q((R))$. It will be easy to check $Q((R^{(m)})) = Q((R))$ for any $m \in \mathbb{N}$.

Let $N(R) := \{k > 0; R_k \neq 0\}$ and let $M_{\geq n} := \{k \in M; k \geq n\}$ for any subset $M \subset \mathbb{N}$ and $n \in \mathbb{N}$. Since $N(R)$ is a semigroup, we know that $N(R)_{\geq n} = (d\mathbb{N})_{\geq n}$ for $n \gg 0$ and some $d = \gcd N(R) \in \mathbb{N}$.

Let $S = \bigoplus_{k \in \mathbb{N}} S_k$ be a graded \mathbb{C} -subalgebra of R . For any $k \geq 0$, let $S_0[S_k]$ denote the graded \mathbb{C} -subdomain generated by S_0 and S_k .

The following propositions are a list of basic properties in commutative algebra.

Proposition 4.1. $R^{(m)} \subset R$ is an integral extension of rings for any integers $m \in \mathbb{N}$. In particular, the Krull dimensions of all $R^{(m)}$ are all the same.

Proposition 4.2. If R is finitely generated over $R_0 = \mathbb{C}$, then the Krull dimension of R equals the transcendence degree of $Q(R)$ over \mathbb{C} .

Proposition 4.3. If R is finitely generated over R_0 , then for all sufficiently divisible $d \in \mathbb{N}$, the graded subalgebra $R^{(d)}$ is finitely generated in degree 1 over R_0 (i.e., $R^{(d)} = R_0[R_1^{(d)}]$).

Proposition 4.4. *If R is finitely generated in degree 1 over R_0 , then the Hilbert function $\text{HF} : k \in \mathbb{N} \mapsto \dim_{\mathbb{C}} R_k \in \mathbb{N}$ has growth order $\delta - 1$, where δ is the Krull dimension of R . Precisely, this means $\text{HF}(k) \sim C \cdot k^{\delta-1}$ as $k \rightarrow \infty$ for some $C > 0$.*

Let X be a smooth projective variety, $Q(X)$ be the rational function field of X and D be a Cartier divisor on X . Let $L = \mathcal{O}_X(D)$ be the corresponding invertible sheaf of D .

From now on, we always set $R(X, L) := R = \bigoplus_{k \in \mathbb{N}} R_k$ and

$$R_k := H^0(X, \mathcal{O}_X(kD)) = H^0(X, kL)$$

be the associated \mathbb{C} -graded algebra. We also set $N(X, L) := N(R)$, $Q((X, L)) := Q((R))$. Note $Q((R))$ is algebraically closed in $Q(X)$ (cf. Proposition 1.4 in [26]).

For any $k \in N(X, L)$, R_k induces a Kodaira meromorphic mapping (cf. examples below Definition 4.6)

$$(4.1) \quad \Phi_{|kL|} : X \dashrightarrow \mathbb{P}(H^0(X, kL)),$$

which is a morphism outside the base locus. When $N(X, L) = \emptyset$, set $\kappa(X, L)$ to be $-\infty$. When $N(X, L) \neq \emptyset$, the Iitaka D -dimension $\kappa(X, L)$ or $\kappa(X, D)$ is defined as either one of the following numbers.

Proposition 4.5. *The following numbers are equal when $N(X, L) \neq \emptyset$:*

1. $\kappa^{(1)}(X, L) := \text{tr. deg}_{\mathbb{C}} R(X, L) - 1 = \text{tr. deg}_{\mathbb{C}} Q((X, L))$;
2. $\kappa^{(2)}(X, L) := \max_{k \in N(X, L)} \dim \Phi_{|kL|}(X)$;
3. $\kappa^{(3)}(X, L) := \max\{\nu \in \mathbb{N}; \limsup_{k \rightarrow \infty} \frac{h^0(X, kL)}{k^\nu} > 0\}$.

Let us briefly explain the idea to the proof of Proposition 4.5. It follows easily from the fact that

$$(4.2) \quad Q((X, L)) = Q(\text{Im } \Phi_{|kL|})$$

for all $k \in N(X, K_X + L)$ sufficiently large (cf. Proposition 1.4 in [26]) that $\kappa^{(1)}(X, L) = \kappa^{(2)}(X, L)$.

To see the reason why $\kappa^{(2)}(X, L) = \kappa^{(3)}(X, L)$, there are, to the best of the authors' knowledge, at least two ways to show this result. On one hand, the first strategy relies on the existence of Iitaka fibration (cf. Theorem 1.12 and Corollary 1.13 in [26]), which aims at reducing the problems on the line bundle L to a big line bundle on a smooth model of $\text{Im } \Phi_{|kL|}$. On the other hand, by fixing a faithful \mathbb{Z}^n -valued valuation $v : Q(X) \setminus \{0\} \rightarrow \mathbb{Z}^n$ on X a priori, the second strategy (cf. Theorem 3.3 + Corollary 1.16 in [22]) considers a general graded algebra $R \subset Q(X)$ (in this case we set $R = R(X, L)$) of almost integral type and applies some basic facts from convex geometry (cf. Theorem 1.14 in [22]) to obtaining that both $\kappa^{(2)}(X, L)$ and $\kappa^{(3)}(X, L)$ coincide with the dimension of its Newton convex body of $\Delta_v(X, L)$ related to $R = R(X, L)$. We will see in the next subsection that how these two methods can be generalized, when considering the graded subalgebra containing multiplier ideal sheaf and assuming X is merely compact (i.e. possibly non-algebraic).

In the special case when $R(X, L)$ is a finitely generated algebra, $\kappa^{(2)}(X, L) = \kappa^{(3)}(X, L)$ can be proved by directly applying Proposition 4.1-4.4. In fact, if $R(X, L)$ is finitely generated, then by choosing d sufficiently divisible, we obtain $\kappa^{(3)}(X, L) = \kappa^{(3)}(X, dL) = \dim R^{(d)} - 1 = \text{tr. deg}_{\mathbb{C}} Q((R^{(d)})) = \kappa^{(2)}(X, dL) = \kappa^{(2)}(X, L)$.

4.2. Generalized Kodaira dimension. In this part, we will still adopt the notation as in subsection 4.1. Given a vector space V , the notation $\mathbb{P}V$ stands for all 1-dimensional quotients of V or all hyperplanes of V^* .

Let X be a compact (connected) complex manifold. Let L be a pseudoeffective line bundle over X equipped with a singular hermitian metric h_L whose curvature current is semi-positive. The following results in this subsection can all be generalized when L is a \mathbb{Q} -bundle.

Let $\mathcal{I}_k(h_L)$ be the k -multiplier ideal sheaf as in (2.1), then

$$(4.3) \quad S_k := H^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L))$$

contains all holomorphic sections s of $\mathcal{O}_X(k(K_X + L))$ on X and

$$(4.4) \quad \|s\|_k := \left(\int_X |s|^{2/k} h_L \right)^{k/2} < \infty.$$

Note although at each k -level $\|\cdot\|_k$ does not satisfy the triangle inequality (hence not a norm), it still can be checked that

$$(4.5) \quad \|s_1 + s_2\|_k \leq 2^k (\|s_1\|_k + \|s_2\|_k)$$

holds for any $s_1, s_2 \in S_k$.

From now on, let us take $R_k = H^0(X, k(K_X + L))$ and $R = \bigoplus_{k \in \mathbb{N}} R_k$. Set

$$(4.6) \quad S = \bigoplus_{k \in \mathbb{N}} S_k \subset R,$$

where S_k is defined as in (4.3). By the basic fact that $\mathcal{I}_k(h_L)$ is a coherent sheaf, it is recognized that S_k is a linear subspace of R_k . Since for $s_m \in S_m$ and $s_l \in S_l$

$$(4.7) \quad \|s_m \cdot s_l\|_{m+l} \leq \|s_m\|_m \cdot \|s_l\|_l$$

holds by Hölder inequality, S becomes a graded \mathbb{C} -subalgebra of R .

Let $Q((X, K_X + L, h_L))$ and $N(X, K_X + L, h_L)$ denote the degree zero part of the quotient ring of S and respectively the corresponding semigroup of S (see section 4.1). Let $\mathbb{P}H^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L))$ be the projective space of all hyperplanes of $H^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L))$.

Definition 4.6. (Definition 2.2 in [32]) *Let X, Y be two complex spaces. A mapping from X to the power set of Y , denoted by $\Phi : X \dashrightarrow Y$, is called a **meromorphic mapping**, if the followings are satisfied:*

1. *The graph $G := \{(x, y) \in X \times Y; y \in \varphi(x)\}$ is an irreducible analytic subset in $X \times Y$;*
2. *The projection map $p_X : G \rightarrow X$ is a proper modification.*

Take $k \in N(X, K_X + L, h_L)$, then basic examples of meromorphic mappings

$$(4.8) \quad \Phi_{|S_k|} : X \dashrightarrow \mathbb{P}H^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L)) = \mathbb{P}S_k$$

will be constructed as follows (cf. Example 2.4.1 and 2.4.2 in [32] when $\mathcal{I}_k(h_L)$ is trivial).

Let $\{s_0, \dots, s_{N_k}\}$ be a basis of S_k and set

$$(4.9) \quad \Sigma := \{x \in X; s_0(x) = \dots = s_{N_k}(x) = 0\}$$

to be a nowhere dense analytic subset of X . Then there is a holomorphic mapping

$$(4.10) \quad \Phi : X \setminus \Sigma \rightarrow \mathbb{P}^{N_k}, x \mapsto \{s \in S_k; s(x) = 0\}$$

and G be the closure of the graph of Φ in $X \times \mathbb{P}^{N_k}$. One can conclude that G is an (irreducible) analytic subset and $p_X : G \rightarrow X$ is a proper modification. Hence

G induces a meromorphic map denoted by (4.8). Let $\text{Im } \Phi_{|S_k|}$ be the image of the projection of G onto \mathbb{P}^{N_k} . By Remmert's proper mapping theorem, $\text{Im } \Phi_{|S_k|}$ is a projective algebraic variety in \mathbb{P}^{N_k} .

Proposition 4.7. $Q(\text{Im } \Phi_{|S_k|}) = Q((X, K_X + L, h_L))$ for all $k \in N(X, K_X + L, h_L)$ sufficiently large enough.

Proof. By the very basic fact that the (meromorphic) function field $Q(X)$ is finitely generated (cf. Theorem 3.1 in [32]), any subfield of $Q(X)$ will be finitely generated. As a consequence, $Q(X, K_X + L, h_L)$ is finitely generated over \mathbb{C} . Then there exists $M > 0$ so that $Q((X, K_X + L, h_L)) = Q((S_0[S_k])) = Q(\text{Im } \Phi_{|S_k|})$ for all $k \in N(X, K_X + L, h_L)_{\geq M}$ (cf. (1.2 i) in [26]). \square

Proposition 4.8. $Q((X, K_X + L, h_L))$ is algebraically closed in $Q(X)$.

Proof. Let us first show S is integrally closed in R . Take any $r \in R_\nu$ ($\nu \in \mathbb{N}$) satisfying the following equation

$$(4.11) \quad r^k + s_1 r^{k-1} + \cdots + s_{k-1} r + s_k = 0$$

where $s_j \in S_{\nu j}$ for $j = 1, \dots, k$, we claim that $r \in S_\nu$ or equivalently

$$(4.12) \quad \|r\|_\nu \leq 2^{\nu k^2} (k+1) \max_{j=1, \dots, k} \{\|s_j\|_{\nu j}^{1/j}\} < \infty.$$

Indeed, if by contradiction (4.12) does not hold, then

$$(4.13) \quad \|r\|_\nu^j > 2^{\nu k^2} (k+1) \|s_j\|_{\nu j}$$

holds for any $j \in [1, k]$. According to (4.5) and equation (4.11), we get

$$\begin{aligned} \|r\|_\nu^k &= \|r^k\|_{\nu k} \leq 2^{\nu k} (\|s_1 r^{k-1}\|_{\nu k} + \|r^k + s_1 r^{k-1}\|_{\nu k}) \\ &\leq 2^{\nu k} (\|s_1 r^{k-1}\|_{\nu k} + 2^{\nu k} (\|r^k + s_1 r^{k-1} + s_2 r^{k-2}\|_{\nu k} + \|s_2 r^{k-2}\|_{\nu k})) \leq \dots \\ (4.14) \quad &\leq 2^{\nu k^2} (\|s_1 r^{k-1}\|_{\nu k} + \cdots + \|s_k\|_{\nu k}). \end{aligned}$$

It follows from (4.7), (4.4) and (4.13) that

$$(4.15) \quad \|s_j r^{k-j}\|_{\nu k} \leq \|s_j\|_{\nu j} \cdot \|r^{k-j}\|_{\nu(k-j)} = \|s_j\|_{\nu j} \cdot \|r\|_\nu^{k-j} \leq \frac{1}{2^{\nu k^2} (k+1)} \|r\|_\nu^k$$

for $j \in [1, k]$. Thanks to (4.14) and (4.15), we eventually get the contradiction:

$$(4.16) \quad \|r\|_\nu^k \leq 2^{\nu k^2} \cdot k \cdot \frac{1}{2^{\nu k^2} (k+1)} \|r\|_\nu^k = \frac{k}{k+1} \|r\|_\nu^k$$

as $r \neq 0$. Hence (4.12) has been proved.

The rest of the proof will be almost the same as Proposition 1.4 in [26]. Let ξ be the $Q(X)$ -scheme $\text{Spec } Q(X)$ and $\xi \rightarrow X$ be the inclusion (see Ex2.7 in GTM52 for the details of this map). Let $(K_X + L)_\xi$ be the pull back of $K_X + L$. Since S is integrally closed in R , R is integrally closed in $R(\xi, (K_X + L)_\xi)$, we know that S is integrally closed in $R(\xi, (K_X + L)_\xi)$. Therefore, $Q((S))$ is algebraically closed in $S^{\#-1} R(\xi, (K_X + L)_\xi)$ by (1.2) in [26]. Note $Q(X) = R(\xi, (K_X + L)_\xi)_0 \subset S^{\#-1} R(\xi, (K_X + L)_\xi)$, we can now obtain that $Q((S))$ is algebraically closed in $Q(X)$. \square

Consider the following three numbers when $N(X, K_X + L, h_L) \neq \emptyset$:

$$(4.17) \quad \kappa^{(1)}(X, K_X + L, h_L) := \text{tr} \cdot \deg_{\mathbb{C}} Q((X, K_X + L, h_L))$$

$$(4.18) \quad \kappa^{(2)}(X, K_X + L, h_L) := \max_{k \in N(X, K_X + L, h_L)} \dim \text{Im } \Phi_{|S_k|}$$

$$(4.19) \quad \kappa^{(3)}(X, K_X + L, h_L) := \max\{m \in \mathbb{N}; \limsup_{k \rightarrow \infty} \frac{h^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L))}{k^m} > 0\}.$$

Note $\kappa^{(3)}(X, K_X + L, h_L)$ has already been introduced in [40].

The following proposition illustrates how these numbers vary under blow-ups.

Proposition 4.9. *Let $\mu : X' \rightarrow X$ be a blow-up of X with smooth center Z of $\text{codim} \geq 2$, then inequality*

$$(4.20) \quad \kappa^{(j)}(X, K_X + L, h_L) = \kappa^{(j)}(X', K_{X'} + L', h_{L'})$$

*holds for $j = 1, 2, 3$, where $(L', h_{L'}) = (\mu^*L, \mu^*h_L)$.*

Proof. By definition of $\kappa^{(j)}(X, K_X + L, h_L)$, since $H^0(X', k(K_{X'} + L') \otimes \mathcal{I}_k(h_{L'})) = H^0(X, \mu_*(kK_{X'} \otimes kL' \otimes \mathcal{I}_k(h_{L'})))$, it will be sufficient for us to show

$$(4.21) \quad \mu_*(\mathcal{O}_{X'}(kK_{X'} + kL') \otimes \mathcal{I}_k(h_{L'})) = \mathcal{O}_X(kK_X + kL) \otimes \mathcal{I}_k(h_L)$$

for any $k \in \mathbb{N}$. Let us write $h_L = e^{-2\varphi_L}$ on some open coordinate $U \subset X$ which trivializes L , then $h_{L'} = e^{-2\mu^*\varphi_L}$ on $\mu^{-1}(U)$.

By the definition of multiplier ideal sheaves, given any open subset $U \subset X$, the coherent analytic sheaf $\mathcal{O}_X(kK_X + kL) \otimes \mathcal{I}_k(h_L)(U)$ consists of all holomorphic k -canonical (i.e. $K_U^{\otimes k}$ -valued) forms f on U such that

$$(4.22) \quad \int_V c_n(f \wedge \bar{f})^{1/k} e^{-2\varphi_L} < \infty$$

holds for any $V \Subset U$. The change of variable formula yields that

$$(4.23) \quad \int_{\mu^{-1}(V)} c_n(\mu^*f \wedge \overline{\mu^*f})^{1/k} e^{-2\mu^*\varphi_L} = \int_V c_n(f \wedge \bar{f})^{1/k} e^{-2\varphi_L} < \infty.$$

Hence μ^*f is the pull-back holomorphic k -canonical form on $\mu^{-1}(U)$ such that (4.23) holds. Therefore, $\mu_*(kK_{X'} \otimes kL' \otimes \mathcal{I}_k(h_{L'})) \supset kK_X \otimes kL \otimes \mathcal{I}_k(h_L)$.

On the other side, if we a priori know that a holomorphic k -canonical form μ^*f on $\mu^{-1}(U)$ satisfies (4.23), then f must be a holomorphic k -canonical form on U by the fact that μ is an isomorphism outside $\mu^{-1}(Z)$. Actually, this can be done according to the $\text{codim} \geq 2$ extension theorem for holomorphic functions. Moreover, it must also satisfy (4.22). As a result, $\mu_*(kK_{X'} \otimes kL' \otimes \mathcal{I}_k(h_{L'})) \subset kK_X \otimes kL \otimes \mathcal{I}_k(h_L)$ holds. This finishes the proof of (4.21). \square

Now let us turn to compare $\kappa^{(1)}$, $\kappa^{(2)}$ and $\kappa^{(3)}$. Our next goal is to show these three numbers all coincide with the dimension of an appropriate Newton-Okounkov body associated to X , L and h_L , which also equals the so-called generalized Kodaira dimension. To this end, let us first recall some basic constructions of the Newton-Okounkov body of any given graded algebra $S \subset Q(X)$, where $Q(X)$ is the rational function field of a projective manifold X . A systematic study of the Newton-Okounkov body can be referred to [25] and [22], which was based on the pioneering work of A. Okounkov (cf. [29]). The following notations and basic facts are mainly taken from [22].

Let $P \subset \mathbb{Z}^n$ be a semigroup. Let G be the subgroup of \mathbb{Z}^n generated by P , L be the subspace of \mathbb{R}^n spanned by P . Let C be the smallest closed convex cone with apex at the origin generated by P . The regularization \tilde{P} of P is defined to be the semigroup $G \cap C$ contained in L . Now assume that the cone C is strongly convex (i.e. the linear subspace contained in C is only zero) and set $\dim L = q + 1$. Fix a rational half-space $M \subset L$ (i.e. ∂M can be spanned by rational vectors) containing P . Take a linear map $\pi_M : L \rightarrow \mathbb{R}$ so that $\ker \pi_M = \partial M$, $\pi_M(L \cap \mathbb{Z}^n) = \mathbb{Z}$ and $\pi_M(M \cap \mathbb{Z}^n) = \mathbb{Z}_{\geq 0}$.

Let $H_P(k) := \text{Card}(P \cap \pi_M^{-1}(k))$ and $H_{\tilde{P}}(k) := \text{Card}(\tilde{P} \cap \pi_M^{-1}(k))$ be the Hilbert function of P and \tilde{P} respectively, where $\text{Card}(\cdot)$ stands for the cardinality of a set. Then the convex body (i.e. a compact convex set, which can be confirmed due to the fact that C is strongly convex) $\Delta(P) := \tilde{P} \cap \pi_M^{-1}(1)$ of dimension q is called the *Newton-Okounkov body* of the semigroup P . Let us fix the notations $m(P)$, $\text{ind}(P)$ standing for the index of the subgroup $\pi_M(G)$ in \mathbb{Z} , and the index of the subgroup $G \cap \partial M$ in $\mathbb{Z}^{n-1} \times \{0\}$ respectively.

Let X be a projective manifold and $Q(X)$ be the rational function field. Let $S = \bigoplus_{k \in \mathbb{N}} S_k \subset Q(X)$ be a graded algebra of *almost integral type* (see the precise definition in section 2.3 in [22]). The basic example for such S includes all graded subalgebras of $R(X, L)$ associated to an arbitrary line bundle L over X (cf. Theorem 3.7 in [22]). Fix a faithful \mathbb{Z}^n -valued valuation $v : Q(X)^* \rightarrow \mathbb{Z}^n$ (i.e. $v(Q(X)^*) = \mathbb{Z}^n$) with respect to the total ordering of \mathbb{Z}^n (e.g. for $p = (p_1, \dots, p_n)$, $q = (q_1, \dots, q_n) \in \mathbb{Z}^n$, say $p > q$ iff for some $1 \leq r < n$ we have $p_i = q_i$ for $i = 1, \dots, r$ and $p_{r+1} > q_{r+1}$). Then v naturally induces a valuation $v_t : Q(X)[t]^* \rightarrow \mathbb{Z}^{n+1}$ extending v (see the details of the construction in section 2.4 in [22]). The valuation v_t maps non-zero elements of $S[t] := \bigoplus_{k \in \mathbb{N}} S_k t^k$ to a semigroup, denoted by $P(S)$, of integral points contained in $\mathbb{Z}^n \times \mathbb{Z}_{\geq 0}$. Fix the rational half-space $M = \mathbb{R}^n \times \mathbb{R}_{\geq 0}$ and take $\pi_M : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ to be the projection map to the last coordinate. It can be shown that the cone associated to $P(S)$ is strongly convex since S is of almost integral type (cf. Theorem 2.30 in [22]). Then one can define the convex body $\Delta(P(S)) = \tilde{P}(S) \cap (\mathbb{Z}^n \times \{1\})$ to be the *Newton-Okounkov body*, denoted by $\Delta(S)$, of the algebra S . Let us fix the notations $m(S)$, $\text{ind}(S)$ standing for the indices $m(P(S))$, $\text{ind}(P(S))$ for the semigroup $P(S)$ respectively.

We will adopt the following two important properties of Newton-Okounkov bodies in our context.

Lemma 4.10 (Theorem 2.31 in [22]). *Let $S = \bigoplus_{k \in \mathbb{N}} S_k$ be an algebra of almost integral type with the Newton-Okounkov body $\Delta(S)$. Put $m = m(S)$, $q = \dim \Delta(S)$ and the Hilbert function $H_S(k) := \dim S_k$. Then the q -th growth coefficient of the function $H_S(m \cdot)$*

$$a_q := \lim_{k \rightarrow \infty} \frac{H_S(mk)}{k^q}$$

exists and $a_q = \text{Vol}_q(\Delta(S))/\text{ind}(S) > 0$.

Lemma 4.11 (a part of Theorem 3.3 in [22]). *Let S be an algebra of almost integral type in $Q(X)$ and S_k be the k -th subspace of the algebra S . Let $Y_k := \text{Im } \Phi_k$, where Φ_k is the Kodaira map defined in (4.8) associated to the linear subspace S_k . If p is sufficiently large and divisible by $m(S)$, then $\dim Y_p$ is independent of p and equals the dimension of the Newton-Okounkov body $\Delta(S)$.*

Let X now be a (possibly non-algebraic) compact complex manifold. The algebraic dimension $a(X)$ of X is defined to be the transcendental degree of $Q(X)$, namely the field of meromorphic functions, over \mathbb{C} .

Definition 4.12 (Algebraic reduction, cf. p. 25 in [32]). *A surjective morphism $a : X' \rightarrow A$ is called an algebraic reduction if it satisfies the following conditions:*

- 1) X' is also a complex manifold and bimeromorphically equivalent to X .
- 2) A is a projective manifold and $\dim A = a(X)$.
- 3) a induces an isomorphism between $Q(X)$ and $Q(A)$.

An algebraic reduction always exists (cf. p. 24-25 in [32]), and is unique up to a bimeromorphic equivalence. In addition, by the very basic fact Corollary 1.10 in [32], the fibers of $a : X' \rightarrow A$ are connected (i.e. $a : X' \rightarrow A$ is an analytic fiber space) from the above condition 3).

Let F be a holomorphic line bundle over X . The following result will be a useful tool for the study of the asymptotic behavior of a graded linear system associated to X and F . For completeness, we will also give a sketch of proof of the following lemma, whose original idea comes from [36].

Lemma 4.13 (Theorem 1 in [36]). *Let $a_0 \in \mathbb{N}$ such that $a_0 F$ is effective. Then there exists a smooth projective variety A (independent of a_0) and an algebraic reduction of X such that there exists a \mathbb{Q} -effective divisor D over A and*

$$(4.24) \quad H^0(A, kD) = H^0(X, ka_0 F)$$

holds for $k > 0$ sufficiently divisible.

Proof. With the existence of the algebraic reduction of X (cf. Definition 4.12) and the neat model of any holomorphic fibrations between compact complex manifolds (cf. Lemma 1.3 in [4] or Lemma 2 in [36]), we may assume there exists a proper modification $\mu : X' \rightarrow X$ and an algebraic reduction $a : X' \rightarrow A$, so that every a -exceptional divisor is also μ -exceptional. We decompose the divisor $\mu^*(a_0 F)$ by (the dimension of the image of each irreducible component of $\mu^* F$ under the algebraic reduction a)

$$(4.25) \quad \mu^*(a_0 F) = N' + \sum_{j=1}^J b_j D'_j + R',$$

where N' is an effective divisor so that $a(N'_0) = A$ for each irreducible component N'_0 of N' (thus N' is non-polar, in the sense of Definition 1.2 in [4]), each D'_j is a prime divisor whose image under a is of codimension one (hence also a prime divisor in A , according to section 9.1.3 in [17]) with coefficients $b_j \in \mathbb{Z}_{>0}$ and R' stands for an a -exceptional divisor.

Let us denote $D_j := a(D'_j)$ for $j = 1, \dots, J$ and assume all D_l ($l = 1, \dots, L$) are not equal to each other with $L \leq J$, while D_{j_1} ($j_1 > L$) equals one of the D_{j_2} ($j_2 = 1, \dots, L$) (since it might happen that $D_p = D_q$ for $p \neq q \in \{1, \dots, J\}$). Then we consider the following decomposition

$$(4.26) \quad a^* D_l = \sum_{j=1}^J c_{j,l} D'_j + \sum_{m=1}^{M_l} d_{m,l} E'_l + R'_l, l = 1, \dots, L$$

with prime divisors E'_l for $l = 1, \dots, L$ which do not contain any D'_j for $j = 1, \dots, J$, a -exceptional divisors R'_l for $l = 1, \dots, L$ and all integers $c_{j,l} \geq 0, d_{m,l} > 0$ and

$c_{j,l} > 0$ for at least one $j \in \{1, \dots, J\}$. For $j \in \{1, \dots, J\}$, we set $f_{j,l} = b_j$ if $c_{j,l} > 0$ and $f_{j,l} = 0$ if $c_{j,l} = 0$. Then

$$(4.27) \quad G'_l := \sum_{j=1}^J f_{j,l} D'_j > 0$$

will be the maximal divisor whose support is contained in all D'_j ($j = 1, \dots, J$) and $a^* D_l$, so that $\mu^*(a_0 F) - G'_l$ is still effective.

Consider the \mathbb{Q} -effective divisor

$$D := \sum_{l=1}^L g_l D_l$$

with $g_l = \min_{j=1, \dots, J} \frac{b_j}{c_{j,l}} \in \mathbb{Q}$ if $M_l = 0$; $g_l = 0$ if $M_l \neq 0$ ($l = 1, \dots, L$). If $g_l = 0$, then it can be shown that G'_l will be partially supported on the fibers of a (see its definition in Definition 1.21 in [4]). In this case, we set $T'_l := 0$. If $g_l > 0$, then all $f_{j,l}$ are not zero. Set $T'_l := g_l R'_l$ then $G'_l - g_l \mu^*(D_l) + T'_l$ will also be a \mathbb{Q} -effective divisor partially supported on the fibers of a , thanks to (4.26) and (4.27). Denote by $T' := \sum_{l=1}^L T'_l$ the a -exceptional (hence also μ -exceptional) \mathbb{Q} -effective divisor. Summing up with the index $l = 1, \dots, L$, thus we obtain $P' := \mu^*(a_0 F) - N' - a^* D + T'$ will be a \mathbb{Q} -effective divisor partially supported on the fibers of a according to (4.25).

Choosing $k \in \mathbb{N}$ sufficiently divisible, we will see that

$$H^0(X, ka_0 F) = H^0(X', k\mu^*(a_0 F) + kT') = H^0(X', kP' + kN' + ka^* D) = H^0(A, kD)$$

holds, by the basic property of exceptional divisors (cf. Lemma 3 in [36]), non-polar divisors (cf. Lemma 4 in [36]) and partially supported divisors (cf. Lemma 1 in [36] or Lemma 1.22 in [4]). \square

We are now in position to show that the aforementioned three numbers defined from (4.17) to (4.19) all coincide, which will all be denoted by $\kappa(X, K_X + L, h_L)$ from now on.

Proposition 4.14. $\kappa^{(1)}(X, K_X + L, h_L) = \kappa^{(2)}(X, K_X + L, h_L) = \kappa^{(3)}(X, K_X + L, h_L)$.

Proof. The first equality follows easily from the fact that

$$(4.28) \quad Q((X, K_X + L, h_L)) = Q(\text{Im } \Phi_{|S_k|})$$

for all $k \in N(X, K_X + L, h_L)$ sufficiently large.

Let us then show that

$$(4.29) \quad \kappa^{(2)} = \kappa_a^{(2)}, \kappa^{(3)} = \kappa_a^{(3)}$$

for any integers $a \in \mathbb{N}$, where

$$(4.30) \quad \kappa_a^{(2)} := \max_{k \in \mathbb{N}} \dim \text{Im } \Phi_{|S_{ak}|}$$

and

$$(4.31) \quad \kappa_a^{(3)} := \max\{m \in \mathbb{N}; \limsup_{k \rightarrow \infty} \frac{h^0(X, ak(K_X + L) \otimes \mathcal{I}_{ak}(h_L))}{k^m} > 0\}.$$

For $\kappa^{(2)}$, one just need to use the fact $Q((S^{(a)})) = Q((S))$ and the first equality $\kappa^{(2)}(X, K_X + L, h_L) = \kappa^{(1)}(X, K_X + L, h_L) = \text{tr} \cdot \deg_{\mathbb{C}} Q((S))$. For $\kappa^{(3)}$, by definition, it is clear that $\kappa^{(3)} \geq \kappa_a^{(3)}$. On the other side, if there exists $\{k_\nu\}_{\nu=1}^\infty$ such that

$$(4.32) \quad N_\nu := h^0(X, k_\nu(K_X + L) \otimes \mathcal{I}_{k_\nu}(h_L)) \geq C k_\nu^{\kappa^{(3)}}$$

for some $C > 0$, then $h^0(X, a k_\nu(K_X + L) \otimes \mathcal{I}_{a k_\nu}(h_L)) \geq N_\nu \geq C k_\nu^{\kappa^{(3)}}$ and $\kappa^{(3)} \leq \kappa_a^{(3)}$. Hence $\kappa^{(3)} = \kappa_a^{(3)}$.

Applying Lemma 4.13 with $F = K_X + L$, there exist a smooth projective variety A and a \mathbb{Q} -effective divisor D on A so that $H^0(A, kD) = H^0(X, k a_0(K_X + L)) = H^0(X', k a_0(K_{X'} + L'))$ holds for some integer $a_0 > 0$, any $k \in \mathbb{N}$ sufficiently divisible. Furthermore, the following diagram

$$(4.33) \quad \begin{array}{ccc} X' & \xrightarrow{\mu} & X \\ \downarrow & & \\ A & & \end{array}$$

holds, where $\mu : X' \rightarrow X$ is a proper modification and $X' \rightarrow A$ is a surjective holomorphic map with connected fibers. We may assume $H^0(A, k a D) = H^0(X, k a a_0(K_X + L))$ holds for some other integer a and any integers k .

Therefore, $S^{(a a_0)} \subset W := \bigoplus_{k \in \mathbb{N}} H^0(A, k a D)$ will be an algebra of almost integral type according to Theorem 3.7 in [22]; thus we can consider $\Delta(S^{(a a_0)})$, which is defined to be the Newton-Okounkov body of $S^{(a a_0)}$. We highlight that $\Delta(S^{(a a_0)})$ might depend on the choice of the algebraic reduction and A as pointed out in Remark 1 in [36] (however, its dimension does not rely on the choice of A).

Identifying $S'_{k a a_0} := H^0(X', k a a_0(K_{X'} + L') \otimes \mathcal{I}_{k a a_0}(h_{L'}))$ as a linear subspace V_k of $H^0(A, k a D)$ via an isomorphism θ_k for any $k \in \mathbb{N}$ (note $S'_{k a a_0} = S_k^{(a a_0)}$ by Proposition 4.9), then $V_k V_l \subset V_{k+l}$ and we have the following diagram

$$(4.34) \quad \begin{array}{ccc} X' & \xrightarrow{\Phi_{|S'_{k a a_0}|}} & \text{Im} \Phi_{|S'_{k a a_0}|} \subset \mathbb{P}^N \\ \downarrow & & \theta_k^* \downarrow \\ A & \xrightarrow{\Phi_{|V_k|}} & \text{Im} \Phi_{|V_k|} \subset \mathbb{P}^N \end{array}$$

with θ_k^* an isomorphism between $\text{Im} \Phi_{|S'_{k a a_0}|}$ and $\text{Im} \Phi_{|V_k|}$ (indeed, $\theta_k^* \in \text{Aut}(\mathbb{P}^N)$) induced by θ_k . Hence $\kappa^{(2)}(X, K_X + L, h_L) = \max_{k \in \mathbb{N}} \dim \text{Im} \Phi_{|V_k|}$. Now applying Lemma 4.11 and Lemma 4.10 respectively to the graded algebra $S^{(a a_0)} \simeq \bigoplus_{k \in \mathbb{N}} V_k$ of almost integral type, we then obtain that both $\kappa_{a a_0}^{(2)}$ and $\kappa_{a a_0}^{(3)}$ are equal to $\dim \Delta(S^{(a a_0)})$. Our claim follows in the end as $\kappa_{a a_0}^{(2)} = \kappa^{(2)}$ and $\kappa_{a a_0}^{(3)} = \kappa^{(3)}$, thanks to (4.29). \square

In the special case when S is finitely generated,

$$\kappa^{(2)}(X, K_X + L, h_L) = \kappa^{(3)}(X, K_X + L, h_L)$$

can be obtained immediately as mentioned in the previous subsection. However, it is still in question that whether S defined as in (4.6) is finitely generated or not. In [21], the authors has reduced this problem to the study of the singularities of admissible Bergman metrics.

A dominating meromorphic map $f : X \dashrightarrow Y$ between a compact complex manifold X and a projective algebraic variety Y is determined by the subfield $K = Q(Y) \subset Q(X)$ up to bimeromorphic equivalence. For such f , there exist two

proper modifications $g' : X' \rightarrow X$ and $h' : Y' \rightarrow Y$ such that Y' is a projective manifold and the induced map $f' : X' \dashrightarrow Y'$ is actually a morphism. This is obtained by choosing X' as a non-singular model (whose existence is due to Hironaka, cf. Theorem 2.12 in [32]) of the graph of the meromorphic mapping $X \dashrightarrow Y'$ and g' as the projection onto the factor X . In this way, f' is called a representative of f or the function field K . Recall that f' is a fiber space if and only if K is algebraically closed in $Q(X)$ (cf. Corollary 1.10 in [32]).

We can now state the following main result of this subsection, whose main idea comes from (the proof of) Theorem 1.11 in [26].

Theorem 4.15. *Let X be a compact complex manifold and $L \rightarrow X$ be a pseudo-effective line bundle equipped with a singular metric h_L whose curvature current is semi-positive. Assume that h_L has analytic singularities (see its definition in section 1) and $\kappa(X, K_X + L, h_L) \geq 0$. Let $f^\flat : X^\flat \rightarrow Y^\flat$ be a representative of*

$$(4.35) \quad f(= \Phi_{|S_k|}) : X \dashrightarrow \text{Im } \Phi_{|S_k|} =: Y$$

for any $k \in N(X, K_X + L, h_L)$ sufficiently large so that Proposition 4.7 holds. Then $f^\flat : X^\flat \rightarrow Y^\flat$ is an analytic fiber space, $\kappa(X, K_X + L, h_L) = \dim Y^\flat$ and $\kappa(F, K_F + L_F, h_L|_F) = 0$, where F is the very general (which means, outside a countable union of analytic subsets) fiber of f^\flat .

Proof. According to Proposition 4.8, one may find $f^\flat : X^\flat \rightarrow Y^\flat$ as an algebraic fiber space and $\kappa^{(2)}(X, K_X + L, h_L) = \dim Y^\flat$. By the analytic singularities assumption on h_L , we see that by taking a resolution of singularities of $f^{b*}\varphi_L$ ($h_L = e^{-\varphi_L}$ locally) one may assume

$$(4.36) \quad f^{b*}\varphi_L \sim \alpha \sum \lambda_j \log |g_j|,$$

where $\alpha \in \mathbb{R}_{>0}$, g_j are local generators of the invertible sheaves $\mathcal{O}_X(-D_j)$ and $D = \sum \lambda_j D_j$ is a normal crossing divisor on X^\flat which also contains all the components of the exceptional divisors. Then $\mathcal{I}_k(f^{b*}\varphi_L)$ is computed by

$$(4.37) \quad \mathcal{I}_k(f^{b*}\varphi_L) = \mathcal{O}_{X'}(-\sum ([k\alpha\lambda_j] - k + 1)D_j)$$

and therefore locally free.

Let E_l denote the Cartier divisor on X^\flat associated to the invertible sheaf $\mathcal{O}_{X^\flat}(l(K_{X^\flat} + L^\flat) \otimes \mathcal{I}_l(f^{b*}h_L))$ and $l \in \mathbb{N}$. Let H be a very ample divisor on Y^\flat , we claim that

$$(4.38) \quad f^{b*}\mathcal{O}_{Y^\flat}(H) \subset \mathcal{O}_{X^\flat}(E_k)$$

holds with our choice of k .

Indeed, let $1, \sigma_1, \dots, \sigma_N$ be a basis of $H^0(Y^\flat, \mathcal{O}_{Y^\flat}(H)) = \{\sigma \in Q(Y^\flat); (\sigma) + H \geq 0\}$. Here, for a Cartier divisor D , " $D \geq 0$ " means all the coefficients of D is non-negative, i.e. D is effective. By our choice of k so that $Q((S_0[S_k])) = Q((X, K_X + L, h_L)) \supset Q(Y^\flat)$ (cf. Proposition 4.7), there exist $N > 0$ and $\xi_0, \dots, \xi_N \in H^0(X^\flat, \mathcal{O}_{X^\flat}(E_k))$ such that $\xi_0 \neq 0$ and $\sigma_j = \xi_j/\xi_0$, $j = 0, \dots, N$. Hence we get

$$(4.39) \quad 0 \leq (\xi_j) + E_k = f^{b*}((\sigma_j) + H) + ((\xi_0) + E_k - f^{b*}H).$$

Since H is very ample, we have $\bigcap_{j=1}^N f^{b*}((\sigma_j) + H) = \emptyset$; thus $(\xi_0) + E_k - f^{b*}H \geq 0$. Hence multiplying by ξ_0 gives $f^{b*}\mathcal{O}_{Y^\flat}(H) \subset \mathcal{O}_{X^\flat}(E_k)$, which confirms our claim.

Since $\kappa^{(2)}(X, K_X + L, h_L) \geq 0$, we have $H^0(X^\flat, k_1(K_{X^\flat} + L^\flat) \otimes \mathcal{I}_{k_1}(f^{b*}h_L)) \neq 0$ for some $k_1 \in N(X, K_X + L, h_L)$. It follows from the restriction map

$$H^0(X^\flat, k_1(K_{X^\flat} + L^\flat) \otimes \mathcal{I}_{k_1}(f^{b*}h_L)) \rightarrow H^0(X_y^\flat, k_1(K_{X_y^\flat} + L_y^\flat) \otimes \mathcal{I}_{k_1}(f^{b*}h_L)|_{X_y^\flat})$$

that $\kappa^{(2)}(F, K_F + L_F, h_L|_F) \geq 0$ for general fibers $F = X_y^\flat$, thanks to the fact that $\mathcal{I}_{k_1}(f^{b*}h_L|_{X_y}) = \mathcal{I}_{k_1}(f^{b*}h_L)|_{X_y}$ holds generally. Therefore, to prove $\kappa^{(2)}(F, K_F + L_F, h_L|_F) = 0$ holds for very general F , it only suffices to show the coherent sheaf

$$(4.40) \quad \mathcal{F}_a := f_*^\flat \mathcal{O}(a(K_{X^\flat} + L^\flat) \otimes \mathcal{I}_a(f^{b*}h_L)) =: f_*^\flat \mathcal{O}(E_a)$$

has rank at most 1 for every $a \in N(X, K_X + L, h_L)$.

Thanks to (4.38), there exist injections

$$(4.41) \quad \mathcal{O}(E_a) \otimes f^{b*} \mathcal{O}_{Y^\flat}(bH) \subset \mathcal{O}(E_a) \otimes \mathcal{O}(bE_k)$$

for any $b \in \mathbb{N}$ (since $\mathcal{O}(E_a)$ is locally free). By the basic relation $\mathcal{I}_k(f^{b*}h_L)^{\otimes b} \cdot \mathcal{I}_a(f^{b*}h_L) \subset \mathcal{I}_{a+kb}(f^{b*}h_L)$ and the left exactness of the functor f_*^\flat , (4.41) also induces injections

$$(4.42) \quad \mathcal{F}_a \otimes \mathcal{O}(bH) \hookrightarrow f_*^\flat \mathcal{O}((a+kb)(K_X + L) \otimes \mathcal{I}_k(f^{b*}h_L)^{\otimes b} \otimes \mathcal{I}_a(f^{b*}h_L)) \hookrightarrow \mathcal{F}_{a+kb}$$

for all b . One may obtain a commutative diagram

$$(4.43) \quad \begin{array}{ccc} H^0(Y^\flat, \mathcal{F}_a \otimes \mathcal{O}(bH)) & \hookrightarrow & H^0(Y^\flat, \mathcal{F}_{a+kb}) = H^0(X^\flat, \mathcal{O}(E_{a+kb})) \\ \downarrow \beta & & \downarrow \gamma \\ \mathcal{F}_a \otimes \mathcal{O}(bH) \otimes_{\mathcal{O}_{Y^\flat}} \mathcal{O}_{Y^\flat, y} / \mathfrak{m}_y & \hookrightarrow & \mathcal{F}_{a+kb} \otimes_{\mathcal{O}_{Y^\flat}} \mathcal{O}_{Y^\flat, y} / \mathfrak{m}_y \end{array}$$

from (4.42). Since $Q((X, K_X + L, h_L)) = Q(Y)$ (cf. Proposition 4.7), $\text{Im } \gamma$ is a vector space of dimension ≤ 1 . Since β is surjective for $b \gg 1$, thus \mathcal{F}_a has rank at most 1. \square

5. PROOF OF COROLLARY 1.6

In this section, for a holomorphic vector bundle E (resp. coherent sheaf \mathcal{E}), the notation $\mathbb{P}(E)$ (resp. $\mathbb{P}(\mathcal{E})$) stands for the projective fiber space containing all 1-dim quotients of fibers of E (resp. the set of all quotient invertible sheaves of \mathcal{E} , cf. (2.8) and (2.9) in [32]).

To give the proof of Corollary 1.6, we first consider some results on generalized Kodaira dimensions from the viewpoint of birational geometry. The following proposition is a slight modification of Theorem 5.11 in [32]. For completeness, we give details of its proof here.

Proposition 5.1. *Let $f : X \rightarrow Y$ be a surjective and proper holomorphic map between two compact complex manifolds X and Y with connected fibers. Let $L \rightarrow X$ be a \mathbb{Q} -line bundle equipped with a singular metric h_L with semi-positive curvature current. Then there exists a set $\Sigma \subset Y$ whose complement has measure zero, such that*

$$(5.1) \quad \kappa(X, K_X + L, h_L) \leq \kappa(F, K_F + L|_F, h_L|_F) + \dim Y,$$

where $F \subset f^{-1}(\Sigma)$ denotes the sufficiently general fiber of f .

Proof. Let us assume $\kappa(X, K_X + L, h_L) \neq -\infty$ without loss of generality. Then for sufficiently large and divisible $k \in \mathbb{N}$, we set

$$(5.2) \quad \mathcal{F}_k := f_*(k(K_X + L) \otimes \mathcal{I}_k(h_L)) \neq 0,$$

which is coherent over Y by Grauert's direct image theorem. Moreover, by Grauert's theorem (cf. Theorem 10.6 and Theorem 10.7 in [3]), there exists a Zariski open dense subset $U_k \subset Y_0$ such that $\mathcal{F}_k|_{U_k}$ is locally free and the following base change property

$$(5.3) \quad \mathcal{F}_{k,y} \otimes \mathcal{O}_{Y,y}/\mathfrak{m}_y = H^0(X_y, (kK_{X_y} + kL|_{X_y}) \otimes \mathcal{I}_k(h_L)|_{X_y})$$

holds for $y \in U_k$. As mentioned in section 2, we already obtain that

$$(5.4) \quad \mathcal{I}_k(h_L)|_{X_y} = \mathcal{I}_k(h_L|_{X_y})$$

for all $y \in \Sigma_k$ where $\Sigma_k \subset Y$ is a subset with full measure. Let $\mathbb{P}(\mathcal{F}_k)$ be the projective fiber space associated to \mathcal{F}_k , then $f|_{f^{-1}(U_k)}$ factors through a meromorphic map $h_k : X \dashrightarrow \mathbb{P}(\mathcal{F}_k)$ and a morphism $g_k : \mathbb{P}(\mathcal{F}_k) \rightarrow Y$, i.e. $f|_{f^{-1}(U_k)} = g_k \circ h_k$ (cf. (2.10) in [32]).

On one side, by (5.3) and (5.14), $h_k|_{X_y}$ for $y \in \Sigma_k \cap U_k$ is given by the following meromorphic map

$$(5.5) \quad \Phi_{k,y} : X_y \dashrightarrow \mathbb{P}H^0(X_y, (kK_{X_y} + kL|_{X_y}) \otimes \mathcal{I}_k(h_L|_{X_y}))$$

with respect to a *sublinear* series of the divisor $kK_{X_y} + L|_{X_y}$ (cf. (2.10) in [32]). It follows that

$$(5.6) \quad \dim h_k(X) = \dim Y + \dim \operatorname{Im} \Phi_{k,y}$$

holds for $y \in \Sigma_k \cap U'_k$, where $U'_k \subset U_k$ is another Zariski open dense subset (cf. Remark 5.1).

On the other side, identifying

$$(5.7) \quad H^0(Y, \mathcal{F}_k) = H^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L))$$

we obtain a meromorphic map

$$(5.8) \quad h : \mathbb{P}(\mathcal{F}_k) \dashrightarrow \mathbb{P}H^0(\mathbb{P}(\mathcal{F}_k), \mathcal{O}_{\mathbb{P}(\mathcal{F}_k)}(1)) = \mathbb{P}H^0(Y, \mathcal{F}_k) = \mathbb{P}^{N_k}$$

such that $h \circ h_k$ is the Kodaira meromorphic mapping Φ_k on X with respect to $|H^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L))|$, here $N_k = \dim H^0(X, k(K_X + L) \otimes \mathcal{I}_k(h_L)) - 1$.

Therefore, there is a generically surjective (cf. Definition 2.6 in [32]) meromorphic map $h : h_k(X) \dashrightarrow \operatorname{Im} \Phi_k$. Taking k sufficiently large and divisible, we conclude from (5.6) and Proposition 4.14 that

$$(5.9) \quad \begin{aligned} \kappa(X, K_X + L, h_L) &= \dim \operatorname{Im} \Phi_k \leq \dim h_k(X) = \dim Y + \dim \operatorname{Im} \Phi_{k,y} \leq \\ &\dim Y + \kappa(X_y, K_{X_y} + L|_{X_y}, h_L|_{X_y}) \end{aligned}$$

for $y \in \Sigma$. □

Remark 5.1. We might give some details of (5.6) as follows.

Let $\Gamma_k := \overline{h_k(f^{-1}(U_k))}$ be the closure of the image of h_k in $\mathbb{P}(\mathcal{F}_k)$, which is a complex analytic subspace of $\mathbb{P}(\mathcal{F}_k)$ (or the projection of the graph of h_k onto the second factor $\mathbb{P}(\mathcal{F}_k)$). When considering the generically surjective morphism $\tilde{g}_k : \Gamma_k \rightarrow Y$ induced by the restriction of g_k on Γ_k , there is a Zariski open dense subset $U'_k \subset U_k$ such that $\dim \tilde{g}_k^{-1}(y) = \dim \Gamma_k - \dim Y$ is constant for any $y \in U'_k$. Since $h_k|_{X_y}$ is given by (5.5) for any $y \in \Sigma_k \cap U'_k$, we conclude that

$$\dim \operatorname{Im} \Phi_{k,y} = \dim h_k(X_y) = \dim \tilde{g}_k^{-1}(y) = \dim \Gamma_k - \dim Y$$

holds for sufficiently general y .

The following lemma can be seen as a quantitative version of Lemma 3.1 in [35].

Lemma 5.2. *Let $f : X \rightarrow Y$ be a surjective holomorphic map between a compact complex manifold X and a projective manifold Y with connected fibers. Let $L \rightarrow X$ be a \mathbb{Q} -line bundle equipped with a singular metric h_L with semi-positive curvature current and $A \rightarrow Y$ be an ample \mathbb{Q} -line bundle. If $\kappa(X, K_{X/Y} + L, h_L) \geq 0$ and h_L has analytic singularities, then*

$$(5.10) \quad \kappa(X, K_{X/Y} + L + f^*A, h_L) = \kappa(F, K_F + L|_F, h_L|_F) + \dim Y,$$

where $F = X_y$ for sufficiently general $y \in Y$.

Proof. By multiplying a sufficiently large and divisible integer k_0 , we may assume that $H^0(X, (k_0 K_{X/Y} + k_0 L) \otimes \mathcal{I}_{k_0}(h_L)) \neq 0$ and $k_0 A$ is a very ample line bundle over Y .

Now we consider the meromorphic map

$$(5.11) \quad \Phi := \Phi|_{V_{k_0}} : X \dashrightarrow \mathbb{P}V_{k_0}$$

by enlarging k_0 so that Proposition 4.7 holds, where $V_{k_0} = H^0(X, (k_0 K_{X/Y} + k_0 L + k_0 f^*A) \otimes \mathcal{I}_{k_0}(h_L))$. Upon a modification $\mu : X' \rightarrow X$, where X' is obtained as a smooth model of the graph of the meromorphic map $\Phi|_{V_{k_0}}$, we obtain that

$$\Phi' := \Phi'|_{V'_{k_0}} : X' \rightarrow \text{Im } \Phi'|_{V'_{k_0}} = \text{Im } \Phi|_{V_{k_0}} \subset \mathbb{P}V'_{k_0} = \mathbb{P}V_{k_0}$$

is an analytic fiber space (i.e. Φ' is a representative of Φ , cf. Theorem 4.15), where $V'_{k_0} := H^0(X', k_0(K_{X'/Y} + L' + \mu^* f^* A) \otimes \mathcal{I}_{k_0}(h'_L)) = V_{k_0}$, $(L', h'_L) = (\mu^* L, \mu^* h_L)$.

We also obtain

$$\tilde{\kappa}(F, K_F + L|_F, h_L|_F) = \tilde{\kappa}(F', K'_F + L'|_{F'}, h'_L|_{F'})$$

$$(5.12) \quad \tilde{\kappa}(X, K_{X/Y} + L + f^*A, h_L) = \tilde{\kappa}(X', K_{X'/Y} + L' + (f')^*A, h_{L'})$$

by Proposition 4.9, where F (resp. F') is the general fiber of Φ (resp. Φ'). Additionally we have $\dim \text{Im } \Phi' = \tilde{\kappa}(X', K_{X'/Y} + L' + (f')^*A, h_{L'})$. Viewing $H^0(Y, k_0 A) = H^0(X', k_0(f')^*A) \hookrightarrow V'_{k_0}$, the following commutative diagram

$$(5.13) \quad \begin{array}{ccc} \mathbb{P}V'_{k_0} & \dashrightarrow & \mathbb{P}H^0(Y, k_0 A) \\ \Phi' \uparrow & & \uparrow \\ X' & \xrightarrow{f' \circ \mu} & Y \end{array}$$

holds. As a consequence of (5.13), the general fiber G' of Φ' is contracted by f' , hence we obtain a fiber space $\Phi'|_{F'} : F' \rightarrow \text{Im } \Phi'|_{F'}$ whose general fiber is G' (note F' is connected and G' is also connected thanks to Proposition 4.8). It is already known that $\mathcal{I}_{k_0}(h_{L'})|_{F'} = \mathcal{I}_{k_0}(h_{L'}|_{F'})$ holds for sufficiently general fibers F' . Therefore, $\Phi'|_{F'}$ is determined by a *sublinear* series of $|H^0(F', (k_0 K_{F'} + k_0 L'|_{F'}) \otimes \mathcal{I}_{k_0}(h_{L'}|_{F'}))|$, which contains elements that can be extended to V'_{k_0} .

On one hand, we obtain

$$\tilde{\kappa}(F', K_{F'} + L|_{F'}, h_{L'}|_{F'}) \geq \dim \text{Im } \Phi'|_{F'} = \dim F' - \dim G' =$$

$$(5.14) \quad \dim \text{Im } \Phi' - \dim Y = \tilde{\kappa}(X', K_{X'/Y} + L' + (f')^*A, h_{L'}) - \dim Y.$$

On the other hand, applying Proposition 5.1 to $\Phi'|_{F'}$ we get

$$\begin{aligned} \tilde{\kappa}(F', K_{F'} + L'|_{F'}, h_{L'}|_{F'}) &= \tilde{\kappa}(F', K_{F'} + (L' + (f')^*A)|_{F'}, h_{L'}|_{F'}) \leq \\ &\dim \text{Im } \Phi'|_{F'} + \tilde{\kappa}(G', K_{G'} + (L' + (f')^*A)|_{G'}, h_{L'}|_{G'}) = \end{aligned}$$

$$(5.15) \quad \dim \text{Im } \Phi'|_{F'} = \tilde{\kappa}(X', K_{X'/Y} + L' + (f')^*A, h_{L'}) - \dim Y.$$

To see why $\tilde{\kappa}(G', K_{G'} + (L' + (f')^*A)|_{G'}, h_{L'}|_{G'}) = 0$ in (5.15) (where we have used the assumption on h_L), one can just apply Theorem 4.15 to the generalized Iitaka fibration Φ' . In the end, our claim of (5.10) follows easily by (5.14), (5.15) and (5.12) and Proposition 4.14. \square

We are now in position to accomplish the proof of Corollary 1.6.

Proof. (Proof of Corollary 1.6) Let us first fix an ample bundle A over Y such that $A - K_Y$ is sufficiently ample and satisfies the assumption (2.10) in Lemma 2.4. Since K_Y is a big line bundle, we may write

$$(5.16) \quad kK_Y = A + E$$

for some large $k \in \mathbb{N}$ with an effective line bundle E . We may assume that $f_*((2kK_{X/Y} + 2kL) \otimes \mathcal{I}_{2k}(h_L)) \neq 0$, otherwise there is nothing to prove. Then Lemma 2.4 already implies that $\kappa(X, K_{X/Y} + L + \frac{1}{2k}f^*A, h_L) \geq 0$, hence one can apply Lemma 5.2 to obtain that

$$\begin{aligned} \kappa(X, K_X + L, h_L) &= \kappa(X, K_{X/Y} + L + f^*K_Y, h_L) \geq \\ \kappa(X, (K_{X/Y} + L + \frac{1}{2k}f^*A) + \frac{1}{2k}f^*A, h_L) &= \kappa(F, K_F + L|_F, h_L|_F) + \dim Y \end{aligned}$$

holds for general F . For the other direction, it will just be an immediate consequence of Proposition 5.1. \square

Acknowledgements: The authors would sincerely thank the anonymous referee(s) for his/her suggestions to the original version of this paper, especially for Remark 2.3. The first author would like to thank Prof. Z. W. Wang for some fruitful discussion about this paper and Dr. J. Y. Wang for introducing some background about the Iitaka conjecture and some of his work on this topic.

REFERENCES

- [1] B. Berndtsson and M. Păun. Bergman kernels and the pseudoeffectivity of relative canonical bundles. *Duke Math. J.*, **145**(2): 341–378, 2008.
- [2] B. Berndtsson and M. Păun. A Bergman kernel proof of the Kawamata subadjunction theorem. ArXiv0804.3884v2.
- [3] J. Bertin, J.-P. Demailly, L. Illusie and C. Peters. Introduction to Hodge theory. SMF/AMS Texts and Monographs 8.
- [4] F. Campana. Orbifolds, special varieties and classification theory. *Ann. Inst. Fourier (Grenoble)*, **54**(3): 499–630, 2004.
- [5] J.Y. Cao. Ohsawa-Takegoshi extension theorem for compact Kähler manifolds and applications. Volume **21**, pages 19–38. Cham., 2017. Springer INdAM Ser.
- [6] J.-P. Demailly. *Analytic methods in algebraic geometry*. Surv. Mod. Math. 1, International Press, Somerville, 2012.
- [7] J.-P. Demailly. Extension of holomorphic functions defined on non reduced analytic subvarieties. In *The legacy of Bernhard Riemann after one hundred and fifty years. Vol. I*, volume 35 of *Adv. Lect. Math. (ALM)*, pages 191–222. Int. Press, Somerville, MA, 2016.
- [8] J.-P. Demailly. Extension of holomorphic functions defined on non reduced analytic subvarieties.
- [9] J.-P. Demailly, T. Peternell and M. Schneider. Pseudo-effective line bundles on compact Kähler manifolds. *Internat. J. Math.*, **12**(6): 689–741, 2001.
- [10] F.S. Deng, Z.W. Wang, L.Y. Zhang and X.Y. Zhou. New characterizations of plurisubharmonic functions and positivity of direct image sheaves. ArXiv: 1809.10371v1, to appear in *American J. Math.*.
- [11] Y. Deng. Applications of the Ohsawa-Takegoshi extension theorem to direct image problems. ArXiv: 1703.07279.

- [12] O. Fujino. Foundations of the minimal model program. Volume **35** of *MSJ Memoirs*, Mathematical Society of Japan, Tokyo, 2017.
- [13] O. Fujino. Iitaka conjecture: An introduction. *Springer Briefs in Mathematics*, Singapore: Springer, 2020.
- [14] O. Fujino. Notes on the weak positivity theorems. In Algebraic varieties and automorphism groups, volume **75**, *Adv. Stud. Pure Math.*, pages 73–118. *Math. Soc. Japan*, Tokyo, 2017.
- [15] O. Fujino. On subadditivity of the logarithmic Kodaira dimension. *J. Math. Soc. Japan*, **69**(4): 1565–1581, 2017.
- [16] O. Fujino. Corrigendum to "On subadditivity of the logarithmic Kodaira dimension". *J. Math. Soc. Japan*, **72**(4): 1181–1187, 2020.
- [17] H. Grauert and R. Remmert. Coherent analytic sheaves. Grundlehren Math. Wiss., **265**, Springer-Verlag, 1984.
- [18] Q.A. Guan and X.Y. Zhou. A proof of Demailly's strong openness conjecture. *Ann. of Math.*, **182**(2): 605–616, 2015.
- [19] Q.A. Guan and X.Y. Zhou. A solution of an L^2 extension problem with an optimal estimate and applications. *Ann. of Math.*, **181**(2): 1139–1208, 2015.
- [20] C. Hacon, M. Popa and C. Schnell. Algebraic fiber spaces over abelian varieties: around a recent theorem by Cao and Păun. *Contemp. Math.*, **712**: 143–195, 2018.
- [21] B.J. He and X.Y. Zhou. Metrics on twisted pluricanonical bundles and finite generation of twisted canonical rings. *Pure Appl. Math. Q.*, **19**(2): 451–485, 2023.
- [22] K. Kaveh and A. G. Khovanskii. Newton-Okounkov bodies, semigroups of integral points, graded algebras and intersection theory. *Ann. of Math. (2)*, **176**: 925–978, 2012.
- [23] D. Kim. L^2 extension of holomorphic functions for log canonical pairs. *J. Math. Pures Appl.*, **177**: 198–213, 2023.
- [24] R. Lazarsfeld. *Positivity in Algebraic Geometry I and II*. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, 48–49, Springer-Verlag, Berlin, 2004.
- [25] R. Lazarsfeld and M. Mustata. Convex bodies associated to linear series. *Ann. Sci. Éc. Norm. Supér.*, **42**: 783–835, 2009.
- [26] S. Mori. Classification of higher-dimensional varieties. In Algebraic geometry, Bowdoin, Proc. Sympos. Pure Math. 46, Part 1, pages 269–331. *Amer. Math. Soc.*, 1985.
- [27] N. Nakayama. *Zariski-decomposition and Abundance*. MSJ Memoirs, 14, Math. Soc. Japan, Tokyo, 2004.
- [28] T. Ohsawa. *On the extension of L^2 holomorphic functions. IV. A new density concept*. Geometry and analysis on complex manifolds, 157–170, World Sci. Publ., River Edge, NJ, 1994.
- [29] A. Okounkov. Brunn-Minkowski inequality for multiplicities. *Invent. Math.*, **125**: 405–411, 1996.
- [30] M. Păun and S. Takayama. Positivity of twisted relative pluricanonical bundles and their direct images. *J. Algebraic Geom.*, **27**(2): 211–272, 2018.
- [31] Y.-T. Siu. Invariance of plurigenera and torsion-freeness of direct image sheaves of pluricanonical bundles. Finite or infinite dimensional complex analysis and applications. Advances in Complex Analysis and its Applications 2. Dordrecht: Kluwer Academic Publishers, 2004: 45–83.
- [32] K. Ueno. *Classification theory of algebraic varieties and compact complex spaces*. Lecture Notes in Mathematics 439, Springer-Verlag, Berlin-New York, 1975.
- [33] E. Viehweg. Weak positivity and the additivity of Kodaira dimension for certain algebraic fiber spaces. *Adv. Studies in Pure Math. 1, Algebraic Varieties and Analytic Varieties*, pages 329–353, 1983.
- [34] E. Viehweg. Weak positivity and the additivity of the Kodaira dimension. II. The local Torelli map. *Progr. Math.*, **39**: 567–589, 1983.
- [35] J.Y. Wang. On the Iitaka conjecture $C_{n,m}$ for Kähler fiber spaces. *Ann. Fac. Sci. Toulouse Math. (6)*, **30**(4): 813–897, 2021.
- [36] X.J. Wu. Note on asymptotic behaviour of the canonical ring. arXiv:2209.02759v1.
- [37] X.Y. Zhou and L.F. Zhu. An optimal L^2 extension theorem on weakly pseudoconvex Kähler manifolds. *J. Differential Geom.*, **110**(1): 135–186, 2018.
- [38] X.Y. Zhou and L.F. Zhu. Optimal L^2 extension of sections from subvarieties in weakly pseudoconvex manifolds. *Pacific J. Math.*, **309**(2): 475–510, 2020.
- [39] X.Y. Zhou and L.F. Zhu. Siu's lemma, optimal L^2 extension and applications to twisted pluricanonical sheaves. *Math. Ann.*, **377**: 675–722, 2020, no.1-2.

- [40] X.Y. Zhou and L.F. Zhu. Subadditivity of generalized Kodaira dimensions and extension theorems. *Internat. J. Math.*, **31**:36 pp, 2020, no.12.

BOJIE HE: SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING, 100871, P. R. CHINA

Email address: `hbj@amss.ac.cn`

XIANGYU ZHOU: INSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, P. R. CHINA

Email address: `xyzhou@math.ac.cn`