Explicit Orthogonal Arrays and Universal Hashing with Arbitrary Parameters

Nicholas Harvey Arvin Sahami

Abstract

Orthogonal arrays are a type of combinatorial design that were developed in the 1940s in the design of statistical experiments. In 1947, Rao proved a lower bound on the size of any orthogonal array, and raised the problem of constructing arrays of minimum size. Kuperberg, Lovett and Peled (2017) gave a non-constructive existence proof of orthogonal arrays whose size is near-optimal (i.e., within a polynomial of Rao's lower bound), leaving open the question of an algorithmic construction. We give the first explicit, deterministic, algorithmic construction of orthogonal arrays achieving near-optimal size for all parameters. Our construction uses algebraic geometry codes.

In pseudorandomness, the notions of t-independent generators or t-independent hash functions are equivalent to orthogonal arrays. Classical constructions of t-independent hash functions are known when the size of the codomain is a prime power, but very few constructions are known for an arbitrary codomain. Our construction yields algorithmically efficient t-independent hash functions for arbitrary domain and codomain.

1 Introduction

Orthogonal arrays are a concept in the design of statistical experiments, first proposed by C. R. Rao in the 1940s. A detailed exposition of this subject can be found in reference books [12] [30]. An orthogonal array with parameters [s, m, n, t] is a matrix with s rows, m columns and entries in $[n] = \{1, \ldots, n\}$ such that, for every set of t columns, those columns contain each tuple in $[n]^t$ exactly s/n^t times. A long history of research considers the problem of, given m, n and t, finding an orthogonal array with small s.

If n is a prime power, a classical construction due to Bush¹ in 1952 [12, Theorem 3.1] [2, §3] gives an orthogonal array with exactly $s = n^t$ rows if $m \le n$. This is exactly optimal (the definition requires that s/n^t is a positive integer). For m > n, a simple modification of this construction has $s \le (nm)^t$ rows; however this is no longer optimal.

Indeed, a lower bound of Rao^2 [12, Theorem 2.1] implies that every orthogonal array has

$$s \geq \binom{m}{\lfloor t/2 \rfloor} (n-1)^{\lfloor t/2 \rfloor} \geq (cmn/t)^{ct}, \tag{1}$$

¹This is essentially equivalent to a Reed-Solomon code, and predates it by eight years.

²For the case n = 2, this can be found in [5, Section 8]. In the special case of linear orthogonal arrays, Rao's bound is equivalent to the dual of the generalized Hamming bound on codes, and predates it by three years.

for some constant c > 0. In contrast, Bush's upper bound lacks the denominator of t, and only applies when n is a prime power. In general, there are very few constructions of orthogonal arrays when n is not a prime power; see [12, Table 12.6].

Some important problems are the constructions of these arrays with the optimum values of m for a given t and s.

Rao, 1947 [24]

Research Problem 2.33. For fixed values of n and t, and large m, how far are the Rao bounds from the truth?

Hedayat, Sloane, Stufken 1999 [12]

(Note: notation has been adjusted to match ours.)

Previous constructions. One might imagine that a simple product construction can reduce the general case to the prime power case. This is possible (and was known to Bose in 1950 [1, Section 2]) but it is suboptimal. First factor n into a product of prime powers $\prod_{\ell \leq d} p_{\ell}^{a_{\ell}}$. For each ℓ , build an orthogonal array $A^{(\ell)}$ which in which n is taken to be $p_{\ell}^{a_{\ell}}$. Then, define A to be an entry-wise Cartesian product of $A^{(1)}, \ldots, A^{(d)}$. This gives an orthogonal array, but it is unfortunately somewhat large. Applying Rao's lower bound separately to each $A^{(\ell)}$, the number of rows of A is at least $\prod_{\ell \leq d} (cmp_{\ell}^{a_{\ell}}/t)^{ct} = (cm/t)^{cdt}n^{ct}$. This does not asymptotically match Rao's lower bound for all parameters due to the factor d in the exponent. So this construction cannot in general prove that Rao's lower bound is tight. Note that d can be $\Omega(\log(n)/\log\log n)$ when n is the product of the first d primes.

If n is a prime power, then it is known how to explicitly construct orthogonal arrays with $s \leq (Cmn/t)^{Ct}$ for some constant C. We will call such a construction *near-optimal*, meaning that it matches Rao's lower bound up to the value of the constant C. The anonymous reviewers of this manuscript have informed us of this result, which is stated in [4, Theorem 3.4] using the language of pseudorandom generators. That construction, like our results in Section 4, is based on the use of algebraic geometry codes.

If n is not restricted to be a prime power, then Kuperberg, Lovett and Peled [19] give a non-explicit construction, for all m, n, t, of orthogonal arrays that are also near-optimal, so $s \leq (Cmn/t)^{Ct}$ for some constant C. However their approach is randomized, and they only prove an exponentially small lower bound on the probability of success. One of their open questions is whether an algorithmic version of their construction exists [19, pp. 968].

Our results. We also give a *near-optimal* construction of orthogonal arrays. Our construction is explicit, algorithmic, works for all m, n, t, and is apparently unrelated to the construction of Kuperberg, Lovett and Peled.

Theorem 1 (Informal). For any m, n, t (with n not necessarily a prime power), there is an explicit description of an orthogonal array with parameters [s, m, n, t] where $s = (cmn/t)^{35t}$, and c is a universal constant. The array can be constructed by a deterministic algorithm with runtime poly(sm).

To prove this, we give an abstract construction based on coding theory. In Section 3 we instantiate that construction with Reed-Solomon codes, obtaining a simple, easily implementable, deterministic construction with $s = O(m+n)^{6t}$. In Section 4 we instantiate the construction instead with algebraic geometry codes to obtain a deterministic, near-optimal construction with $s = O(mn/t)^{35t}$. In Section 6 we instantiate the construction instead with random linear codes to obtain a non-explicit construction that has near-optimal size, and can be efficiently constructed by a randomized algorithm.

1.1 Hash Functions

Hash functions of various sorts are crucial tools in pseudorandomness and randomized algorithms. In this work we focus on *t-independent* functions. Let $h : [m] \to [n]$ be a random function with some distribution. We say that $h : [m] \to [n]$ is *t*-independent if

$$\Pr[h(a_1) = \alpha_1 \wedge \dots \wedge h(a_t) = \alpha_t] = \frac{1}{n^t}$$

$$\forall \text{ distinct } a_1, \dots, a_t \in [m], \ \forall \alpha_1, \dots, \alpha_t \in [n]$$

(Obviously $t \leq m$ is necessary.) This is equivalent to the random variables $\{h(a) : a \in [m]\}$ being uniformly distributed and t-wise independent. If this property is satisfied by a function h that is uniformly chosen from a multiset \mathcal{H} , then we also say that \mathcal{H} is t-independent. This property is also called *strongly universal*_t, or simply *strongly universal* if t = 2.

Wegman and Carter [34, Section 1] gave a simple construction of a *t*-independent family \mathcal{H} when *n* is a prime power³ and m = n. In this construction, \mathcal{H} corresponds to the polynomials of degree t - 1 over the field \mathbb{F}_n , so $|\mathcal{H}| = n^t$ and the space required to represent a member of \mathcal{H} is $O(t \log n)$ bits. In general, if *n* is a prime power and *m* is arbitrary, then a modified construction has $|\mathcal{H}| = (mn)^{O(t)}$, so members of \mathcal{H} can be represented in $O(t \log(mn))$ bits.

Orthogonal arrays and t-independent hash families are mathematically equivalent notions, as was observed by Stinson⁴ [27, Theorem 3.1] [28, Theorem 5.2].

Claim 2. Let M be an $s \times m$ matrix with entries in [n]. Let $\mathcal{H} = \{h_1, \ldots, h_s\}$ be a multiset of $[m] \to [n]$ functions where $h_i(j) = M_{i,j} \ \forall j \in [m]$. Then \mathcal{H} is t-independent iff M is a [s, m, n, t] orthogonal array.

The proof is immediate from the definitions. Thus Wegman and Carter's construction of t-independent hash functions is identical to Bush's 1952 construction of orthogonal arrays.

There are some differences in the goals of the research communities working on orthogonal arrays and t-independent hashing. One difference is computational. Researchers in orthogonal arrays tend to be interested in showing existence, or explicit construction, of the entire matrix M. In contrast, researchers in hash functions are additionally interested in quickly sampling a hash function from \mathcal{H} and quickly evaluating the hash function. The following theorem states our main result for hash functions.

Theorem 3 (Informal). For any parameters m, n, t, there exists a t-independent hash family for which functions can be represented in $O(t \log(mn))$ bits, can be evaluated in time O(t), and can be constructed in expected time $O(t + (n + m)^{\epsilon})$ for any $\epsilon > 0$. Assuming the generalized Riemann hypothesis (GRH), the expected construction time improves to O(t) + polylog(nm).

³Carter and Wegman also defined the notion of *weakly* universal hash functions, and gave a construction that allows n to be an arbitrary integer [3, Proposition 7].

⁴Another connection between design theory and derandomization is in Karp and Wigderson [16, §9].

In contrast, the Wegman-Carter construction requires that n is a prime power, also uses $O(t \log(nm))$ bits, also can be evaluated in time O(t), and can be constructed in expected time O(t) + polylog(nm). (If m > n, the construction must work in a field extension of \mathbb{F}_n of size at least m, so random sampling is used to find an irreducible polynomial.) Thus the efficiency of our hash function matches theirs, assuming GRH.

1.2 Applications

There are many algorithms that require t-independent hash functions for t > 2. Examples include cuckoo hashing [6], distinct element estimation [15], MinHash [13], and the leftover hash lemma, which is used to construct seeded extractors [29], etc. More recently, a mechanism for maximizing Nash social welfare in the percentage fee model uses t-independent hash functions where n not a prime power [7].

2 The abstract construction using codes

In this section we present an abstract construction of orthogonal arrays, even when n is not a prime power, using linear codes over finite fields. The subsequent sections instantiate this construction using particular codes. The Hamming distance between vectors x and y is denoted $\Delta(x, y) = |\{ i : x_i \neq y_i \}|.$

Lemma 4. Let m, n, t be integers with $n \ge 2$ and $2 \le t \le m$. Let q be a prime power satisfying $q \equiv 1 \pmod{n}$. Suppose that $C \subseteq \mathbb{F}_q^m$ is a linear code of dimension k whose dual has minimum distance at least t + 1. Let $b \in \mathbb{F}_q^m$ satisfy $\Delta(b, u) \ge m - \tau$ for all $u \in C$. Then BUILDOA, shown in Algorithm 1, will produce an [s, m, n, t] orthogonal array with $s = n^{\tau}q^k$.

An intuitive explanation of this lemma is as follows. If $u \in C$ were chosen uniformly at random, our hypotheses on C would imply that the coordinates of u will be uniformly distributed and t-wise independent. Thus, if for each $u \in C$, we insert u as a new row in the array M, then the columns of M would also be uniformly distributed and t-wise independent, which is equivalent to M being an orthogonal array. The catch, of course, is that the entries of u take values in \mathbb{F}_q , whereas the entries of M must take values in [n]. If q were a multiple of n, then this problem would be easily resolved: we could instead insert u modulo n (entrywise), as a new row in M. The resulting matrix M is easily seen to be an orthogonal array. More generally, the same construction would work using any $\frac{q}{n}$ -to-1 map $f : \mathbb{F}_q \to [n]$ applied entry-wise to u to create a row of M.

However, the main scenario of interest is the one in which n is a composite number, so q (being a prime power) certainly cannot be a multiple of n. Instead, we have $q \equiv 1 \pmod{n}$, so we require a new approach to create rows of M from the vectors $u \in C$. The idea is to choose a "bad value" β to eliminate from \mathbb{F}_q , so that the number of remaining "good values" is a multiple of n, and these good values can be mapped to integers in [n] in a way that preserves the uniform distribution. This mapping, which we denote ϕ_{β} , between good values and [n] can be completely arbitrary, except that it needs to be a $\frac{q-1}{n}$ -to-1 map in order to ensure uniformity. For example, if we identify the elements in \mathbb{F}_q with the integers in $\{0, \ldots, q-1\}$, then one may check that

$$\phi_{\beta}(x) = 1 + \left(\left((x + q - 1 - \beta) \mod q \right) \mod n \right) \tag{2}$$

Algorithm 1 BUILDOA uses the code C and vector b to construct an orthogonal array M. Here ϕ_{β} is an arbitrary $\frac{q-1}{n}$ -to-1 map from $\mathbb{F}_q \setminus \{\beta\}$ to [n], as in (2).

1: function BUILDOA (integer m, n, t, q, τ , code C, vector b) 2: Let M be a matrix with m columns and no rows for $u \in C$ do 3: Let $Z = \{z_1, \ldots, z_\ell\} \subseteq [m]$ 4: be the set of indices with $u_{z_i} = b_{z_i}$ 5:for $v \in [n]^{\ell}$ do 6: $\triangleright v$ specifies how to fix all bad values in u 7: Define the vector $w(u, v) \in [n]^m$ by 8: 9: $w(u,v)_j = \begin{cases} v_i & \text{if } j = z_i \text{ for some } i \in [\ell] \\ \phi_{b_i}(u_j) & \text{otherwise, i.e., if } u_j \neq b_j \end{cases}$ Add $n^{\tau-\ell}$ copies of the row vector w(u, v)10: to the matrix M11: end for 12:end for 13:14: end function

is indeed a $\frac{q-1}{n}$ -to-1 map from $\mathbb{F}_q \setminus \{\beta\}$ to [n].

For future purposes, it will be convenient to allow this bad value to depend on the coordinate. Using the given vector b, we let b_j be the bad value for coordinate j. To eliminate these bad values from u, we must find all indices j with $u_j = b_j$ and remap u_j to a new value. Intuitively, we would like to do so uniformly at random. However the construction must be deterministic, so instead we create multiple copies of u in which each such u_j has been replaced with every possible value in [n]. In Algorithm 1, the vector v specifies how the bad entries of u will be fixed, whereas the vector w(u, v) is the modified copy of u that incorporates all the fixes and maps all entries to [n]. To control the number of copies that are created, we must bound the number of bad indices. This is ensured by the condition $\Delta(b, u) \ge m - \tau$: every $u \in C$ has at most τ bad indices. Lastly, to ensure that every two codewords in C generate the same number of rows in M, we add $n^{\tau-\ell}$ copies of each modified vector w(u, v), where $\ell = m - \Delta(b, u)$ is the number of indices at which u and b agree.

Proof (of Lemma 4). Let us view C as a multiset \mathcal{H}_q of functions mapping [m] to [q] (cf. Claim 2). Since the dual of C has minimum distance at least t + 1, it follows that \mathcal{H}_q is t-independent [10, Exercise 2.13] [30, Theorem 10.17].

BUILDOA produces a matrix M with m columns, where each entry is in [n]. Let s be the total number of rows added. Since each vector $u \in C$ produces exactly n^{τ} rows in M, and C has dimension k, we have

$$s = n^{\tau} |C| = n^{\tau} q^k,$$

as required. By Claim 2 again, we will view M as a multiset $\mathcal{H} = \{h_1, \ldots, h_s\}$ of $[m] \to [n]$ functions. It remains to show that \mathcal{H} is *t*-independent.

Consider a function $h \in \mathcal{H}_q$. For a sequence $y \in [n]^m$, define the function $f'_{h,y}: [m] \to [n]$ by

$$f_{h,y}'(j) = \begin{cases} \phi_{b_j}(h(j)) & \text{if } h(j) \neq b_j \\ y_j & \text{if } h(j) = b_j. \end{cases}$$

Here the "bad" hash values are replaced with entries of y, and the other hash values are mapped by ϕ into [n]. Let \mathcal{H}' be the multiset made by adding one copy of $f'_{h,y}$ (counting multiplicities) for each $y \in [n]^m$ and $h \in \mathcal{H}_q$, i.e.,

$$\mathcal{H}' = \left\{ f'_{h,y} : y \in [n]^m, \ h \in \mathcal{H}_p \right\}.$$

Note that each function $h \in \mathcal{H}_q$ gives rise to exactly n^m functions in \mathcal{H}' , although they are not necessarily unique. It is clear that \mathcal{H}' is the family made by repeating each function in \mathcal{H} exactly $n^{m-\tau}$ times. Therefore \mathcal{H}' is t-independent if and only if \mathcal{H} is t-independent.

We will show that \mathcal{H}' is t-independent by showing that, for any fixed $i_1, \ldots, i_t \in [m]$, the random variables $h'(i_1), \ldots, h'(i_t)$ are independent and uniformly distributed over [n] when $h' \in \mathcal{H}'$ is chosen uniformly at random. Note that by definition of \mathcal{H}' , uniformly sampling a function h' from \mathcal{H}' is (in distribution) equivalent to uniformly sampling a function $h \in \mathcal{H}_q$, and a sequence $N = (N_1, \ldots, N_m)$ of independent and uniformly distributed random variables in [n], and then returning $h' = f'_{h,N}$.

To show that $h'(i_1), \ldots, h'(i_t)$ are t-wise independent⁵, we will show that they are a deterministic function of random variables that are themselves t-wise independent. That is, the pairs

$$(h(i_1), N_{i_1}), \ldots, (h(i_t), N_{i_t})$$

are easily seen to be t-wise independent. We obtain the values $h'(i_1), \ldots, h'(i_t)$ by applying the function $f: \mathbb{F}_q \times [n] \times \mathbb{F}_q \to [n]$, defined by

$$f(x, r, \beta) = \begin{cases} \phi_{\beta}(x) & \text{if } x \neq \beta \\ r & \text{if } x = \beta \end{cases}$$

Notice that by definition of $f'_{h,N}$, we have

$$h'(i_j) = f(h(i_j), N_{i_j}, b_{i_j}), \quad \forall j \in [t].$$

Since f and b are deterministic, it follows that $h'(i_1), \ldots, h'(i_t)$ are t-wise independent.

To complete the proof that \mathcal{H}' is *t*-independent, we must also show that each h'(i) is uniformly distributed. To see this, note that for each $i \in [m]$ and $\nu \in [n]$

$$\Pr[h'(i) = \nu] = \Pr[h(i) = b_i \wedge N_i = \nu] + \Pr[h(i) \in \phi_{b_i}^{-1}(\nu)]$$
$$= \frac{1}{q} \cdot \frac{1}{n} + \frac{(q-1)}{nq} = \frac{1}{n}$$

since h(i) is uniformly distributed, and ϕ is a $\frac{q-1}{n}$ -to-1 map.

To summarize, for any $i_1, \ldots, i_t \in [m], h'(i_1), \ldots, h'(i_t)$ are independent and uniformly distributed random variables over [n]. That is, \mathcal{H}' is t-independent. As previously argued, this implies that \mathcal{H} is t-independent, which is what we intended to prove.

⁵Here the term "t-wise independent" has the meaning from probability theory, where it does not imply that the random variables are uniform. We use the term "t-independent" to indicate the meaning in pseudorandomness, where the random variables must additionally be uniformly distributed.

Corollary 5. Let m, n, t be integers with $n \ge 2$ and $2 \le t \le m$. Let q be a prime power satisfying $q \equiv 1 \pmod{n}$. Let $C_1 \subsetneq C_2 \subseteq \mathbb{F}_q^m$ be linear codes such that the dual of C_1 has minimum distance at least t+1. Let k be the dimension of C_1 and d_2 be the minimum distance of C_2 . Then there is an algorithm to produce an [s, m, n, t] orthogonal array with $s = n^{m-d_2}q^k$.

Proof. Pick any $b \in C_2 \setminus C_1$. Then $\Delta(b, u) \ge d_2$ for all $u \in C_1$. Thus we may apply Lemma 4, taking $C := C_1$ and $\tau := m - d_2$.

3 Construction using Reed-Solomon codes

In this section, we instantiate the abstract bound using Reed-Solomon codes.

Theorem 6. There exists a universal constant c > 0 such that the following is true. For any integers $n, m, t \ge 2$ with $t \le m$, there exists an explicit [s, m, n, t] orthogonal array with

$$s \leq n^t \cdot (c(n+m))^{5t}$$
.

Assuming the generalized Riemann hypothesis,

$$s \leq n^t ((n+m)\ln(n+m))^{2t}.$$

This bound matches (up to constants in the exponent) the non-constructive bound in the conference article of Kuperberg, Lovett and Peled [18]. This bound is not near-optimal (i.e., $(mn/t)^{O(t)})$ for all parameters, but it is in many regimes. (For example, it is near-optimal if $t \leq \min\{n, m\}$, or $t \leq m^{0.99}$, or $n \geq m^{0.01}$. Also, if $t = \Omega(m)$, then the trivial bound of $s \leq n^m$ is near-optimal.) Section 4 presents an improved construction that is near-optimal for all parameters, matching the non-constructive results of the journal article [19].

The prime. The first step of the proof is to find a prime p satisfying $p \equiv 1 \pmod{n}$ and p > m. We will use the following result.

Theorem 7 (Linnik's theorem). There exist universal constants $L, c_L > 0$ such that the following is true. For any integer $n \ge 2$, and any positive integer a coprime with n, there exists a prime p satisfying (i) $p \equiv a \pmod{n}$, and (ii) $p \le c_L \cdot n^L$.

The constant L is known as Linnik's constant. Although Linnik in 1944 did not provide an explicit value for L, the most recent developments. have shown that $L \leq 5.5$ in 1992 [11], L < 5.2 in 2011 [35], and $L \leq 5$ in [36, Theorem 2.1]. Assuming the generalized Riemann hypothesis [20], the conclusion (ii) can be strengthened to $p \leq (n \ln n)^2$.

Corollary 8. For any constants $L, c_L > 0$ satisfying Linnik's theorem, the following is true. For any integer $n \ge 2$ and real $m \ge 2$, there exists a prime p satisfying (i) $p \equiv 1 \pmod{n}$, (ii) $p \le c_L(n+m)^L$, and (iii) m < p.

Proof. Let $\eta = n \cdot \lceil m/n \rceil$. Note that $m \leq \eta < m+n$. Let p be the prime obtained by Theorem 7 with η instead of n, and with a = 1 (which is trivially coprime with n). The theorem ensures that $\eta \mid p-1$; since $n \mid \eta$, we have $n \mid p-1$, which establishes (i). Since $\eta \mid p-1$ and p-1 > 0, we must have $m \leq \eta < p$, which establishes (ii). Lastly, we have $p \leq c_L \cdot \eta^L < c_L \cdot (m+n)^L$, which establishes (ii).

A prime p satisfying the conditions of Corollary 8 can be found by exhaustive search. Simply test the primality of all integers p satisfying $p \equiv 1 \pmod{\eta}$ and $p \leq c_L \cdot \eta^L$, of which there are at most η^{L-1} . Since the primality of x can be tested in $O(\log^7 x)$ time, the overall runtime is $O(\eta^{L-1}\log^7 \eta) = \tilde{O}((n+m)^4)$.

The codes. The next step of the proof is to find codes of appropriate parameters that can be used with Corollary 5. Let q equal the prime p chosen above. We will use Reed-Solomon codes in \mathbb{F}_q^m . The following theorem states their basic properties, with apparently excessive detail, in order to draw a parallel with Theorem 11 below.

Theorem 9. Let $m \leq q$. There exists a sequence of linear codes $C_0, \ldots, C_{m-1} \subseteq \mathbb{F}_q^m$, where C_a has parameters $[m, k_a, d_a]_q$, such that the following statements hold.

 $C_a \subsetneq C_{a+1} \quad \forall \ 0 \le a < m-1 \tag{3}$

$$d_a = m - a \quad \forall \ 0 \le a < m \tag{4}$$

$$k_a = a + 1 \quad \forall \ 0 \le a < m \tag{5}$$

$$k_a^{\perp} + d_a^{\perp} = m + 1 \quad \forall \ 0 \le a < m \tag{6}$$

Here k_a^{\perp} and d_a^{\perp} respectively denote the dimension and distance of C_a^{\perp} , the dual of C_a .

Proofs of the claims in this theorem may be found in standard references, e.g., [10, Claim 5.2.1] [25, Proposition 11.4] [31, Exercise 1.1.20].

We will apply Corollary 5 to the codes $C_{t-1} \subsetneq C_t$. It follows from (6) that

$$m+1 = k_{t-1}^{\perp} + d_{t-1}^{\perp} = (m-k_{t-1}) + d_{t-1}^{\perp},$$

and therefore

$$d_{t-1}^{\perp} = k_{t-1} + 1 = t + 1.$$

Thus Corollary 5 yields an [s, m, n, t] orthogonal array with

$$s = n^{m-d_t} p^{k_{t-1}} = n^t p^t \le n^t \left(c_L (n+m)^5 \right)^t.$$
(7)

This proves Theorem 6.

4 Construction using algebraic geometry codes

In this section we replace the Reed-Solomon codes with algebraic geometry codes, which gives a near-optimal orthogonal array for all parameters. This matches the non-constructive bound shown in the journal article [19], and proves Theorem 1.

Theorem 10. There exists a universal constant c > 0 such that the following is true. For any integers $n, m, t \ge 2$ with $t \le m$, there exists an [s, m, n, t] orthogonal array with

$$s \leq n^{5t} (c(n+11+m/t)^5)^{6t}$$

Moreover, this array can be constructed by a deterministic algorithm in poly(sm) time.

To motivate the proof, let us reflect on the construction of the previous section. A standard bound on all [m, k, d] linear codes is the *Singleton bound*, which states that $k + d \le m + 1$; if equality holds it is called an *MDS code*. In Section 3, to apply Corollary 5 we needed $d_{t-1}^{\perp} \ge t + 1$. Furthermore, to obtain an exponent of O(t) in (7), we wanted $d_t \ge m - O(t)$ and $k_{t-1} = O(t)$. Since Reed-Solomon codes and their duals are MDS codes, we have exactly $d_{t-1}^{\perp} = t + 1$, $d_t = m - t$ and $k_{t-1} = t$. This yields an orthogonal array with $s = n^t p^t$ rows, with the main shortcoming being the undesirably large field size of p = poly(n + m).

The construction of this section attains an improved bound by reducing the field size to poly(n + m/t). We cannot hope to use MDS codes anymore, since it is believed that they do not exist when the field size is less than m-1. Instead, we will use codes that only approximately satisfy the Singleton bound, but do so over a much smaller field. Algebraic geometry codes (AG codes) provide a tradeoff suitable for our purposes.

Algebraic geometry codes. These are a general class of codes that involve evaluations of rational functions over algebraic curves [9]. Reed-Solomon codes are a simple special case using evaluations of polynomials. We begin with a detailed statement of the codes that can be constructed from a general curve.

Let q be a prime power. Let X be a curve over \mathbb{F}_q (see [31, §2.1.2]). Let g be its genus (see [31, pages 87 and 91]). Let N be its number of rational points (see [31, pp. 127], [21, Definition VII.6.14], or [23, Section A.2]).

Theorem 11. Let m < N. There exists a sequence of linear codes $C_0, \ldots, C_{m-1} \subseteq \mathbb{F}_q^m$, where C_a has parameters $[m, k_a, d_a]_q$, such that the following statements hold.

$$C_a \subseteq C_{a+1} \quad \forall \ 0 \le a < m-1$$
[31, Remark 4.1.7]
(8)

$$d_a \ge m - a \quad \forall \ 0 \le a < m$$
[31, Theorem 4.1.1] (9)

$$k_a \ge a - g + 1 \quad \forall \ 0 \le a < m,$$

and equality occurs for $a \ge 2g - 1$ (10)
[31, Remark 4.1.4]

$$k_a^{\perp} + d_a^{\perp} \ge m - g + 1 \qquad \forall \ 0 \le a < m$$
[31, Exercise 4.1.27] (11)

Naively one might imagine that $C_a \subsetneq C_{a+1}$ for all a. However that is not necessarily the case since (10) does not exactly specify the dimension: for small a it only provides a lower bound. However, this will not be problematic for our purposes. We will only consider C_a for $a \ge 2g - 1$, in which case (10) determines the dimension exactly.

Clearly, for a curve of genus 0, each code C_a must be an MDS code, since adding (9) and (10) shows that the Singleton bound is tight. More generally, the AG code construction is most efficient when it is based on a curve for which g/N is as small as possible. However this ratio cannot be arbitrarily small. It is known⁶ that $g/N \ge c/\sqrt{q}$ for some constant c > 0and sufficiently large g. In fact, this bound is asymptotically tight. Various explicit curves

⁶This follows from the Drinfeld-Vlăduț theorem, but is weaker since we do not need exact constants.

asymptotically matching this bound have been discovered by Tsfasman, Vlăduţ and Zink [33], Garcia and Stichtenoth [8], and others.

Using such curves, and letting $m = \Omega(N)$, one may see from (9) and (10) that the Singleton bound nearly holds, but there is a "defect" of $g = O(m/\sqrt{q})$. The rough idea of our analysis is to choose parameters such that $\sqrt{q} = \Theta(m/t)$, so that the defect is only O(t), which will be acceptable for our purposes.

There is a slight complication. The use of algebraic curves in coding theory primarily concerns limiting behaviour of the rate/distance tradeoff as the code length m tends to infinity. Since we wish to construct orthogonal arrays for *all* parameters m, n, t, we must ensure that curves exist with values of N that are sufficiently dense. It is not necessarily true that all curve families that have previously been used with AG codes will be suitable for our purposes. Fortunately, the modular curves discussed in the following theorem are suitably dense.

Theorem 12 ([17], or [32, page 18]). Let $p \neq \ell$ be distinct primes with $p \neq 11$. Then the classical modular curve $X_0(11\ell)$ over \mathbb{F}_{p^2} has at least $N = (p-1)(\ell+1)$ rational points and has genus $g = \ell$. Furthermore, the generator matrix of the codes made from such curves can be constructed in time poly(m).

We remark that the celebrated work of [33] also used the classical modular curve $X_0(\cdot)$ over \mathbb{F}_{p^2} , but they did not require the genus to be a multiple of 11.

Using AG codes for orthogonal arrays. Next we explain how the AG codes can be used to construct orthogonal arrays. Let p be the smallest prime satisfying

$$p \equiv 1 \pmod{n}$$
 and $p \ge 11 + m/t$.

Clearly $p \neq 11$. By Linnik's theorem (Corollary 8), $p \leq c_L(n+11+m/t)^5$. Let $q = p^2$ and note that $q \equiv 1 \pmod{n}$.

We choose ℓ to be the smallest prime such that $\ell \geq t$; by Bertrand's postulate $\ell \leq 2t$. By Theorem 12, the classical modular curve $X_0(11\ell)$ over \mathbb{F}_q satisfies

$$N = (p-1)(\ell+1) > m$$
 and $g = \ell \le 2t.$ (12)

Given this curve, Theorem 11 guarantees existence of a sequence of codes with various parameters. We will only use the codes C_u and C_{u+1} , where

$$u = 2\ell - 1 + t = 2g - 1 + t. \tag{13}$$

The following claim performs some simple calculations in preparation for using Corollary 5.

Claim 13.

- 1. $C_u \subsetneq C_{u+1}$. 2. $d_u^{\perp} \ge t+1$.
- 3. $k_u \leq 3t$.
- 4. $d_{u+1} \ge m 5t$.

Proof.

- 1. By (13) we have u > 2g 1, so (10) holds with equality for $a \in \{u, u + 1\}$, and therefore $k_u < k_{u+1}$.
- 2. $d_u^{\perp} \ge m k_u^{\perp} g + 1 = k_u g + 1 \ge u 2g + 2 = t + 1$, by (11), (10), and (13).
- 3. By (13) we have u > 2g 1, so (10) holds with equality for a = u. Thus

$$k_u = u - g + 1 = (2g - 1 + t) - g + 1 \le 3t,$$

by (10), (13), and (12).

4.
$$d_{u+1} \ge m - u - 1 = m - (2g - 1 + t) - 1 \ge m - 5t$$
, by (9), (13), and (12).

We apply Corollary 5 to C_u and C_{u+1} with $k \leq 3t$ and $d_2 \geq m - 5t$. Thus, there is an [s, m, n, t] orthogonal array with

$$s \leq n^{5t}q^{3t} \leq n^{5t} (c(n+11+m/t)^5)^{6t}.$$

Regarding the algorithmic efficiency, Theorem 12 above states that the generator matrix for C_u and C_{u+1} can be constructed in time poly(m). Given those matrices, a vector $v \in C_{u+1} \setminus C_u$ (as required by Corollary 5) can be found in poly(sm) time. Since Algorithm 1 also runs in poly(sm), it follows that the construction of Corollary 5 runs in poly(sm). This concludes the proof of Theorem 10.

5 Implementing a *t*-independent hash function

Although orthogonal arrays and t-independent hash functions are mathematically equivalent (see Claim 2), a key difference is the efficiency of their implementations. A hash function needs to be represented in little space, and the algorithms for constructing and evaluating it should be fast.

In Algorithm 2, we present pseudocode for a *t*-independent hash function based on the orthogonal array construction of Sections 2 and 3. We assume that UNIFORM(S) returns a uniformly random element of the finite set S in O(1) time.

Theorem 14. In the pseudocode of Algorithm 2:

- CONSTRUCTOR has expected runtime $O(t + (n+m)^{\epsilon})$ for every $\epsilon > 0$.
- EVAL has runtime O(t).
- The space for a HASH object is $O(t \log(n+m))$ bits.

Assuming the generalized Riemann hypothesis, the time for CONSTRUCTOR is $O(t + \log(n+m))$.

The proof involves the following notation, for $x \in \mathbb{R}$, $\eta, a \in \mathbb{Z}$, $\eta \ge 2$,

$$S_{x,\eta,a} = \{ i \in \mathbb{N} : i \equiv a \pmod{\eta}, 1 < i \leq x \}$$

$$\pi(x;\eta,a) = |\{ p \in S_{x,\eta,a} : p \text{ is prime } \}|$$

as well as the Euler totient function $\phi(\eta)$, which counts the number of positive integers up to η that are relatively prime to η . Throughout this section, we assume that a and η are coprime.

Algorithm 2 Pseudocode for a *t*-independent hash function. This pseudocode involves the constant ν (from Theorem 15), which is effectively computable (see Remark 18).

class HASH:

dictionary D $\triangleright D$ has size at most t, so it can be implemented as an array int n, m, t, pint a[0..t-1]

CONSTRUCTOR(int n, m, t)

Store parameters n, m, t in the object Let $\eta \leftarrow n \cdot \lceil m/n \rceil$ Find a prime $p \in [\eta + 1, \eta^{\nu}]$ such that $p \equiv 1 \pmod{\eta}$ \triangleright Generate coefficients of a random polynomial of degree-(t-1)for $i \leftarrow 0 \dots t-1$ $\mid a_i \leftarrow \text{UNIFORM}(\mathbb{F}_p)$ $D \leftarrow \text{empty dictionary}$

 \triangleright Evaluate the hash function at input $x \in [m]$

```
EVAL(int x)
```

```
 \begin{array}{l} \mathbf{if} \ x \in \operatorname{keys}(D) \\ | & \rhd \text{ The hash value of } x \text{ is already in the dictionary} \\ \mathbf{return} \ D[x] \\ \vartriangleright \ \operatorname{Evaluate the chosen polynomial at } x \ (\operatorname{over} \ \mathbb{F}_p) \\ \operatorname{Let} \ y \leftarrow \sum_{i=0}^{t-1} a_i x^i \\ \mathbf{if} \ y = x^t \quad \rhd \text{ Here } x^t \text{ is the "bad value" } b(x) \\ | & \rhd \text{ Choose a replacement value randomly} \\ \operatorname{Let} \ U \leftarrow \text{UNIFORM}([n]) \\ & \rhd \text{ Store the hash value of } x \text{ in the dictionary} \\ D[x] \leftarrow U \\ \mathbf{return} \ U \\ & \rhd \text{ Return } \phi_{b(x)}(y) \\ \mathbf{return } 1 + (((y + p - 1 - x^t) \mod p) \mod n) \end{array}
```

Proof. It is not difficult to see that this pseudocode implements a t-independent hash function, since it implements the construction of Sections 2 and 3. We now summarize the main ideas.

The key task of the CONSTRUCTOR is to find a prime p satisfying $p \equiv 1 \pmod{n}$ and $\eta + 1 \leq p \leq \eta^{\nu}$, for some constant ν defined in Theorem 15. The runtime of that step is discussed extensively below. It then randomly generates the coefficients of a uniformly random degree-(t-1) polynomial, which is analogous to uniformly selecting the function $h \in \mathcal{H}_q$ in the proof of Lemma 4. (Here q = p.) The "bad vector" b is defined to be the evaluations of the degree-t monomial, i.e., $b(x) = x^t$. This is a codeword in C_t but not in C_{t-1} , and so it can be used as in the proof of Theorem 6. The EVAL function calculates h(x), and checks whether it equals the forbidden value b(x). If so, it replaces it with a random value and caches that in the

dictionary D. Otherwise it returns $\phi_{b(x)}(h(x))$.

We claim that there are at most t inputs $x \in [m]$ for which $\sum_{i=0}^{t-1} a_i x^i = x^t$ (i.e., h(x) = b(x)). This follows since the polynomial h(x) - b(x) is of degree t and hence has at most t roots. This implies that the dictionary D will have size at most t. Each key in the dictionary is in [m] and each value is in [n], so the space required for D is $O(t \log(n+m))$ bits. Since $p \leq (n+m)^{\nu}$, and each $a_i \in \mathbb{F}_p$, the space required for all other parameters of the HASH object is $O(t \log(n+m))$ bits.

EVAL can clearly be implemented in O(t) time, even if the dictionary D is implemented as an unsorted array.

The main challenge of the analysis is the time required to find the prime p. In Section 3 the time to find the prime p was negligible compared to the time to build the orthogonal array, so a (deterministic) algorithm with runtime $O((n+m)^4)$ was sufficient. To obtain an improved runtime for CONSTRUCTOR, we will use random sampling. Theorem 15 below shows that there exists a constant ν such that, assuming $x \geq \eta^{\nu}$, we have

$$\pi(x;\eta,a) \geq \frac{c_{\epsilon}x}{\phi(\eta)\ln(x)\eta^{\epsilon}} \geq \frac{c_{\epsilon}x}{\eta^{1+\epsilon}\ln x},$$

using the trivial bound $\phi(\eta) \leq \eta$.

Let us now fix $x = \eta^{\nu}$ and a = 1. The algorithm repeatedly samples integers at random from the set $S_{x,\eta,a}$, testing each for primality. The expected number of iterations until finding a prime is

$$\frac{|S_{x,\eta,a}|}{\pi(x;\eta,a)} \leq \frac{x/\eta}{c_{\epsilon}x/(\eta^{1+\epsilon}\ln x)} = \frac{\eta^{\epsilon}\ln x}{c_{\epsilon}},\tag{14}$$

since $x \ge \eta$. Each primality test requires O(polylog(x)) time. Using that $\eta \le (n+m)$, we have shown that the expected time to find the prime p is $O((n+m)^{\epsilon})$ for every $\epsilon > 0$.

Assuming the generalized Riemann hypothesis, Corollary 16 yields

$$\pi(x;\eta,a) \geq \frac{x}{2\eta\ln x},$$

so the expected number of iterations until finding the prime p is

$$\frac{|S_{x,\eta,a}|}{\pi(x;\eta,a)} \le \frac{2x/\eta}{x/\eta \ln x} = 2\ln(x),$$

which is $O(\log(n+m))$ since $x = \eta^{\nu}$.

The next theorem follows from known results in the analytic number theory literature, as we explain below.

Theorem 15 (Quantitative Linnik's Theorem). There exists a universal constant $\nu \geq 1$ (independent of η, ε) and a constant $c_{\varepsilon} > 0$ such that, for any $\varepsilon > 0$, any $x \geq \eta^{\nu}$,

$$\pi(x;\eta,a) \geq c_{\varepsilon}\left(\frac{x}{\phi(\eta)\ln x}\cdot\frac{1}{\eta^{\varepsilon}}\right).$$

The distribution of the primes in an arithmetic progression is closely tied to the wellstudied *L*-functions of characters mod η . As with many results in analytic number theory, there are different cases depending on whether the *L*-functions have so-called "exceptional zeros". Whether or not these zeros exist is unknown, and relates to the generalized Riemann hypothesis. We will use the following result, which is a corollary of [14, Theorems 18.6, 18.7].

Corollary 16. There exist universal constants ν , c_1 , c_2 , c_3 (independent of x, η , a) such that, if $x \ge \eta^{\nu}$ and $\eta \ge c_3$, then the following holds.

• If there exists a real character $\chi \pmod{\eta}$ whose L-function has a real zero β satisfying $1 - \beta \leq c_1/2 \log \eta$ then

$$\pi(x;\eta,a) \geq c_2 \cdot \frac{x}{\phi(\eta)} \cdot (1-\beta).$$

• If there exists no such character χ , then

$$\pi(x;\eta,a) \geq \frac{1}{2} \cdot \frac{x}{\phi(\eta) \ln x}$$

The generalized Riemann hypothesis implies that the second case holds.

The results in [14] are stated in terms of the von Mangoldt function Λ and its sum ψ :

$$\Lambda(n) = \begin{cases} \ln p & \text{if } n = p^k \text{ for a prime } p \text{ and integer } k \ge 1 \\ 0 & \text{otherwise.} \end{cases}$$
$$\psi(x; \eta, a) = \sum_{\substack{n \le x \\ n \equiv a \pmod{\eta}}} \Lambda(n).$$

Corollary 16 follows from those results using the simple inequality $\psi(x; \eta, a) \leq \pi(x; \eta, a) \cdot \ln(x)$.

Let us now return to the proof of Theorem 15. If the second case of Corollary 16 applies, then Theorem 15 is immediate. If the first case of Corollary 16 applies, then we must ensure that the exceptional zero is significantly less than 1. The following is a classic result of Siegel [26] [22, Corollary 11.15].

Theorem 17 (Siegel, 1935). For every $\epsilon > 0$ there exists a constant c_{ϵ} such that, if χ is a real character (mod η) with $L(s, \chi)$ having a real zero β then

$$\beta < 1 - c_{\epsilon} \eta^{-\epsilon}.$$

Thus, in the first case of Corollary 16, it holds that $1 - \beta \ge c_{\epsilon} \eta^{-\epsilon}$, which completes the proof of Theorem 15.

Remark 18. The numerical value of constants ν , c_L , L, c_1 , c_2 , c_3 satisfying the above properties can be explicitly computed "given the time and will"; see [14, Pages 427, 428 and property 18.90]. Such constants are called "effectively computable" in the number theory literature. In contrast, the value of c_{ϵ} is "ineffective" for $\epsilon < \frac{1}{2}$ [14, Theorem 5.28], meaning that the proof of the result does not yield a way to compute the constant. However, note that implementing Algorithms 1 and 2 only requires knowing the value of the constants ν , c_L , L. The constant c_{ϵ} only appears in the analysis of Theorem 14. Specifically, from (14), it appears inside the O(.)bound on the runtime of CONSTRUCTOR.

6 Construction using random linear codes

Sections 3 and 4 use deterministic code constructions. In this section we show that a random construction of linear codes can also be used to prove an non-explicit form of Theorem 1, using an efficient, randomized algorithm.

Lemma 19. There exists a universal constant c > 0 such that the following is true. For any $m, n, t \in \mathbb{N}$, there exists a prime number p, a linear code $C \subseteq \mathbb{F}_p^m$ and a function $b: [m] \to \mathbb{F}_p$ such that

- $p \equiv 1 \pmod{n}$ and $p \leq c_L (n + (me/t)^3)^L$.
- the dimension of C is at most 2t.
- the distance of C^{\perp} is at least t + 1.
- $\Delta(b, v) \ge m 3t \ \forall v \in C.$

Moreover, there exists a randomized algorithm that outputs p, C and b satisfying these conditions and has expected runtime $(mn/t)^{O(t)}$.

The code C and vector b are then provided as input to Algorithm 1. Applying Lemma 4, we obtain an [s, m, n, t] orthogonal array with $s \leq n^{3t}p^{2t} = (mn/t)^{O(t)}$. The expected runtime is $(mn/t)^{O(t)}$.

The proof of Lemma 19 requires the following lemma. Here, the matrix M is simply the generator matrix for the dual of a code meeting the Gilbert–Varshamov bound. A non-algorithmic form of this lemma may be found in [30, Theorem 10.11].

Lemma 20. Let $m, t \in \mathbb{N}$ and let p be a prime power. Assume that

$$\sum_{i=1}^{t} \binom{m}{i} (p-1)^{i} \leq p^{\ell}/4.$$
(15)

Then there exists a randomized algorithm that, with probability at least 3/4, outputs an $\ell \times m$ matrix M such that any t columns of M are linearly independent.

The proof is classical, but for completeness we include it below.

Proof (of Lemma 19). Define

$$\ell = 2t$$
 and $s = 3t$.

Let p be a prime satisfying both

$$p \equiv 1 \pmod{n}$$
 and $(me/t)^3 < p_1$

By Corollary 8, there exists such a p with

$$p \leq c_L (n + (me/t)^3)^L.$$
 (16)

We can find p deterministically with runtime $(cnm/t)^c$ for some constant c.

Next we claim that (15) is satisfied. Using $t \leq (m-1)/2$ and p > 2, the LHS is at most $t\binom{m}{t}p^t$. Dividing both sides by p^{ℓ} , we get

$$t\binom{m}{t} p^{t-\ell} \leq t(me/t)^t p^{-t} \quad (\text{using } \ell = 2t)$$

$$< t(me/t)^{-2t} \quad (\text{using the lower bound on } p)$$

$$< te^{-4t} \leq 1/4e,$$

using that $m \ge e \cdot t$. Since this is less than 1/4, (15) is satisfied.

We will repeatedly use Lemma 20 to generate a matrix M until all subsets of t columns are linearly independent. This takes time $O(\binom{m}{t}(mt)^c)$ for some constant c. Let C be the list of vectors in \mathbb{F}_p^m obtained by taking all linear combinations of the rows of M. Then C is a linear code of dimension at most ℓ and dual distance at least t + 1; see, e.g., [30, Theorem 10.4]. The time to construct C is $O(|C|m\ell)$.

Lastly, we will repeatedly generate $b \in \mathbb{F}_p^m$ uniformly at random until $\Delta(b, v) \geq m-s \ \forall v \in C$. For any $I \in \binom{[m]}{s}$, let E_I be the event that there exists v for which $v_i = b_i \ \forall i \in I$. We will show that $\Pr[\bigcup_I E_I] < 1$. Clearly

$$\Pr\left[\mathcal{E}_{I}\right] \leq |C| \cdot p^{-s} \leq p^{\ell-s} = p^{-t}.$$

Thus, by a union bound,

$$\Pr\left[\cup_{I} E_{I}\right] \leq \binom{m}{s} p^{-t}$$
$$\leq (me/s)^{s} p^{-t}$$
$$= 3^{-3t} (me/t)^{3t} p^{-t}$$
$$< 3^{-3t} < 1,$$

by the definition s = 3t and the lower bound $p > (me/t)^3$. Thus, the expected number of trials until generating b is a constant. Each trial can be executed in time |C|m. Overall, the expected runtime is $(mn/t)^{O(t)}$.

Proof (of Lemma 20). Let M be a uniformly random matrix of size $\ell \times m$. Let $V \subseteq \mathbb{F}_p^m$ be the non-zero vectors of support size at most t. For $v \in V$, let E_v be the event that Mv = 0. Then $\Pr[E_v] = p^{-\ell}$, so

$$\Pr\left[\bigcup_{v\in V} E_v\right] \leq |V|p^{-\ell} = \sum_{i=1}^t \binom{m}{i} (p-1)^i p^{-\ell}.$$

By the lemma's hypothesis, this is at most 1/4. If this event does not occur then every linear dependence among the columns of M involves more than t columns.

7 Open Questions

There are several open questions related to this work.

- 1. The orthogonal array constructed by Theorem 1 will have duplicate rows, in general. (Similarly, the hash family \mathcal{H} constructed by Theorem 3 will, in general, be a multiset.) Can it be modified to eliminate the duplicate rows?
- 2. In Theorem 14, the expected runtime of CONSTRUCTOR is $O(t + (n+m)^{\epsilon}) \forall \epsilon > 0$. Can it be improved to O(t) + polylog(n+m), without assuming the generalized Riemann hypothesis?
- 3. The hash function construction of Section 5 was based on the Reed-Solomon construction of Section 3. Can the construction of Section 4 be used instead? Although there are algorithms to construct the generator matrix in time poly(m) (see Theorem 12), we would like to evaluate the hash function in time poly(t) while using only space $O(t \log(n+m))$. At present we are unaware of an AG code construction that can be used for this purpose.
- 4. Although there are explicitly known values of L for which Linnik's theorem (Theorem 7) holds, at present it seems that there is no explicitly known pair (L, c_L) such that the theorem holds.

Acknowledgements

We thank Jan Vondrak for suggesting this problem to us and for some initial discussions. We thank Lior Silberman and Greg Martin for very helpful discussions on analytic number theory.

References

- K. A. Bush. A generalization of a theorem due to MacNeish. The Annals of Mathematical Statistics, pages 293–295, 1952.
- [2] K. A. Bush. Orthogonal arrays of index unity. Ann. Math. Statist., 23:426–434, 1952.
- [3] J. Lawrence Carter and Mark N. Wegman. Universal classes of hash functions. Journal of Computer and System Sciences, 18:143–154, 1979.
- [4] Kuan Cheng and Xin Li. Efficient document exchange and error correcting codes with asymmetric information. In Proceedings of the ACM-SIAM Symposium on Discrete Algorithms (SODA), 2021.
- [5] Benny Chor, Oded Goldreich, Johan Hastad, Joel Friedman, Steven Rudich, and Roman Smolensky. The bit extraction problem or t-resilient functions. In Proceedings of the IEEE Symposium of Foundations of Computer Science, 1985.
- [6] Jeffrey S Cohen and Daniel M Kane. Bounds on the independence required for cuckoo hashing, 2009. Manuscript.
- [7] Shahar Dobzinski, Sigal Oren, and Jan Vondrak. Fairness and incentive compatibility via percentage fees. In Proceedings of the 24th ACM Conference on Economics and Computation, pages 517–535, 2023.

- [8] Arnaldo Garcia and Henning Stichtenoth. A tower of Artin-Schreier extensions of function fields attaining the Drinfeld-Vladut bound. *Inventiones mathematicae*, 121(1):211–222, 1995.
- [9] Valerii Denisovich Goppa. Algebraico-geometric codes. *Izvestiya Rossiiskoi Akademii* Nauk. Seriya Matematicheskaya, 46(4):762–781, 1982.
- [10] Venkat Guruswami, Atri Rudra, and Madhu Sudan. Essential Coding Theory. 2018. Manuscript.
- [11] D. R. Heath-Brown. Zero-free regions for Dirichlet L-functions, and the least prime in an arithmetic progression. *Proceedings of the London Mathematical Society*, 3(64):265–338, 1992.
- [12] A. S. Hedayat, N. J. A. Sloane, and John Stufken. Orthogonal Arrays: Theory and Applications. Springer, 1999.
- [13] Piotr Indyk. A small approximately min-wise independent family of hash functions. *Journal of Algorithms*, 38(1):84–90, 2001.
- [14] Henryk Iwaniec and Emmanuel Kowalski. Analytic number theory. American Mathematical Society, 2004.
- [15] Daniel M Kane, Jelani Nelson, and David P Woodruff. An optimal algorithm for the distinct elements problem. In Proceedings of the 29th Symposium on Principles of Database Systems (PODS), pages 41–52, 2010.
- [16] Richard M. Karp and Avi Wigderson. A fast parallel algorithm for the maximal independent set problem. *Journal of the ACM (JACM)*, 32(4):762–773, 1985.
- [17] G. Katsman, M. Tsfasman, and S. Vlăduţ. Modular curves and codes with a polynomial construction. *IEEE Transactions on Information Theory*, 30(2):353–355, 1984.
- [18] Greg Kuperberg, Shachar Lovett, and Ron Peled. Probabilistic existence of rigid combinatorial structures. In *Proceedings of the forty-fourth annual ACM symposium on Theory* of computing, pages 1091–1106, 2012.
- [19] Greg Kuperberg, Shachar Lovett, and Ron Peled. Probabilistic existence of regular combinatorial structures. *Geometric and Functional Analysis*, 27(4):919–972, 2017.
- [20] Youness Lamzouri, Xiannan Li, and Kannan Soundararajan. Conditional bounds for the least quadratic non-residue and related problems. *Mathematics of Computation*, 84(295):2391–2412, 2015.
- [21] Dino Lorenzini. An Invitation to Arithmetic Geometry. American Mathematical Society, 1996.
- [22] Hugh L. Montgomery and Robert C. Vaughan. *Multiplicative Number Theory I*, chapter 11. Cambridge University Press, 2006.

- [23] Harald Niederreiter and Chaoping Xing. Rational Points on Curves over Finite Fields: Theory and Applications. Cambridge University Press, 2001.
- [24] C. R. Rao. Factorial experiments derivable from combinatorial arrangements of arrays. Supplement to the Journal of the Royal Statistical Society, 9(1):128–139, 1947.
- [25] Ron Roth. Introduction to Coding Theory. Cambridge University Press, 2006.
- [26] Carl Siegel. Uber die classenzahl quadratischer zahlkörper. Acta Arithmetica, (1):83–86, 1935.
- [27] D. R. Stinson. Combinatorial techniques for universal hashing. Journal of Computer and System Sciences, 48:337–346, 1994.
- [28] D. R. Stinson. On the connections between universal hashing, combinatorial designs and error-correcting codes. *Congressus Numerantium*, pages 7–28, 1996.
- [29] D. R. Stinson. Universal hash families and the leftover hash lemma, and applications to cryptography and computing. J. Combin. Math. Combin. Comput., 42:3–31, 2002.
- [30] D. R. Stinson. Combinatorial Designs: Constructions and Analysis. Springer, 2004.
- [31] Michael Tsfasman, Serge Vlăduţ, and Dmitry Nogin. Algebraic Geometric Codes: Basic Notions. American Mathematical Society, 2007.
- [32] Michael Tsfasman, Serge Vlăduţ, and Dmitry Nogin. Algebraic Geometric Codes: Advanced Chapters. American Mathematical Society, 2019.
- [33] Michael A. Tsfasman, S. G. Vlăduţ, and Th. Zink. Modular curves, Shimura curves, and Goppa codes, better than Varshamov-Gilbert bound. *Mathematische Nachrichten*, 109(1):21–28, 1982.
- [34] Mark N. Wegman and J. Larence Carter. New hash functions and their use in authentication and set equality. *Journal of Computer and System Sciences*, 22:265–279, 1981.
- [35] Triantafyllos Xylouris. On Linnik's constant. Acta Arithmetica, 150(1):65–91, 2011.
- [36] Triantafyllos Xylouris. Uber die Nullstellen der Dirichletschen L-Funktionen und die kleinste Primzahl in einer arithmetischen Progression. PhD thesis, 2011.