

CONFORMAL METRICS WITH FINITE TOTAL Q-CURVATURE REVISITED

MINGXIANG LI

ABSTRACT. Given a conformal metric with finite total Q-curvature on \mathbb{R}^n for $n \geq 4$, we show that the sign of scalar curvature near infinity control not only the upper bound but also the lower bound of Q-curvature integral which is a new phenomenon. Meanwhile, for general complete non-compact four-manifolds with simple ends, we also obtain similar control of the lower bound of Q-curvature integral.

1. INTRODUCTION

For compact surfaces, the celebrated Chern-Gauss-Bonnet formula [7] intricately links the surface's geometry to its topology. Analogously, for complete open surfaces, this connection is elucidated by the works of Cohn-Vossen [8], Huber [12], and Finn [10]. We also suggest interested readers referring to [17] and [23] for additional insights. When it comes to higher-dimensional manifolds, is there a similar connection to explore? This is the Problem 11 in chapter 7 of [27] established by Schoen and Yau. Some progress has been made by Poor [26] and Greene-Wu [11] under the assumption of non-negative sectional curvature.

In higher-dimensional manifolds, Q-curvature which is firstly introduced by Branson [2] is a natural generalization of Gaussian curvature. In particular, for a four dimensional manifold (M, g) , Q-curvature is defined by

$$Q_g := -\frac{1}{6}(\Delta_g R_g - R_g^2 + 3|Ric_g|^2)$$

where Δ_g is the Laplace-Beltrami operator, Ric_g and R_g are the Ricci curvature and scalar curvature of (M, g) respectively. Combing with the Paneitz operator [25], for a conformal metric $\tilde{g} = e^{2u}g$, there holds a conformally invariant equation

$$P_g u + Q_g = Q_{\tilde{g}} e^{4u}$$

where the Paneitz operator P_g is defined by

$$P_g = \Delta_g^2 - \operatorname{div}_g \left(\left(\frac{2}{3} R_g g - 2 Ric_g \right) d \right).$$

With the assistance of Q-curvature, Chang, Qing, and Yang [3] established an analogous Cohn-Vossen inequality for a complete manifold $(\mathbb{R}^4, e^{2u}|dx|^2)$.

2020 *Mathematics Subject Classification.* Primary: 53C18, Secondary: 58J90.

Key words and phrases. Q-curvature, Cohn-Vossen inequality, Volume comparison.

Fang [9] and Ndiaye-Xiao [24] further extended this to higher dimensional cases.

Precisely, for a complete conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer, the Q-curvature satisfies the conformally invariant equation:

$$(1.1) \quad (-\Delta)^{\frac{n}{2}} u = Qe^{nu}.$$

More information about the equation (1.1) can be found in [1], [4], [6], [13], [14], [15], [16], [20], [22], [29], [31], [32] and the references therein. For brevity, we consider the smooth metrics throughout this paper.

We say the metric g has finite total Q-curvature if

$$\int_{\mathbb{R}^n} |Q|e^{nu} dx < +\infty$$

and "near infinity" means outside a compact set.

Theorem 1.1. (See [3], [9], [24]) *Given a complete conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q-curvature. Suppose that the scalar curvature $R_g \geq 0$ near infinity. Then there holds*

$$\int_{\mathbb{R}^n} Qe^{nu} dx \leq \frac{(n-1)!|\mathbb{S}^n|}{2}$$

where $|\mathbb{S}^n|$ denotes the volume of standard n -sphere.

Our first result is to show that the integral of Q-curvature under the same assumption of Theorem 1.1 is non-negative.

Theorem 1.2. *Given a complete conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q-curvature. Suppose that the scalar curvature $R_g \geq 0$ near infinity. Then there holds*

$$0 \leq \int_{\mathbb{R}^n} Qe^{nu} dx \leq \frac{(n-1)!|\mathbb{S}^n|}{2}$$

where $|\mathbb{S}^n|$ denotes the volume of standard n -sphere.

Moreover, should the scalar curvature exhibit a strictly positive lower bound near infinity, the inequality in Theorem 1.1 attains equality.

Theorem 1.3. *Given a complete conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q-curvature. Suppose that the scalar curvature $R_g \geq C$ near infinity for some constant $C > 0$. Then there holds*

$$\int_{\mathbb{R}^n} Qe^{nu} dx = \frac{(n-1)!|\mathbb{S}^n|}{2}.$$

Drawing on the aforementioned findings and the results in [4], we can effectively control the integral of the Q-curvature over four-manifolds with simple ends. A detailed definition of simple ends can be found in Section 4.

Theorem 1.4. *Given a complete four dimensional manifold (M, g) with only finitely many conformally flat simple ends and finite total Q -curvature. Suppose that the scalar curvature is non-negative at each end. Then there holds*

$$\chi(M) - \#\{\text{ends}\} \leq \frac{1}{32\pi^2} \int_M (|W|^2 + 4Q_g) d\mu_g \leq \chi(M)$$

where $\chi(M)$, W and $\#\{\text{ends}\}$ denote the Euler number, Weyl tensor and the number of ends respectively.

Theorem 1.5. *Given a complete four dimensional manifold (M, g) with only finitely many conformally flat simple ends and finite total Q -curvature. Suppose that the scalar curvature $R_g \geq C$ for some constant $C > 0$ at each end. Then there holds*

$$\int_M (|W|^2 + 4Q_g) d\mu_g = 32\pi^2 \chi(M)$$

where $\chi(M)$ and W denote the Euler number and Weyl tensor respectively.

In fact, even without assuming completeness, we still have control over the integral of Q -curvature.

Theorem 1.6. *Given a conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q -curvature.*

(1) *If the scalar curvature $R_g \geq 0$ near infinity, there holds*

$$0 \leq \int_{\mathbb{R}^n} Qe^{nu} dx \leq (n-1)!|\mathbb{S}^n|.$$

(2) *If the scalar curvature $R_g \geq C > 0$ near infinity, there holds*

$$\frac{(n-1)!|\mathbb{S}^n|}{2} \leq \int_{\mathbb{R}^n} Qe^{nu} dx \leq (n-1)!|\mathbb{S}^n|.$$

(3) *If the scalar curvature $0 \leq R_g \leq C < +\infty$ near infinity, there holds*

$$\int_{\mathbb{R}^n} Qe^{nu} dx = (n-1)!|\mathbb{S}^n| \quad \text{or} \quad 0 \leq \int_{\mathbb{R}^n} Qe^{nu} dx \leq \frac{(n-1)!|\mathbb{S}^n|}{2}.$$

Making use of Theorem 1.6 and Theorem 1.1, we finish the proofs of Theorem 1.2 and Theorem 1.3.

Meanwhile, with help of Theorem 1.6 and the classification theorem for $Q = (n-1)!$ in (1.1) established in [5], [20], [22], [31], and [32], we are able to confirm the Conjecture 2 in [16].

Corollary 1.7. *Given a conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q -curvature. Suppose that the scalar curvature $R_g \geq 0$ near infinity and the Q -curvature*

$$Q_g \geq (n-1)!,$$

then there holds

$$\int_{\mathbb{R}^n} e^{nu} dx \leq |\mathbb{S}^n|$$

with equality holds if and only if $u(x) = \log \frac{2\lambda}{\lambda^2 + |x - x_0|^2}$ for some constant $\lambda > 0$ and $x_0 \in \mathbb{R}^n$.

When imposing a constraint on the integral of scalar curvature, we also obtain similar control of the Q -curvature integral.

Theorem 1.8. *Given a conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q -curvature.*

(1) *Suppose that the negative part of scalar curvature R_g^- satisfies*

$$(1.2) \quad \int_{\mathbb{R}^n} (R_g^-)^{\frac{n}{2}} e^{nu} dx < +\infty.$$

Then there holds

$$0 \leq \int_{\mathbb{R}^n} Q e^{nu} dx \leq (n-1)! |\mathbb{S}^n|.$$

(2) *Suppose that scalar curvature R_g satisfies*

$$(1.3) \quad \int_{\mathbb{R}^n} |R_g|^{\frac{n}{2}} e^{nu} dx < +\infty.$$

Then there holds

$$\int_{\mathbb{R}^n} Q e^{nu} dx = 0 \quad \text{or} \quad \int_{\mathbb{R}^n} Q e^{nu} dx = (n-1)! |\mathbb{S}^n|.$$

Moreover, if the volume is finite i.e. $e^{nu} \in L^1(\mathbb{R}^n)$ in addition, then there holds

$$\int_{\mathbb{R}^n} Q e^{nu} dx = (n-1)! |\mathbb{S}^n|.$$

Remark 1.9. Part (1) in Theorem 1.8 is a generalization of Part (1) in Theorem 1.6. Some results similar to Part (2) of Theorem 1.8 are also obtained by Lu-Wang [21], Wang-Zhou [30], and Li-Wang [18] with help of different methods.

With help of Theorem 1.8 and the classification theorem for $Q = (n-1)!$ in (1.1), we are able to partially confirm the Conjecture 1 in [16].

Corollary 1.10. *Given a conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q -curvature. Suppose that the scalar curvature R_g and Q -curvature Q_g satisfy*

$$\int_{\mathbb{R}^n} |R_g|^{\frac{n}{2}} e^{nu} dx < +\infty, \quad Q_g \leq (n-1)!.$$

Then there holds

$$\int_{\mathbb{R}^n} e^{nu} dx \geq |\mathbb{S}^n|$$

with equality holds if and only if $u(x) = \log \frac{2\lambda}{\lambda^2 + |x - x_0|^2}$ for some constant $\lambda > 0$ and $x_0 \in \mathbb{R}^n$.

Now, let's briefly outline the structure of this paper. In Section 2, we establish some crucial estimates for normal solutions. Leveraging these estimates, we proceed to prove Theorem 1.6 and Theorem 1.8 in Section 3. Finally, in Section 4, we address general manifolds with finitely many simple ends, demonstrating Theorem 1.4 and Theorem 1.5.

Acknowledgment. The author would like to thank Professor Xingwang Xu for helpful discussions.

2. NORMAL SOLUTIONS ON \mathbb{R}^n

We say the solution $u(x)$ to (1.1) with $Qe^{nu} \in L^1(\mathbb{R}^n)$ is normal if $u(x)$ satisfies the integral equation

$$(2.1) \quad u(x) = \frac{2}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} \log \frac{|y|}{|x-y|} Q(y) e^{nu(y)} dy + C$$

where $|\mathbb{S}^n|$ denotes the volume of standard sphere \mathbb{S}^n and C is a constant which may be different from line to line throughout this paper. In particular, the metric $g = e^{2u}|dx|^2$ with finite total Q-curvature said to be normal if $u(x)$ is normal. More details about normal solutions, one may refer to Section 2 of [14].

In order to make the metric g to be normal, Chang, Qing and Yang [3] give a geometric criterion about scalar curvature.

Lemma 2.1. (*Theorem 1.4 in [3], Theorem 1.1 in [28]*) *Given a smooth conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q-curvature. If the scalar curvature $R_g \geq 0$ near infinity, then the metric g is normal.*

S. Wang and Y. Wang [28] established a criterion under the control of the integral of negative part of scalar curvature.

Lemma 2.2. (*Theorem 1.1 in [28], Theorem 4.2 in [14]*) *Given a smooth conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n where $n \geq 4$ is an even integer with finite total Q-curvature. If the negative part of scalar curvature R_g^- satisfies*

$$\int_{\mathbb{R}^n} (R_g^-)^{\frac{n}{2}} e^{nu} dx < +\infty,$$

then the metric g is normal.

Recently, the author [14] introduced a volume entropy defined by

$$\tau(g) := \lim_{R \rightarrow \infty} \sup \frac{\log V_g(B_R(0))}{\log |B_R(0)|}$$

where $B_R(0)$ denotes the Euclidean ball with radius R centered at the origin and $V_g(B_R(0))$, $|B_R(0)|$ represent the volume of $B_R(0)$ related to the metric g and Euclidean metric respectively. With help of such volume entropy $\tau(g)$, a necessary and sufficient condition of the normal metric is established.

Theorem 2.3. (Theorem 1.1 in [14]) *Given a complete metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n with finite total Q -curvature where $n \geq 4$ is an even integer. The metric g is normal if and only if $\tau(g)$ is finite. Moreover, if $\tau(g)$ is finite, one has*

$$\tau(g) = 1 - \frac{2}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} dy.$$

For brevity, set the notation α_0 as follows

$$\alpha_0 := \frac{2}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} dy.$$

Given a measurable set E , the average integral for function φ over E is defined as

$$\oint_E \varphi dx := \frac{1}{|E|} \int_E \varphi dx.$$

For $r > 0$, the notation $\bar{u}(r)$ is defined by

$$\bar{u}(r) := \frac{1}{|\partial B_r(0)|} \int_{\partial B_r(0)} u(x) d\sigma(x).$$

The following lemma is easy but useful in our following proofs.

Lemma 2.4. *For any integer $n \geq 3$, $0 < k \leq n-2$ and $r > 0$, $|y| > 0$, there holds*

$$\oint_{\partial B_r(0)} \left(\frac{|y|}{|x-y|} \right)^k d\sigma(x) \leq 1.$$

Proof. By Hölder's inequality, one has

$$\oint_{\partial B_r(0)} \left(\frac{|y|}{|x-y|} \right)^k d\sigma(x) \leq \left(\oint_{\partial B_r(0)} \left(\frac{|y|}{|x-y|} \right)^{n-2} d\sigma(x) \right)^{\frac{k}{n-2}}.$$

Thus we only need to prove the case $k = n-2$. For $n \geq 3$, $|x-y|^{n-2}$ is the Green's function. With help of mean value property of harmonic functions, we obtain the formula

$$\oint_{\partial B_r(0)} \frac{1}{|x-y|^{n-2}} d\sigma(x) = \min\{r^{2-n}, |y|^{2-n}\}$$

which has been established in Page 249 of [19] by Lieb and Loss. With help of such formula, one has

$$\oint_{\partial B_r(0)} \left(\frac{|y|}{|x-y|} \right)^{n-2} d\sigma(x) \leq 1.$$

Thus we finish the proof. \square

Lemma 2.5. *As $r \rightarrow \infty$, the normal solutions $u(x)$ to (2.1) satisfies*

$$r^2 \cdot \oint_{\partial B_r(0)} (-\Delta)u d\sigma \rightarrow (n-2)\alpha_0.$$

Proof. Firstly, since $Qe^{nu} \in L^1(\mathbb{R}^n)$, for any $\epsilon > 0$, there exists $R_\epsilon > 0$ such that

$$(2.2) \quad \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q|e^{nu} dy < \epsilon.$$

With help of Fubini's theorem, a direct computation yields that

$$\begin{aligned} & r^2 \cdot \oint_{\partial B_r(0)} (-\Delta)u d\sigma - (n-2)\alpha_0 \\ &= \frac{2(n-2)}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} \oint_{\partial B_r(0)} \frac{|x|^2}{|x-y|^2} d\sigma(x) dy \\ & \quad - \frac{2(n-2)}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} dy \\ &= \frac{2(n-2)}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} \oint_{\partial B_r(0)} \frac{|x|^2 - |x-y|^2}{|x-y|^2} d\sigma(x) dy \\ &= \frac{2(n-2)}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} \oint_{\partial B_r(0)} \frac{|y|^2 - 2y \cdot (y-x)}{|x-y|^2} d\sigma(x) dy \\ &= \frac{2(n-2)}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} \oint_{\partial B_r(0)} \frac{|y|^2}{|x-y|^2} d\sigma(x) dy \\ & \quad - \frac{4(n-2)}{(n-1)!|\mathbb{S}^n|} \oint_{\partial B_r(0)} \int_{\mathbb{R}^n} Q(y)e^{nu(y)} \frac{y \cdot (y-x)}{|x-y|^2} dy d\sigma(x). \end{aligned}$$

We will deal with these two terms one by one. Firstly, we split the second term into two parts as follows

$$\begin{aligned} & \oint_{\partial B_r(0)} \int_{\mathbb{R}^n} \frac{y \cdot (y-x)}{|x-y|^2} Q(y)e^{nu(y)} dy d\sigma(x) \\ &= \oint_{\partial B_r(0)} \int_{B_{R_\epsilon}(0)} \frac{y \cdot (y-x)}{|x-y|^2} Q(y)e^{nu(y)} dy d\sigma(x) \\ & \quad + \oint_{\partial B_r(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{y \cdot (y-x)}{|x-y|^2} Q(y)e^{nu(y)} dy d\sigma(x) \\ &=: II_1 + II_2. \end{aligned}$$

Making use of Fubini's theorem, there holds

$$|II_2| \leq C \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q(y)|e^{nu(y)} \oint_{\partial B_r(0)} \frac{|y|}{|x-y|} d\sigma(x) dy.$$

Applying Lemma 2.4 and the estimate (2.2), one has

$$(2.3) \quad |II_2| \leq C\epsilon.$$

With help of dominated convergence theorem, we have

$$(2.4) \quad II_1 \rightarrow 0, \quad \text{as } r \rightarrow \infty.$$

Then due to arbitrary choice of ϵ , one has

$$(2.5) \quad \oint_{\partial B_r(0)} \int_{\mathbb{R}^n} \frac{y \cdot (y - x)}{|x - y|^2} Q(y) e^{nu(y)} dy d\sigma(x) \rightarrow 0$$

as $r \rightarrow \infty$.

As for the first term, we also split it into two terms as before

$$\begin{aligned} & \int_{\mathbb{R}^n} Q(y) e^{nu(y)} \oint_{\partial B_r(0)} \frac{|y|^2}{|x - y|^2} d\sigma(x) dy \\ &= \int_{B_{R_\epsilon}(0)} Q(y) e^{nu(y)} \oint_{\partial B_r(0)} \frac{|y|^2}{|x - y|^2} d\sigma(x) dy \\ & \quad + \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} Q(y) e^{nu(y)} \oint_{\partial B_r(0)} \frac{|y|^2}{|x - y|^2} d\sigma(x) dy \\ &=: II_3 + II_4. \end{aligned}$$

The dominated convergence theorem yields that

$$\int_{B_{R_\epsilon}(0)} Q(y) e^{nu(y)} \frac{|y|^2}{|x - y|^2} dy \rightarrow 0, \quad \text{as } |x| \rightarrow \infty.$$

Thus we have

$$II_3 \rightarrow 0, \quad \text{as } r \rightarrow \infty.$$

With help of Lemma 2.4 and (2.2), one has

$$|II_4| \leq C\epsilon.$$

Combing these estimates and the arbitrary choice of ϵ , there holds

$$r^2 \cdot \oint_{\partial B_r(0)} (-\Delta) u d\sigma \rightarrow (n-2)\alpha_0, \quad \text{as } r \rightarrow \infty.$$

□

Lemma 2.6. *As $r \rightarrow \infty$, the normal solutions $u(x)$ to (2.1) satisfies*

$$(2.6) \quad r \cdot \bar{u}'(r) \rightarrow -\alpha_0.$$

Proof. A direct computation yields that

$$\begin{aligned} & x \cdot \nabla u(x) \\ &= -\frac{2}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} \frac{x \cdot (x - y)}{|x - y|^2} Q(y) e^{nu(y)} dy \\ &= -\alpha_0 + \frac{2}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} \frac{y \cdot (y - x)}{|x - y|^2} Q(y) e^{nu(y)} dy. \end{aligned}$$

Then, with help of the estimate (2.5), one has

$$r \cdot \bar{u}'(r) = \oint_{\partial B_r(0)} r \cdot \frac{\partial u}{\partial r} d\sigma = \oint_{\partial B_r(0)} x \cdot \nabla u d\sigma \rightarrow -\alpha_0$$

as $r \rightarrow \infty$.

□

Lemma 2.7. *As $r \rightarrow \infty$, the normal solutions $u(x)$ to (2.1) satisfies*

$$r^2 \int_{\partial B_r(0)} |\nabla u|^2 d\sigma \rightarrow \alpha_0^2.$$

Proof. With help of Fubini's theorem, a direct computation yields that

$$\begin{aligned} & |\nabla u|^2(x) \\ &= \left(\frac{2}{(n-1)!|\mathbb{S}^n|} \right)^2 \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{(x-y) \cdot (x-z)}{|x-y|^2 |x-z|^2} Q(y) e^{nu(y)} Q(z) e^{nu(z)} dy dz \\ &= \left(\frac{2}{(n-1)!|\mathbb{S}^n|} \right)^2 \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|x-y|^2 + |x-z|^2 - |y-z|^2}{2|x-y|^2 |x-z|^2} Q(y) e^{nu(y)} Q(z) e^{nu(z)} dy dz \\ &= \alpha_0 \frac{1}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} \frac{1}{|x-z|^2} Q(z) e^{nu(z)} dz \\ &\quad + \alpha_0 \frac{1}{(n-1)!|\mathbb{S}^n|} \int_{\mathbb{R}^n} \frac{1}{|x-y|^2} Q(y) e^{nu(y)} dy \\ &\quad - \left(\frac{2}{(n-1)!|\mathbb{S}^n|} \right)^2 \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|y-z|^2}{2|x-y|^2 |x-z|^2} Q(y) e^{nu(y)} Q(z) e^{nu(z)} dy dz \\ &= \frac{-\Delta u(x)}{n-2} \alpha_0 - \left(\frac{2}{(n-1)!|\mathbb{S}^n|} \right)^2 \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|y-z|^2}{2|x-y|^2 |x-z|^2} Q(y) e^{nu(y)} Q(z) e^{nu(z)} dy dz. \end{aligned}$$

Taking the same strategy as in Lemma 2.5, for any $\epsilon > 0$, there exists $R_\epsilon > 1$ such that

$$(2.7) \quad \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q| e^{nu} dx < \epsilon.$$

A direct computation yields that

$$\begin{aligned} & \left| \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|y-z|^2}{2|x-y|^2 |x-z|^2} Q(y) e^{nu(y)} Q(z) e^{nu(z)} dy dz \right| \\ & \leq \int_{B_{R_\epsilon}(0)} \int_{B_{R_\epsilon}(0)} \frac{|y-z|^2}{2|x-y|^2 |x-z|^2} |Q(y)| e^{nu(y)} |Q(z)| e^{nu(z)} dy dz \\ & \quad + \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{B_{R_\epsilon}(0)} \frac{|y-z|^2}{2|x-y|^2 |x-z|^2} |Q(y)| e^{nu(y)} |Q(z)| e^{nu(z)} dy dz \\ & \quad + \int_{B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|y-z|^2}{2|x-y|^2 |x-z|^2} |Q(y)| e^{nu(y)} |Q(z)| e^{nu(z)} dy dz \\ & \quad + \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|y-z|^2}{2|x-y|^2 |x-z|^2} |Q(y)| e^{nu(y)} |Q(z)| e^{nu(z)} dy dz \\ & =: I_1(x) + I_2(x) + I_3(x) + I_4(x). \end{aligned}$$

We are going to deal these terms one by one. With help of dominated convergence theorem, we have

$$(2.8) \quad I_1(x) \rightarrow 0, \quad \text{as } |x| \rightarrow \infty.$$

For $|x| \geq 2R_\epsilon$, $|y| \leq R_\epsilon$ and $|z| \geq R_\epsilon$, it is not hard to check

$$\frac{|y - z|^2}{2|x - z|^2|x - y|^2} \leq \frac{8|z|^2}{|x|^2|x - z|^2}.$$

Then making use of above estimate, for $|x| \geq 2R_\epsilon \geq 2$, one has

$$\begin{aligned} I_2(x) &\leq \frac{8}{|x|^2} \int_{B_{R_\epsilon}(0)} |Q(y)|e^{nu(y)} dy \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|z|^2|Q(z)|e^{nu(z)}}{|x - z|^2} dz \\ &\leq C \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|z|^2|Q(z)|e^{nu(z)}}{|x - z|^2} dz. \end{aligned}$$

Then for $|x| \geq 2R_\epsilon$, we have

$$\begin{aligned} \oint_{\partial B_r(0)} I_2(x) d\sigma(x) &\leq C \oint_{\partial B_r(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|z|^2|Q(z)|e^{nu(z)}}{|x - z|^2} dz d\sigma(x) \\ &\leq C \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q(z)|e^{nu(z)} \oint_{\partial B_r(0)} \frac{|z|^2}{|x - z|^2} d\sigma(x) dz. \end{aligned}$$

By using Lemma 2.4, for $r \geq 2R_\epsilon$, there holds

$$(2.9) \quad \oint_{\partial B_r(0)} I_2(x) d\sigma(x) \leq C \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q(z)|e^{nu(z)} dz \leq C\epsilon.$$

Similarly, one has

$$(2.10) \quad \oint_{\partial B_r(0)} I_3(x) d\sigma(x) \leq C\epsilon.$$

As for the term I_4 , one has

$$\begin{aligned} I_4(x) &= \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|x - y|^2 + |x - z|^2 - 2(x - y) \cdot (x - z)}{2|x - y|^2|x - z|^2} |Q(y)|e^{nu(y)} |Q(z)|e^{nu(z)} dy dz \\ &= \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q(y)|e^{nu(y)} dy \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{1}{2|x - z|^2} |Q(z)|e^{nu(z)} dz \\ &\quad + \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q(z)|e^{nu(z)} dz \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{1}{2|x - y|^2} |Q(y)|e^{nu(y)} dy \\ &\quad - \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{(x - y) \cdot (x - z)}{|x - y|^2|x - z|^2} |Q(y)|e^{nu(y)} |Q(z)|e^{nu(z)} dy dz \end{aligned}$$

Thus one has

$$\begin{aligned} I_4(x) &\leq \epsilon \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{1}{|x - z|^2} |Q(z)|e^{nu(z)} dz \\ &\quad + \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|Q(y)|e^{nu(y)} |Q(z)|e^{nu(z)}}{|x - y||x - z|} dy dz \\ &\leq \epsilon \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|z|^2}{|x - z|^2} |Q(z)|e^{nu(z)} dz \end{aligned}$$

$$+ \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|y| \cdot |z| \cdot |Q(y)|e^{nu(y)}|Q(z)|e^{nu(z)}}{|x-y||x-z|} dy dz$$

where we have used the fact $R_\epsilon > 1$. Then similarly as before, one has

$$\oint_{\partial B_r(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|z|^2}{|x-z|^2} |Q(z)|e^{nu(z)} dz d\sigma(x) \leq C\epsilon.$$

Using Hölder's inequality and Lemma 2.4, for $|y| > 0$ and $|z| > 0$, one has

$$\oint_{\partial B_r(0)} \frac{|y| \cdot |z|}{|x-y| \cdot |x-z|} d\sigma(x) \leq \left(\oint_{\partial B_r(0)} \frac{|y|^2}{|x-y|^2} d\sigma(x) \right)^{\frac{1}{2}} \cdot \left(\oint_{\partial B_r(0)} \frac{|z|^2}{|x-z|^2} d\sigma(x) \right)^{\frac{1}{2}} \leq 1.$$

By using Fubini's theorem, Hölder inequality as well as Lemma 2.4, one has

$$\begin{aligned} & \oint_{\partial B_r(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \frac{|y| \cdot |z| \cdot |Q(y)|e^{nu(y)}|Q(z)|e^{nu(z)}}{|x-y||x-z|} dy dz d\sigma(x) \\ & \leq \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q(y)|e^{nu(y)}|Q(z)|e^{nu(z)} \oint_{\partial B_r(0)} \frac{|y| \cdot |z|}{|x-y| \cdot |x-z|} d\sigma(x) dy dz \\ & \leq \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} \int_{\mathbb{R}^n \setminus B_{R_\epsilon}(0)} |Q(y)|e^{nu(y)}|Q(z)|e^{nu(z)} dy dz \\ & \leq \epsilon^2 \end{aligned}$$

Thus we obtain that

$$(2.11) \quad \oint_{\partial B_r(0)} I_4(x) d\sigma(x) \leq C\epsilon^2.$$

Combing these estimate (2.8), (2.9), (2.10), (2.11) and the arbitrary choice of ϵ , as $r \rightarrow \infty$, there holds

$$(2.12) \quad \left| \oint_{\partial B_r(0)} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|y-z|^2}{2|x-y|^2|x-z|^2} Q(y)e^{nu(y)}Q(z)e^{nu(z)} dy dz d\sigma(x) \right| \rightarrow 0.$$

With help of (2.12) and Lemma 2.5, one has

$$r^2 \oint_{\partial B_r(0)} |\nabla u|^2 d\sigma \rightarrow \alpha_0^2, \quad \text{as } r \rightarrow \infty.$$

□

Lemma 2.8. *The normal solutions $u(x)$ to (2.1) satisfies*

$$\bar{u}(r) = (-\alpha_0 + o(1)) \log r$$

where $o(1) \rightarrow 0$ as $r \rightarrow \infty$.

Proof. Choose $|x| \geq e^4$ such that $|x| \geq 2 \log |x|$. We split \mathbb{R}^n into three pieces

$$A_1 = B_1(x), \quad A_2 = B_{\log|x|}(0), \quad A_3 = \mathbb{R}^n \setminus (A_1 \cup A_2).$$

For $y \in A_2$ and $|y| \geq 2$, we have $|\log \frac{|x| \cdot |y|}{|x-y|}| \leq \log(2 \log |x|)$. Respectively, for $|y| \leq 2$, $|\log \frac{|x| \cdot |y|}{|x-y|}| \leq |\log |y|| + C$. Thus there holds

$$(2.13) \quad \left| \int_{A_2} \log \frac{|y|}{|x-y|} Q(y) e^{nu(y)} dy + \log |x| \int_{A_2} Q(y) e^{nu(y)} dy \right| \leq C \log \log |x| + C = o(1) \log |x|$$

where $o(1) \rightarrow 0$ as $|x| \rightarrow \infty$. For $y \in A_3$, it is not hard to check

$$\frac{1}{|x|+1} \leq \frac{|y|}{|x-y|} \leq |x|+1.$$

With help of this estimate, we could control the integral over A_3 as

$$(2.14) \quad \left| \int_{A_3} \log \frac{|y|}{|x-y|} Q(y) e^{nu(y)} dy \right| \leq \log(|x|+1) \int_{A_3} |Q| e^{nu} dy.$$

For $y \in B_1(x)$, one has $1 \leq |y| \leq |x|+1$ and then

$$\left| \int_{A_1} \log |y| Q(y) e^{nu(y)} dy \right| \leq \log(|x|+1) \int_{A_1} |Q| e^{nu} dy.$$

Since $Q e^{nu} \in L^1(\mathbb{R}^n)$, notice that $\int_{A_3 \cup A_1} |Q| e^{nu} dy \rightarrow 0$ as $|x| \rightarrow \infty$ and

$$\frac{2}{(n-1)! |\mathbb{S}^n|} \int_{A_2} Q(y) e^{nu(y)} dy = \alpha_0 + o(1).$$

Thus there holds

$$(2.15) \quad u(x) = (-\alpha_0 + o(1)) \log |x| + \frac{2}{(n-1)! |\mathbb{S}^n|} \int_{B_1(x)} \left(\log \frac{1}{|x-y|} \right) Q(y) e^{nu(y)} dy$$

where $o(1) \rightarrow 0$ as $|x| \rightarrow \infty$. For $r \gg 1$, with help of Fubini's theorem, one has

$$\begin{aligned} & \left| \int_{\partial B_r(0)} \int_{B_1(x)} \left(\log \frac{1}{|x-y|} \right) Q(y) e^{nu(y)} dy d\sigma(x) \right| \\ & \leq \int_{\partial B_r(0)} \int_{B_{r+1}(0) \setminus B_{r-1}(0)} |\log |x-y|| \cdot |Q(y)| e^{nu(y)} dy d\sigma(x) \\ & \leq \int_{B_{r+1}(0) \setminus B_{r-1}(0)} |Q(y)| e^{nu(y)} \int_{\partial B_r(0)} |\log |x-y|| d\sigma(x) dy \\ & \leq \int_{B_{r+1}(0) \setminus B_{r-1}(0)} |Q(y)| e^{nu(y)} \int_{\partial B_r(0) \cap B_1(y)} |\log |x-y|| d\sigma(x) dy \\ & \quad + \int_{B_{r+1}(0) \setminus B_{r-1}(0)} |Q(y)| e^{nu(y)} \int_{\partial B_r(0) \cap (\mathbb{R}^n \setminus B_1(y))} |\log |x-y|| d\sigma(x) dy \\ & \leq \int_{B_{r+1}(0) \setminus B_{r-1}(0)} |Q(y)| e^{nu(y)} (C + Cr^{n-1} \log(2r+1)) dy. \end{aligned}$$

Due to $Qe^{nu} \in L^1(\mathbb{R}^n)$, there holds

$$\oint_{\partial B_r(0)} \int_{B_1(x)} \left(\log \frac{1}{|x-y|} \right) Q(y) e^{nu(y)} dy d\sigma(x) = o(1) \log r.$$

With help of (2.15), we finish our proof. \square

The following lemma which has been established in [3] and [24] is also very useful and we repeat its statement for readers' convenience.

Lemma 2.9. (*Lemma 3.2 in [3], Proposition 3.1 in [24]*) *For any number $k > 0$, the normal solutions $u(x)$ to (2.1) satisfies*

$$\oint_{\partial B_r(0)} e^{ku} d\sigma = e^{k\bar{u}(r)} e^{o(1)}$$

where $o(1) \rightarrow 0$ as $r \rightarrow \infty$.

The following lemma has been established in [14]. For reader's convenience, we repeat the proof briefly.

Lemma 2.10. *Given a conformal metric $g = e^{2u}|dx|^2$ on \mathbb{R}^n with finite total Q -curvature where $n \geq 4$ is an even integer. If the metric g is normal and the volume is finite, then there holds*

$$\int_{\mathbb{R}^n} Qe^{nu} dx \geq \frac{(n-1)!|\mathbb{S}^n|}{2}.$$

Proof. By direct computation and Fubini's theorem, for $|x| \gg 1$, one has

$$\begin{aligned} & \int_{B_{|x|/2}(x)} \left| \int_{B_1(z)} \log \frac{1}{|z-y|} Q(y) e^{nu(y)} dy \right| dz \\ & \leq \int_{B_{|x|/2}(x)} \int_{B_1(z)} \frac{1}{|z-y|} |Q(y) e^{nu(y)}| dy dz \\ & \leq \int_{B_{|x|/2}(x)} \int_{B_{|x|/2+1}(x)} \frac{1}{|z-y|} |Q(y) e^{nu(y)}| dy dz \\ & \leq \int_{B_{|x|/2+1}(x)} |Q(y) e^{nu(y)}| \int_{B_{|x|/2}(x)} \frac{1}{|z-y|} dz dy \\ & \leq \int_{B_{|x|/2+1}(x)} |Q(y) e^{nu(y)}| \int_{B_{|x|+1}(0)} \frac{1}{|z|} dz dy \\ & \leq C|x|^{n-1}. \end{aligned}$$

Thus

$$(2.16) \quad \oint_{B_{|x|/2}(x)} \int_{B_1(z)} \log \frac{1}{|z-y|} Q(y) e^{nu(y)} dy dz = O(|x|^{-1}).$$

Meanwhile, for $y \in B_{|x|/2}(x)$, there holds

$$(2.17) \quad \left| \log \frac{|y|}{|x|} \right| \leq \log 2.$$

With help of these estimates (2.16), (2.17) and (2.15), we have

$$(2.18) \quad \int_{B_{|x|/2}(x)} u(y) dy = (-\alpha_0 + o(1)) \log |x|$$

where $o(1) \rightarrow 0$ as $|x| \rightarrow \infty$.

By using Jensen's inequality and (2.18), one has

$$\begin{aligned} \int_{B_{|x|/2}(x)} e^{nu(y)} dy &\geq |B_{|x|/2}(x)| \exp \left(\int_{B_{|x|/2}(x)} nu(y) dy \right) \\ &\geq C|x|^{n-n\alpha_0+o(1)}. \end{aligned}$$

Due to the assumption of finite volume, the above estimate yield that $\alpha_0 \geq 1$ i.e.

$$\int_{\mathbb{R}^n} Qe^{nu} dx \geq \frac{(n-1)!|\mathbb{S}^n|}{2}.$$

□

3. RANGE OF Q-CURVATURE INTEGRAL

In this section, we give the proofs of Theorem 1.6 and Theorem 1.8.

Proof of Theorem 1.6:

The scalar curvature R_g of $g = e^{2u}|dx|$ on \mathbb{R}^n can be written as

$$(3.1) \quad R_g = 2(n-1)e^{-2u}(-\Delta u - \frac{n-2}{2}|\nabla u|^2).$$

Firstly, for all of cases, the scalar curvature is supposed to be non-negative near infinity and then Lemma 2.1 yields that $u(x)$ is normal.

Case (1): $R_g \geq 0$ near infinity.

With help of (3.1), for $r \gg 1$, there holds

$$r^2 \int_{\partial B_r(0)} (-\Delta)u d\sigma(x) \geq \frac{n-2}{2} r^2 \int_{\partial B_r(0)} |\nabla u|^2 d\sigma(x).$$

By letting $r \rightarrow \infty$, Lemma 2.5 and Lemma 2.7 yield that

$$(n-2)\alpha_0 \geq \frac{n-2}{2}\alpha_0^2$$

which is equivalent to

$$(3.2) \quad 0 \leq \alpha_0 \leq 2.$$

Thus one has

$$0 \leq \int_{\mathbb{R}^n} Qe^{nu} dx \leq (n-1)!|\mathbb{S}^n|.$$

Case (2): $R_g \geq C > 0$ near infinity.

By using (3.1), one has

$$r^2 \int_{\partial B_r(0)} (-\Delta)u d\sigma(x) - \frac{n-2}{2} r^2 \int_{\partial B_r(0)} |\nabla u|^2 d\sigma(x)$$

$$\begin{aligned}
&= \frac{r^2}{2(n-1)} \int_{\partial B_r(0)} R_g e^{2u} d\sigma(x) \\
&\geq \frac{Cr^2}{2(n-1)} \int_{\partial B_r(0)} e^{2u} d\sigma(x)
\end{aligned}$$

Making use of Jensen's inequality and Lemma 2.8, there holds

$$\begin{aligned}
&r^2 \int_{\partial B_r(0)} (-\Delta u) d\sigma(x) - \frac{n-2}{2} r^2 \int_{\partial B_r(0)} |\nabla u|^2 d\sigma(x) \\
&\geq \frac{Cr^2}{2(n-1)} e^{2\bar{u}(r)} = \frac{C}{2(n-1)} r^{2-2\alpha_0+o(1)}.
\end{aligned}$$

If $\alpha_0 < 1$, taking $r \rightarrow \infty$ in the above inequality, we obtain a contradiction by using Lemma 2.5 and Lemma 2.7. Thus we must have

$$\alpha_0 \geq 1.$$

Combing with (3.2), we finally obtain our desired result

$$\frac{(n-1)!|\mathbb{S}^n|}{2} \leq \int_{\mathbb{R}^n} Q e^{nu} dx \leq (n-1)!|\mathbb{S}^n|.$$

Case (3): $0 \leq R_g \leq C$ near infinity.

With help of Lemma 2.9 and Lemma 2.8, for $r \gg 1$, we have

$$\begin{aligned}
&r^2 \int_{\partial B_r(0)} (-\Delta u) d\sigma(x) - \frac{n-2}{2} r^2 \int_{\partial B_r(0)} |\nabla u|^2 d\sigma(x) \\
&= \frac{r^2}{2(n-1)} \int_{\partial B_r(0)} R_g e^{2u} d\sigma(x) \\
&\leq \frac{r^2 C}{2(n-1)} \int_{\partial B_r(0)} e^{2u} d\sigma(x) = \frac{r^2 C}{2(n-1)} e^{2\bar{u}(r)} e^{o(1)} = \frac{C}{2(n-1)} r^{2-2\alpha_0+o(1)} e^{o(1)}.
\end{aligned}$$

If $\alpha_0 > 1$, by taking $r \rightarrow \infty$ in the above inequality, Lemma 2.5 and Lemma 2.7 yield that

$$(n-2)\alpha_0 - \frac{n-2}{2}\alpha_0^2 \leq 0$$

which deduces that $\alpha_0 \geq 2$. Combing with (3.2), one finally has

$$\alpha_0 = 2 \quad \text{or} \quad 0 \leq \alpha_0 \leq 1.$$

Thus we finish our proof.

Proof of Theorem 1.8:

Case (1):

Since $(R_g^-)^{\frac{n}{2}} e^{nu} \in L^1(\mathbb{R}^n)$, Lemma 2.2 yields that $u(x)$ is normal. Recalling (3.1) and applying Hölder's inequality, one has

$$r^2 \int_{\partial B_r(0)} (-\Delta u) d\sigma(x) - \frac{n-2}{2} r^2 \int_{\partial B_r(0)} |\nabla u|^2 d\sigma(x)$$

$$\begin{aligned}
&= \frac{r^2}{2(n-1)} \int_{\partial B_r(0)} R_g e^{2u} d\sigma(x) \\
&\geq -\frac{r^2}{2(n-1)} \int_{\partial B_r(0)} R_g^- e^{2u} d\sigma(x) \\
&\geq -\frac{r^2}{2(n-1)} \left(\int_{\partial B_r(0)} (R_g^-)^{\frac{n}{2}} e^{nu} d\sigma(x) \right)^{\frac{n}{2}} \\
&= -\frac{|\mathbb{S}^{n-1}|^{-\frac{n}{2}}}{2(n-1)} \left(r \int_{\partial B_r(0)} (R_g^-)^{\frac{n}{2}} e^{nu} d\sigma(x) \right)^{\frac{n}{2}}.
\end{aligned}$$

Based on the assumption (1.3), there exists a sequence $r_i \rightarrow \infty$ such that

$$(3.3) \quad r_i \int_{\partial B_{r_i}(0)} (R_g^-)^{\frac{n}{2}} e^{nu} d\sigma(x) \rightarrow 0.$$

Choosing such sequence r_i and making use of Lemma 2.5 and Lemma 2.7 as well as (3.3), we obtain that

$$(3.4) \quad (n-2)\alpha_0 - \frac{n-2}{2}\alpha_0^2 \geq 0$$

by taking $r_i \rightarrow \infty$. Thus we finish our proof.

Case (2):

Firstly, Lemma 2.2 yields that u is normal based on the assumption $R_g e^{2u} \in L^{\frac{n}{2}}(\mathbb{R}^n)$. With help of similar method as before, one has

$$\begin{aligned}
&r^2 \int_{\partial B_r(0)} (-\Delta u) d\sigma(x) - \frac{n-2}{2} r^2 \int_{\partial B_r(0)} |\nabla u|^2 d\sigma(x) \\
&= \frac{r^2}{2(n-1)} \int_{\partial B_r(0)} R_g e^{2u} d\sigma(x) \\
&\leq \frac{r^2}{2(n-1)} \int_{\partial B_r(0)} R_g^+ e^{2u} d\sigma(x) \\
&\leq \frac{r^2}{2(n-1)} \left(\int_{\partial B_r(0)} (R_g^+)^{\frac{n}{2}} e^{nu} d\sigma(x) \right)^{\frac{n}{2}} \\
&= \frac{|\mathbb{S}^{n-1}|^{-\frac{n}{2}}}{2(n-1)} \left(r \int_{\partial B_r(0)} (R_g^+)^{\frac{n}{2}} e^{nu} d\sigma(x) \right)^{\frac{n}{2}}
\end{aligned}$$

where R_g^+ is the positive part of scalar curvature. Due to (1.3), there exists a sequence $r_j \rightarrow \infty$ such that

$$r_j \int_{\partial B_{r_j}(0)} (R_g^+)^{\frac{n}{2}} e^{nu} d\sigma(x) \rightarrow 0.$$

By taking $r_j \rightarrow \infty$, Lemma 2.5 and Lemma 2.7 deduce that

$$(3.5) \quad (n-2)\alpha_0 - \frac{n-2}{2}\alpha_0^2 \leq 0.$$

Combing with (3.4), one has

$$(3.6) \quad \alpha_0 = 0 \quad \text{or} \quad 2.$$

If $e^{nu} \in L^1(\mathbb{R}^n)$ in addition, with help of Lemma 2.10, we finally obtain that

$$\alpha_0 = 2.$$

Thus we finish our proof.

4. COMPLETE MANIFOLDS WITH FINITELY MANY SIMPLE ENDS

Given a complete non-compact four dimensional manifold (M, g) , we say that it has finitely many simple ends if

$$M = N \cup \left\{ \bigcup_{i=1}^k E_i \right\}$$

where N is a compact manifold with boundary

$$\partial N = \bigcup_{i=1}^k \partial E_i$$

and each E_i is a conformally flat simple end satisfying

$$(E_i, g) = (\mathbb{R}^4 \setminus B_1(0), e^{2w_i}|dx|^2)$$

for some function w_i .

Consider the conformal metric on the end $(\mathbb{R}^4 \setminus B_1(0), e^{2w}|dx|^2)$ where

$$(4.1) \quad (-\Delta)^{\frac{n}{2}} w(x) = Q(x)e^{4w(x)}, \quad x \in \mathbb{R}^4 \setminus B_1(0).$$

We also say that such metric has finite total Q-curvature if $Qe^{4w} \in L^1(\mathbb{R}^4 \setminus B_1(0))$. Different from the case on \mathbb{R}^4 , we say the solution to (4.1) with finite total Q-curvature is normal if $w(x)$ has the following decomposition

$$(4.2) \quad w(x) = \frac{1}{8\pi^2} \int_{\mathbb{R}^4 \setminus B_1(0)} \log \frac{|y|}{|x-y|} Q(y)e^{4u(y)} dy + \alpha_1 \log |x| + h(x)$$

where α_1 is a constant and $h(\frac{x}{|x|^2})$ is some biharmonic function on $B_1(0)$.

When w is normal, such conformal metric $e^{2w}|dx|^2$ is said to be normal. For brevity, set the notation α_2 as follows

$$\alpha_2 := \frac{1}{8\pi^2} \int_{\mathbb{R}^4 \setminus B_1(0)} Qe^{4u} dy.$$

More details about normal metric on $\mathbb{R}^4 \setminus B_1(0)$ can be found in [4].

Leveraging the definition of w and the lemmas outlined in the preceding section, we are able to establish the following lemmas for future use.

Lemma 4.1. *For the normal solution $w(x)$ to (4.2), as $r \rightarrow \infty$, there holds*

(1)

$$r^2 \oint_{\partial B_r(0)} |\nabla w|^2 d\sigma(x) \rightarrow (\alpha_2 - \alpha_1)^2,$$

(2)

$$r^2 \oint_{\partial B_r(0)} (-\Delta)w d\sigma(x) \rightarrow 2(\alpha_2 - \alpha_1).$$

Proof. For brevity, we set

$$w_1(x) := \frac{1}{8\pi^2} \int_{\mathbb{R}^4 \setminus B_1(0)} \log \frac{|y|}{|x-y|} Q(y) e^{4u(y)} dy,$$

and $b(x) := h(\frac{x}{|x|^2})$. Thus $b(x)$ is biharmonic on $B_1(0)$. A direct computation yields that

$$(4.3) \quad \partial_i h(x) = \frac{\partial_i b}{|x|^2} - \sum_{j=1}^4 \partial_j b \frac{2x_i x_j}{|x|^4}$$

which implies that

$$(4.4) \quad |x| \cdot |\nabla h(x)| \rightarrow 0, \quad \text{as } |x| \rightarrow \infty.$$

With help of Lemma 2.7 and Lemma 2.6, one has

$$(4.5) \quad r^2 \oint_{\partial B_r(0)} |\nabla w_1|^2 d\sigma(x) \rightarrow \alpha_2^2$$

and

$$(4.6) \quad \oint_{\partial B_r(0)} x \cdot \nabla w_1 d\sigma(x) \rightarrow -\alpha_2$$

as $r \rightarrow \infty$. A direct computation yields that

$$\begin{aligned} & r^2 \oint_{\partial B_r(0)} |\nabla w|^2 d\sigma(x) \\ &= r^2 \oint_{\partial B_r(0)} |\nabla w_1|^2 d\sigma(x) + \alpha_1^2 r^2 \oint_{\partial B_r(0)} |\nabla \log |x||^2 d\sigma(x) + r^2 \oint_{\partial B_r(0)} |\nabla h|^2 d\sigma(x) \\ & \quad + 2\alpha_1 r^2 \oint_{\partial B_r(0)} \nabla w_1 \cdot \nabla \log |x| d\sigma(x) + 2r^2 \oint_{\partial B_r(0)} \nabla w_1 \cdot \nabla h d\sigma(x) \\ & \quad + 2\alpha_1 r^2 \oint_{\partial B_r(0)} \nabla h \cdot \nabla \log |x| d\sigma(x) \\ &= r^2 \oint_{\partial B_r(0)} |\nabla w_1|^2 d\sigma(x) + \alpha_1^2 + r^2 \oint_{\partial B_r(0)} |\nabla h|^2 d\sigma(x) \\ & \quad + 2\alpha_1 \oint_{\partial B_r(0)} x \cdot \nabla w_1 d\sigma(x) + 2r^2 \oint_{\partial B_r(0)} \nabla w_1 \cdot \nabla h d\sigma(x) \\ & \quad + 2\alpha_1 \oint_{\partial B_r(0)} x \cdot \nabla h d\sigma(x) \end{aligned}$$

With help of (4.4), one has

$$r^2 \int_{\partial B_r(0)} |\nabla h|^2 d\sigma(x) \rightarrow 0, \quad \text{as } r \rightarrow \infty$$

and

$$\int_{\partial B_r(0)} x \cdot \nabla h d\sigma(x) \rightarrow 0, \quad \text{as } r \rightarrow \infty.$$

Making use of Hölder's inequality and the estimates (4.4), (4.5), one has

$$\left| r^2 \int_{\partial B_r(0)} \nabla w_1 \cdot \nabla h d\sigma \right| \leq \left(r^2 \int_{\partial B_r(0)} |\nabla w_1|^2 d\sigma \right)^{\frac{1}{2}} \left(r^2 \int_{\partial B_r(0)} |\nabla h|^2 d\sigma \right)^{\frac{1}{2}} \rightarrow 0$$

as $r \rightarrow \infty$. Combing these estimates, we have

$$r^2 \int_{\partial B_r(0)} |\nabla w|^2 d\sigma(x) \rightarrow (\alpha_2 - \alpha_1)^2, \quad \text{as } r \rightarrow \infty.$$

Using the same argument as (4.3), it is not hard to check that

$$(4.7) \quad |x|^2 \Delta h(x) \rightarrow 0, \quad \text{as } |x| \rightarrow \infty.$$

With help of Lemma 2.5 and the fact $n = 4$, one has

$$(4.8) \quad r^2 \int_{\partial B_r(0)} (-\Delta) w_1 d\sigma \rightarrow 2\alpha_2, \quad \text{as } r \rightarrow \infty.$$

Combing these estimates (4.7) and (4.8), there holds

$$\begin{aligned} & r^2 \int_{\partial B_r(0)} (-\Delta) w d\sigma(x) \\ &= r^2 \int_{\partial B_r(0)} (-\Delta) w_1 d\sigma(x) + r^2 \alpha_1 \int_{\partial B_r(0)} (-\Delta) \log |x| d\sigma(x) \\ & \quad + r^2 \int_{\partial B_r(0)} (-\Delta) h d\sigma(x) \\ &= r^2 \int_{\partial B_r(0)} (-\Delta) w_1 d\sigma(x) - 2\alpha_1 + r^2 \int_{\partial B_r(0)} (-\Delta) h d\sigma(x) \\ & \rightarrow 2\alpha_2 - 2\alpha_1, \quad \text{as } r \rightarrow \infty. \end{aligned}$$

Thus we finish our proof. \square

Lemma 4.2. *For the normal solution $w(x)$ to (4.2), there holds*

(1)

$$r \cdot \bar{w}'(r) \rightarrow -\alpha_2 + \alpha_1, \quad \text{as } r \rightarrow \infty.$$

(2)

$$\bar{w}(r) = (-\alpha_2 + \alpha_1 + o(1)) \log r$$

where $o(1) \rightarrow 0$ as $r \rightarrow \infty$.

Proof. A direct computation yields that

$$\begin{aligned} r \cdot \bar{w}'(r) &= \oint_{\partial B_r(0)} x \cdot \nabla w_1 d\sigma(x) + \alpha_1 \oint_{\partial B_r(0)} x \cdot \nabla \log |x| d\sigma(x) + \oint_{\partial B_r(0)} x \cdot \nabla h d\sigma(x) \\ &= \oint_{\partial B_r(0)} x \cdot \nabla w_1 d\sigma(x) + \alpha_1 + \oint_{\partial B_r(0)} x \cdot \nabla h d\sigma(x). \end{aligned}$$

By using Lemma 2.6 and the estimate (4.4), one has

$$r \cdot \bar{w}'(r) \rightarrow -\alpha_2 + \alpha_1, \quad \text{as } r \rightarrow \infty.$$

Since $h(\frac{x}{|x|^2})$ is biharmonic in $B_1(0)$, it is easy to see that

$$|h(x)| \leq C, \quad \text{as } |x| \rightarrow \infty.$$

Using such fact and Lemma 2.8, there holds

$$\bar{w}(r) = \bar{w}_1(r) + \alpha_1 \log r + \oint_{\partial B_r(0)} h d\sigma = (-\alpha_2 + \alpha_1 + o(1)) \log r.$$

□

Similar to the case for \mathbb{R}^n , Chang, Qing, and Yang [4] showed that if the scalar curvature is non-negative near infinity, the conformal metric $e^{2w}|dx|^2$ on $\mathbb{R}^4 \setminus B_1(0)$ is also normal. For readers' convenience, we repeat their result as follows.

Proposition 4.3. *(See Proposition 1.12 in [4]) Given a conformal metric $(\mathbb{R}^4 \setminus B_1(0), e^{2w}|dx|^2)$ with finite total Q -curvature. Supposing that the scalar curvature is non-negative near infinity, then the metric is normal.*

Set the isoperimetric ratio for $(\mathbb{R}^4 \setminus B_1(0), e^{2w}|dx|^2)$ as follows

$$I_g(r) := \frac{\left(\int_{\partial B_r(0)} e^{3w} d\sigma(x) \right)^{\frac{4}{3}}}{4(2\pi^2)^{\frac{1}{3}} \int_{B_r(0) \setminus B_1(0)} e^{4w} dx}.$$

Chang, Qing and Yang [4] showed that the limit of such isoperimetric ratio exists as r tends to infinity if the metric is complete and normal.

Proposition 4.4. *(Proposition 1.11 in [4]) Suppose that $(\mathbb{R}^4, e^{2w}|dx|^2)$ is a complete normal metric. Then there holds*

$$\lim_{r \rightarrow \infty} I_g(r) = \lim_{r \rightarrow \infty} (1 + r \cdot \bar{w}'(r)) \geq 0.$$

With the aid of the aforementioned preparations, we aim to give the control of isoperimetric ratio by constraining the lower bound of scalar curvature.

Lemma 4.5. *Given a complete metric $g = e^{2w}|dx|^2$ on $\mathbb{R}^4 \setminus B_1(0)$ with finite total Q -curvature. If the scalar curvature $R_g \geq 0$ near infinity, then there holds*

$$0 \leq \lim_{r \rightarrow \infty} I_g(r) \leq 1.$$

Proof. Firstly, due to $R_g \geq 0$ near infinity, Proposition 4.3 yields that w is normal. Recall the definition R_g for $(\mathbb{R}^4 \setminus B_1(0), e^{2w}|dx|^2)$

$$(4.9) \quad R_g = 6e^{-2w} (-\Delta w - |\nabla w|^2).$$

Based on the assumption that $R_g \geq 0$ near infinity, for $r \gg 1$, there holds

$$r^2 \int_{\partial B_r(0)} (-\Delta)w d\sigma \geq r^2 \int_{\partial B_r(0)} |\nabla w|^2 d\sigma.$$

Making use of Lemma 4.1, letting $r \rightarrow \infty$, we have

$$(4.10) \quad 0 \leq \alpha_2 - \alpha_1 \leq 2.$$

By using Lemma 4.2 and Proposition 4.4, there holds

$$(4.11) \quad \lim_{r \rightarrow \infty} I_g(r) = 1 + \alpha_1 - \alpha_2 \geq 0.$$

Combing with (4.10), we finally obtain that

$$0 \leq \lim_{r \rightarrow \infty} I_g(r) \leq 1.$$

□

Lemma 4.6. *Given a complete metric $g = e^{2w}|dx|^2$ on $\mathbb{R}^4 \setminus B_1(0)$ with finite total Q-curvature. If $R_g \geq C$ for some constant $C > 0$ near infinity, there holds*

$$\lim_{r \rightarrow \infty} I_g(r) = 0.$$

Proof. Due to same reason as before, w is normal. Using Jensen's inequality and Lemma 4.2, for $r \gg 1$, one has

$$\begin{aligned} & r^2 \int_{\partial B_r(0)} (-\Delta)w d\sigma - r^2 \int_{\partial B_r(0)} |\nabla w|^2 d\sigma \\ &= \frac{1}{6} r^2 \int_{\partial B_r(0)} R_g e^{2u} d\sigma \\ &\geq \frac{C}{6} r^2 \int_{\partial B_r(0)} R_g e^{2u} d\sigma \\ &\geq \frac{C}{6} r^2 e^{2\bar{w}(r)} \\ &= \frac{C}{6} r^{2-2\alpha_2+2\alpha_1+o(1)}. \end{aligned}$$

With help of Lemma 4.1, letting $r \rightarrow \infty$ in the above inequality, we must have

$$2 - 2\alpha_2 + 2\alpha_1 \leq 0.$$

Combing with (4.11), we finally obtain that

$$\lim_{r \rightarrow \infty} I_g(r) = 0.$$

□

To prove Theorem 1.4 and Theorem 1.5, we just need to combining the following result which has been established in [4] with Lemma 4.5 and Lemma 4.6 respectively.

Theorem 4.7. *(See Corollary in [4]) Given a complete four dimensional manifold (M^4, g) with only finitely many conformally flat simple ends and finite total Q -curvature. Suppose that the scalar curvature is non-negative at each end. Then there holds*

$$\chi(M) - \frac{1}{32\pi^2} \int_M (|W|^2 + 4Q) d\mu_g = \sum_{i=1}^k \nu_i$$

where in the inverted coordinates centered at each end,

$$\nu_i = \lim_{r \rightarrow \infty} \frac{(\text{vol}(\partial B_r))^{\frac{4}{3}}}{4(2\pi^2)^{\frac{1}{3}} \text{vol}(B_r \setminus B_1(0))}.$$

REFERENCES

- [1] M. Bonk, J. Heinonen, and F. Saksman, Logarithmic potentials, quasiconformal flows, and Q -curvature, *Duke Math. J.* 142(2008), no.2, 197–239.
- [2] T. P. Branson, Differential operators canonically associated to a conformal structure, *Math. Scand.* 57:2 (1985).
- [3] S.-Y. A. Chang, J. Qing, and P. C. Yang, On the Chern-Gauss-Bonnet integral for conformal metrics on \mathbb{R}^4 , *Duke Math. J.* 103, No. 3, 523–544 (2000).
- [4] S.-Y. A. Chang, J. Qing, and P. C. Yang, Compactification of a class of conformally flat 4-manifold, *Invent. Math.* 142 (2000), no. 1, 65–93.
- [5] S.-Y. A. Chang and P. C. Yang, On uniqueness of solutions of n th order differential equations in conformal geometry, *Math. Res. Lett.* 4(1997), no.1, 91–102.
- [6] W. Chen and C. Li, Classification of solutions of some nonlinear elliptic equations, *Duke Math. J.* 63 (1991), no. 3, 615–622.
- [7] S.-S. Chern, A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds, *Ann. of Math.* (2) 45(1944), 747–752.
- [8] S. Cohn-Vossen, Kürzeste Wege und Totalkrümmung auf Flächen, *Compositio Math.* 2 (1935), 69–133.
- [9] H. Fang, On a conformal Gauss–Bonnet–Chern inequality for LCF manifolds and related topics, *Calc. Var.* 23, 469–496 (2005).
- [10] R. Finn, On a class of conformal metrics, with application to differential geometry in the large, *Comment. Math. Helv.* 40 (1965), 1–30.
- [11] R. Greene and H. Wu, C^∞ convex functions and manifolds of positive curvature, *Acta Math.* 137(1976), no.3-4, 209–245.
- [12] A. Huber, On subharmonic functions and differential geometry in the large, *Comment. Math. Helv.* 32 (1957) 13–72.
- [13] M. Li, A Liouville-type theorem in conformally invariant equations, *Math. Ann.* (2023) <https://doi.org/10.1007/s00208-023-02712-9>.
- [14] M. Li, The total Q -curvature, volume entropy and polynomial growth polyharmonic functions, *arXiv:2306.15623*.
- [15] M. Li, Obstructions to prescribed Q -curvature of complete conformal metric on \mathbb{R}^n , *arXiv:2401.03457*.
- [16] M. Li and J. Wei, Higher order Bol’s inequality and its applications, *arXiv:2308.11388*.
- [17] P. Li and L.-F. Tam, Complete surfaces with finite total curvature, *J. Differential Geom.* 33(1991), no.1, 139–168.

- [18] Y. Li and Z. Wang, Manifolds for which Huber's Theorem holds, *Calc. Var.* 63, 86 (2024). <https://doi.org/10.1007/s00526-024-02686-9>
- [19] E. Lieb and M. Loss, *Analysis*, Second edition, *Grad. Stud. Math.*, 14 American Mathematical Society, Providence, RI, 2001. xxii+346 pp.
- [20] C.-S. Lin, A classification of solutions of a conformally invariant fourth order equation in R^n , *Comment. Math. Helv.* 73 (1998), no. 2, 206–231.
- [21] Z. Lu and Y. Wang, On locally conformally flat manifolds with finite total Q-curvature, *Calc. Var. Partial Differential Equations* 56(2017), no.4, Paper No. 98, 24 pp.
- [22] L. Martinazzi, Classification of solutions to the higher order Liouville's equation on \mathbb{R}^{2m} , *Math. Z.* 263 (2009), no. 2, 307–329.
- [23] S. Müller and V. Šverák, On surfaces of finite total curvature, *J. Differential Geom.* 42(1995), no.2, 229–258.
- [24] C. Ndiaye and J. Xiao, An upper bound of the total Q-curvature and its isoperimetric deficit for higher-dimensional conformal Euclidean metrics, *Calc. Var. Partial Differential Equations*, volume 38, pages 1–27 (2010)
- [25] S. A. Paneitz, A quartic conformally covariant differential operator for arbitrary pseudo-Riemannian manifolds, *Symmetry Integrability Geom. Methods Appl.* 4, paper 036..
- [26] W. A. Jr. Poor, Some results on nonnegatively curved manifolds, *J. Differential Geometry* 9(1974), 583–600.
- [27] R. Schoen, S.-T. Yau, *Lectures on differential geometry*, *Conference Proceedings and Lecture Notes in Geometry and Topology*, I. International Press, Cambridge, MA, 1994.
- [28] S. Wang and Y. Wang, Integrability of scalar curvature and normal metric on conformally flat manifolds, *J. Differ. Equations* 265, No. 4, 1353–1370 (2018).
- [29] Y. Wang, The isoperimetric inequality and Q-curvature, *Adv. Math.* 281(2015), 823–844.
- [30] Z. Wang and J. Zhou, Three Circles Theorem for Volume of Conformal Metrics, *Commun. Math. Stat.* (2024). <https://doi.org/10.1007/s40304-024-00394-6>.
- [31] J. Wei and X. Xu, Classification of solutions of higher order conformally invariant equations, *Math. Ann.* 313 (1999), no. 2, 207–228.
- [32] X. Xu, Uniqueness and non-existence theorems for conformally invariant equations, *J. Funct. Anal.* 222 (2005), no.1, 1–28.

MINGXIANG LI, NANJING UNIVERSITY, CHINA
Email address: limx@smail.nju.edu.cn