# arXiv:2405.10368v1 [quant-ph] 16 May 2024

# Trapped-Ion Quantum Simulation of Electron Transfer Models with Tunable Dissipation

Visal So,<sup>1, \*</sup> Midhuna Duraisamy Suganthi,<sup>1, 2, \*</sup> Abhishek Menon,<sup>1</sup> Mingjian Zhu,<sup>1</sup> Roman

Zhuravel,<sup>1</sup> Han Pu,<sup>1</sup> Peter G. Wolvnes,<sup>1,3,4,5</sup> José N. Onuchic,<sup>1,3,4,5</sup> and Guido Pagano<sup>1,†</sup>

<sup>1</sup>Department of Physics and Astronomy, Rice University, Houston, TX 77005, USA

<sup>2</sup>Applied Physics Graduate Program, Smalley-Curl Institute, Rice University, Houston, TX 77005, USA

<sup>3</sup>Department of Chemistry, Rice University, Houston, TX 77005, USA

<sup>4</sup>Center for Theoretical Biological Physics, Rice University, Houston, TX 77005, USA

<sup>5</sup>Department of Biosciences, Rice University, Houston, TX 77005, USA

Electron transfer is at the heart of many fundamental physical, chemical, and biochemical processes essential for life. Exact simulation of reactions in these systems is often hindered by the large number of degrees of freedom and by the essential role of quantum effects. In this work, we experimentally simulate a paradigmatic model of molecular electron transfer using a multi-species trapped-ion crystal, where the donor-acceptor gap, the electronic and vibronic couplings, and the bath relaxation dynamics can all be controlled independently. We employ the ground-state qubit of one ion to simulate the electronic degree of freedom and the optical qubit of another ion to perform reservoir engineering on a collective mode encoding a reaction coordinate. We observe the real-time dynamics of the spin excitation, measuring the transfer rate in several regimes of adiabaticity and relaxation dynamics. The setup allows access to the electron transfer dynamics in the non-perturbative regime, where there is no clear hierarchy among the energy scales in the model, as has been suggested to be optimal for many rate phenomena, including photosynthesis. Our results provide a testing ground for increasingly rich models of molecular excitation transfer processes that are relevant for molecular electronics and light-harvesting systems.

Quantum devices hold the promise to provide an advantage in directly simulating many-body quantum systems [1]. Chemical reaction dynamics provides a wide range of target applications. Fully realistic digitization of the real-time dynamics of molecules on fault-tolerant quantum computers, however, requires qubit numbers and circuit depths that exceed the current state-of-theart [2]. A promising alternative approach is to develop programmable analog quantum simulators [3–5] that map the dynamical degrees of freedom of a molecule directly onto the quantum hardware, therefore providing a more direct but problem-specific quantum advantage.

One outstanding challenge is modeling the real-time electron transfer (ET) dynamics in molecular systems embedded in biological environments. In these systems, the energy differences between the electronic states, molecular vibrational energies, and their mutual couplings are all of the same order of magnitude. This requires simulating electronic excitations while taking into account a large number of nuclear degrees of freedom. Additionally, reactions at low temperatures in many molecular systems, ranging from myoglobin ligand recombination [6] to charge transport in DNA strands [7], suggest that quantum effects play a key role.

In many regimes, the reaction dynamics can be treated using imaginary-time path-integral methods [8–10]. It has also proven expedient to treat the nuclear and electronic degrees of freedom using a mix of quantum and classical dynamics [11], but the limits of this approach are not always clear. When quantum coherences between the electronic and vibrational degrees of freedom [12, 13] are relevant, such approaches are only approximate. Methods based on the hierarchical equations of motion approach [14] or real-time path-integral evaluations have also made progress in those regimes [15, 16].

Recently, the high degree of control and tunability of programmable quantum platforms such as trapped ions, superconducting qubit, and photonic simulators have been used to experimentally simulate models of vibrationally-assisted energy transfer [17], conical intersections [18–20], noise-assisted excitation transfer [21, 22], electron transfer driven by polarized light [23], and molecular vibrational dynamics [24].

In this work, we show that a trapped-ion quantum simulator with independent control of unitary and dissipative processes can successfully simulate a paradigmatic ET model. This is achieved by manipulating two different atomic ion species and employing both ground-state and optical qubits, combining spin and spin-motion coherent manipulation with sympathetic cooling of a collective bosonic mode. This programmable open quantum system enables the measurement of the time-resolved dynamics of the system in contact with an engineered bosonic bath, accessing non-perturbative regimes where electronic and vibrational excitations, their mutual coupling, and the relaxation rate are all of the same order of magnitude.

An effective model that describes ET is the celebrated spin-boson model [25]. Here, the electronic degrees of freedom are mapped onto a two-level system coupled to a bath of harmonic vibrations encoded in a collection of bosonic modes. This model involves one two-level sys-

<sup>\*</sup> These authors gave equal contributions.

<sup>&</sup>lt;sup>†</sup> pagano@rice.edu



FIG. 1. (a) Experimental setup: A  $^{171}$ Yb<sup>+</sup>  $_{-}^{172}$ Yb<sup>+</sup> ion crystal confined in a harmonic potential with Coulomb interactions defining normal modes of motion. The ground-state qubit of  $^{171}$ Yb<sup>+</sup> encodes the spin degree of freedom and is coherently manipulated by two counter-propagating 355 nm Raman beams (purple arrows, with green arrows showing the light polarization). The optical qubit of  $^{172}$ Yb<sup>+</sup> is addressed with a 435 nm laser (blue arrow) and, together with a 935 nm repumper (brown line in the inset), is used for sympathetic cooling. Insets: simplified level scheme for  $^{171}$ Yb<sup>+</sup> and  $^{172}$ Yb<sup>+</sup> .(b) Donor (red) and acceptor (blue) surfaces defined by the Hamiltonian (1) with parameters ( $V_x, g, \Delta E$ ) = (0.06, 1.6, 1.56) $\omega$  shown as a function of the reaction coordinate y with their respective non-interacting harmonic wavefunctions. The bath is represented by vibrational modes with a finite linewidth  $\gamma$ . The color hue reflects the weights of the spin population at each position y. (c) Donor population dynamics governed by unitary (purple circles) and dissipative (blue circles) evolution with ( $V_x, g, \Delta E$ ) = (0.18, 1, 1) $\omega$  compared to the numerical results (Eq. (2), solid lines) with  $\gamma = 0$  (purple) and  $\gamma = 0.014\omega$  (blue), respectively, including spin decoherence ( $\gamma_z = 0.0013\omega$ ) and motional dephasing ( $\gamma_m = 0.0013\omega$ ) (see Methods). Error bars are the statistical standard error around the mean.

tem, encoding the electron donor and acceptor states and a reaction coordinate encoded in a single bosonic mode, which is, in turn, itself coupled to a continuous bath of harmonic oscillators [26, 27]. Despite its simplicity, this model allows experimental access to paradigmatic ET regimes by measuring the real-time dynamics of the two-level system and extracting the transfer rate as a function of its coupling to the bosonic mode, the electronic donor-acceptor coupling, their energy difference, and the relaxation rate. The central system is described by the following Hamiltonian [26, 28], which is a variant of the Rabi model [29] in quantum optics ( $\hbar = 1$ ):

$$H_{\rm s} = \frac{\Delta E}{2}\sigma_z + V_x\sigma_x + \frac{g}{2}\sigma_z(a^{\dagger} + a) + \omega a^{\dagger}a, \quad (1)$$

where  $\sigma_{x,z}$  are the Pauli matrices and  $a^{\dagger}(a)$  is the creation (annihilation) operator of the bosonic mode at frequency  $\omega$ . The reaction coordinate is expressed in terms of the position operator as  $y = y_0(a^{\dagger} + a)/2$ , with  $y_0 = \sqrt{1/2m\omega}$ and m being the particle mass. In this model, when  $V_x = 0$ , the energy spectrum is described by two harmonic wells assigned to the donor and acceptor states,  $|D\rangle \equiv |\uparrow\rangle_z$  and  $|A\rangle \equiv |\downarrow\rangle_z$  separated by a relative energy shift (a.k.a. exothermicity)  $\Delta E$ . The electronic coupling  $V_x$  mixes the states associated with the donor and acceptor surfaces. The spin-boson coupling g displaces the two coupled surfaces along the reaction coordinate, as shown in Fig. 1b. In electron transfer, this is akin to the nuclear coupling that gives rise to the activation energy of a typical electron transfer reaction, which is the core of the Marcus theory [30] in chemistry and polaron theory in solid state physics [31].

Crucially, the full ET Hamiltonian  $H_{ET} = H_{\rm s} + H_{\rm b} +$  $H_{\rm sb}$  must also include bath degrees of freedom  $H_{\rm b}$ , generally modeled as a large collection of harmonic oscillators, and a linear coupling  $H_{\rm sb}$  between the bath and the system's bosonic degree of freedom [26]. The bath correlation functions and their effect on the system can be described by a continuous spectral density function  $J(\omega)$ . One way to create an analog for the structured bath spectral densities of biological environments using trapped ions is to employ multiple phononic modes naturally hosted in an ion crystal [32, 33]. Here, we take a different approach by exploiting the fact that, under certain conditions, a harmonic environment with a continuous spectral density can be obtained by cooling a spectator ion [34]. In section I of Supplementary Information, we prove that sympathetic cooling can effectively simulate an Ohmic spectral density  $J(\omega) \sim \omega$ , a common choice in electron transfer literature. The cooling process can be described by a master equation in terms of Lindbladian super-operators  $\mathcal{L}_c[\rho]$ , where c is a generic jump operator:

$$\frac{\partial \rho}{\partial t} = -i[H_{\rm s},\rho] + \gamma(\bar{n}+1)\mathcal{L}_a[\rho] + \gamma \bar{n}\mathcal{L}_{a^{\dagger}}[\rho], \quad (2)$$

$$\mathcal{L}_c[\rho] = c\rho c^{\dagger} - \frac{1}{2} \{ c^{\dagger} c, \rho \}.$$
(3)

Here,  $\rho$  is the density matrix of the spin-boson system,  $\gamma$  is the motional relaxation rate, and  $\bar{n}$  is the phonon population determined by the temperature of the bath  $k_B T = \omega / \log(1 + 1/\bar{n}).$ 

The dynamics of the spin and the bosonic observables predicted by Eq. (2) are essentially indistinguishable from those of the system in Eq. (1) in contact with an Ohmic bath, provided that the damping is weak ( $\gamma \ll \omega$ ) and the bath thermal energy is larger than the relaxation rate ( $\gamma\beta \ll 1$ , with  $\beta = 1/k_BT$ ) [34]. As shown in the following, these conditions can be realized experimentally with a trapped-ion system, where the dynamics is determined by five parameters ( $\omega, \Delta E, V_x, g, \gamma$ ) that can be all tuned independently. Notably, all the timescales associated with these parameters are faster than the spin and motional decoherence associated with experimental imperfections (see Fig. 1c and Methods), allowing the full characterization of both the transient dynamics and the steady state of the system under Eq. (2).

The experimental setup consists of one <sup>171</sup>Yb<sup>+</sup> ion and one <sup>172</sup>Yb<sup>+</sup> ion confined in a linear Paul trap. The two-level system is encoded in the ground state qubit in the two <sup>171</sup>Yb<sup>+</sup> clock hyperfine states  $|^{2}S_{1/2}, F = 1, m_{F} = 0\rangle \equiv |\uparrow\rangle_{z}$  and  $|^{2}S_{1/2}, F = 0, m_{F} = 0\rangle \equiv |\downarrow\rangle_{z}$  (see Fig. 1a), separated by a frequency of  $\omega_{\rm hf}/2\pi = 12.642$  GHz. The bosonic mode in the Hamiltonian (1) is encoded in the radial tilt collective mode at frequency  $\omega_{\rm tilt}/2\pi = 3.207$  MHz (see Methods).

We engineer  $H_s$  in Eq. (1) in a driven rotating frame: two  $\pi/2$  pulses are used to map the z spin basis of Eq. (1) onto the y basis. In this configuration, two laser tones resonant with the qubit frequency realize the  $\Delta E$ and  $V_x$  terms. Two additional laser tones at frequencies  $\pm \mu = \pm (\omega_{\text{tilt}} + \delta)$  realize the spin-phonon coupling and the harmonic term in Eq. (1), where  $\delta \equiv -\omega$  is the detuning with respect to the tilt mode [35]. All the terms in  $H_s$  are engineered using a 355 nm pulsed laser addressing the <sup>171</sup>Yb<sup>+</sup> ground-state qubit via stimulated Raman transitions (see Fig. 1a and Methods).

Simulating an independently tunable bath dissipation is achieved by driving the narrow transition from the ground  $|g\rangle \equiv |^2S_{1/2}\rangle$  to the optical metastable state  $|o\rangle \equiv |^2D_{3/2}\rangle$  of a <sup>172</sup>Yb<sup>+</sup> ion. Two tones of a 435 nm laser combined with a 935 nm repumper are used to perform sympathetic cooling [36] on the tilt mode with a cooling rate  $\gamma/2\pi$ , which is tunable over the 50-500 Hz range (see Fig. 1a). This setting is well suited to achieve efficient sympathetic cooling because the fractional mass imbalance of the two ions is very small, and the  $|g\rangle \rightarrow |o\rangle$ transition linewidth allows for large Rabi frequencies at modest laser power while providing negligible crosstalk with the qubit states of <sup>171</sup>Yb<sup>+</sup>.

The experimental protocol consists of the following steps: (i) After Doppler cooling, Raman resolvedsideband cooling is applied to both the radial centerof-mass and tilt modes. The resulting initial tilt mode phonon population is in the  $\bar{n}_0 \sim (0.1-0.3)$  range, which is comparable to  $\bar{n}$  defined in Eq. (2). (ii) Then, by applying a  $\pi/2$  pulse followed by a displacement operator  $\mathcal{D}(-g/2\omega)$ , we initialize the system in the donor state  $|D\rangle \langle D| \otimes \rho_-$ , where  $\rho_- = \sum_n e^{-n\omega/k_B T} |n_-\rangle \langle n_-|$  is a thermal state with temperature  $k_B T \approx \omega/\log(1+1/\bar{n})$ and  $|n_{\pm}\rangle = \mathcal{D}(\pm g/2\omega) |n\rangle$  are displaced Fock states. (iii) We simultaneously apply the laser tones to generate the ET dynamics described by Eq. (2). All the parameters that determine the unitary and the dissipative evolution are calibrated independently (see Methods). *(iv)* At the end of the evolution, after a final  $\pi/2$  pulse, we use state-dependent fluorescence to measure the probability of the system being in the donor state  $P_D = (\langle \sigma_z \rangle + 1)/2$  or the average phonon population  $\langle n \rangle$  of the tilt mode.

The average number of phonons  $\bar{n}$  in the (0.1-0.3) range fulfills the condition  $k_BT \leq \omega$  while making sure that the constraint  $\gamma \ll k_BT$  is also satisfied. In this highly quantum regime, the transfer is dominated by the discrete level structure of the vibrational mode, and the temperature has a limited effect on the transfer rate. This corresponds to the low-temperature, tunnelling-dominated regime of electron transfer.

A crucial parameter for the ET dynamics is the Marcus reorganization energy  $\lambda = g^2/\omega$ , which is the amount of energy required to displace a wavepacket by  $g/\omega$  from the center of the donor surface without transferring to the acceptor surface (see Fig. 1(b)). The reorganization energy, in turn, determines the classical activation energy  $U = (\Delta E + \lambda)^2/4\lambda$ , which is the barrier a wavepacket localized in the donor surface would have to overcome to enter the acceptor surface when the electronic coupling  $V_x$  is negligible.

We individuate and investigate two regimes [28]: a non-adiabatic and a strongly adiabatic transfer regime. In the former, the electronic coupling  $V_x$  is a small perturbation with respect to the other energy scales in the Hamiltonian (1) and is comparable or smaller than the relaxation rate  $(V_x \leq \gamma)$ . When  $V_x$  is also much less than  $\lambda/4$ , namely the activation energy at  $\Delta E = 0$ , the bosonic wavepacket is largely localized on either the donor or the acceptor surface, and the ET can be described by the Fermi golden rule (FGR) leading to characteristic isolated peaks. Conversely, in the strongly adi*abatic* regime, the electronic coupling becomes comparable with the activation energy  $(V_x \sim \lambda/4)$  and greater than the relaxation rate  $(V_x > \gamma)$ , changing the shapes of the BO surfaces. In this regime, the transfer rate is less sensitive to the electronic coupling  $V_x$  and cannot be predicted in terms of the FGR. Increasing  $V_x$  lowers the barrier, and the eigenmodes of Hamiltonian (1)become closer to delocalized wavepackets on the two nonadiabatic surfaces. In this case, one can observe significant oscillations between the donor and acceptor states before the steady state is reached (see, for example, Fig. 1(c)). This corresponds to a Type II or Type III mixed valence compound [37].

**Non-adiabatic regime** - In the non-adiabatic, lowtemperature regime, the transfer is dominated by the vibrational mode structure: both the unitary and dissipative dynamics are frozen unless the donor-acceptor energy difference nearly matches the vibrational energy at  $\Delta E = \ell \omega$ , with  $\ell$  being an integer greater than zero. This vibrational-assisted dynamics [17] results in wellresolved resonances (see Fig. 2). Deep in the non-



FIG. 2. Non-adiabatic transfer regime: (a) The transfer rate  $k_T$  in units of the relaxation rate  $\gamma$  as a function of the donor-acceptor energy gap  $\Delta E$  for  $(V_x, g, \gamma) = (0.056, 1.4, 0.06)\omega$ . The blue points result from an exponential fit of the measured  $P_D(t)$  dynamics, with the error bars being the standard error of the fit. The dark blue solid curve is obtained from the fit of the dynamics predicted by Eq. (2). The FGR prediction (dark red solid line) is calculated using Eq. (4). (b-c) The experimental (b) and numerical (c) density plots of the time-resolved dynamics of  $P_D(t)$  as a function of both  $\Delta E$  and number of vibrational oscillations  $\omega t/2\pi$ . The detuning from the tilt mode is set to  $\delta/(2\pi) = -5$  kHz, and the numerical results include a motional dephasing of  $\gamma_m = 0.001\omega$ . (d) The transfer rate  $k_T$  in units of the relaxation rate  $\gamma$  as a function of the time-resolved dynamics of  $P_D(t)$  as a function of the donor-acceptor energy gap  $\Delta E$  for  $(V_x, g, \gamma) = (0.046, 0.521, 0.025)\omega$ . (e-f) The experimental (e) and numerical (f) density plot of the time-resolved dynamics of  $P_D(t)$  as a function of both  $\Delta E$  and number of vibrational oscillations  $\omega t/2\pi$ , with  $\delta/(2\pi) = -10$  kHz. The numerical results include a motional dephasing  $\gamma_m = 0.0005\omega$  (see Methods).

adiabatic regime, when  $|V_x| \ll \lambda/4$ , the eigenstates of the Hamiltonian  $H_s$  in Eq. (1) are close to uncoupled donor and acceptor vibronic states represented in Fig. 1b, namely  $|D\rangle |n_-\rangle$  and  $|A\rangle |n_+\rangle$ , respectively. In this case, the  $V_x \sigma_x$  term can be treated as a perturbation to the Hamiltonian  $H_0 = H_s - V_x \sigma_x$ . As a result, the transfer undergoes resonant transitions between the uncoupled donor and acceptor vibronic modes, following the FGR [25, 28, 38]:

$$k_T = 2\pi |V_x|^2 \sum_{n_-, n_+} p_{n_-} \operatorname{FC}_{n_-, n_+} \delta(E_{D, n_-} - E_{A, n_+}), \quad (4)$$

where  $p_{n_{-}}$  is the initial phonon populations in the donor state, and  $\operatorname{FC}_{n_{-},n_{+}} = |\langle n_{-} | n_{+} \rangle|^{2}$  is the Franck-Condon factor, namely the overlap between the two displaced Fock wavefunctions. A larger displacement  $g/\omega$  along the reaction coordinate leads to more vibrational states with a non-negligible overlap, there-

fore increasing the number of observable transfer resonances. In this regime, the effect of the bath can be taken into account by replacing the delta functions in Eq. (4) with normalized Lorentzian distributions with full-width-half-max  $\gamma$ , namely  $\delta(E_{D,n-} - E_{A,n+}) \rightarrow$  $(\gamma/2\pi)/[(E_{D,n-} - E_{A,n+})^2 + (\gamma^2/4)].$ 

In Figs. 2(a), we show the transfer rates extracted from the dynamics of the donor population  $P_D(t)$ , shown in Fig. 2(b) (experimental data) and 2(c) (theory) as density plots as a function of  $\Delta E$  and the number of vibrational oscillations  $\omega t/2\pi$ . The transfer rates extracted from an exponential decay fit of  $P_D(t)$  agree with the numerical predictions from the Lindblad master equation in Eq. (2), exhibiting distinct peaks at  $\Delta E = \ell \omega$ . In Fig. 2(a-c), the chosen parameters place the system in the non-adiabatic regime ( $V_x = 0.056\omega$  and  $\lambda/4 = 0.49\omega$ ), which is confirmed by the qualitative agreement between the FGR prediction (solid dark red line), the experimen-



FIG. 3. Adiabatic transfer regime: (a) The transfer rate  $k_T$  measured with  $(V_x, g, \gamma) = (0.18, 0.95, 0.020)\omega$  (red circles) and  $(V_x, g, \gamma) = (0.21, 1.08, 0.038)\omega$  (blue circles). The solid curves are the transfer rates calculated from Eq. (2) using the definition in Eq. (5) and including spin decoherence ( $\gamma_z = 0.0025\omega$ ) and motional dephasing ( $\gamma_m = 0.0013\omega$ ). The transfer rates overlap when scaled in units of the relaxation rate  $\gamma$ . The error bars are calculated using bootstrapping (see Methods). (b) Experimental donor population evolution  $P_D(t)$  versus energy gap  $\Delta E$  and number of vibrational oscillations  $\omega t/2\pi$  with the same parameters as the red circles in (a). Here, the detuning from the tilt mode is set to  $\delta/2\pi = -4$  kHz. (c) Corresponding numerical results with the same parameters as (b).

tal results, and the exact theory. Here, since  $g = 1.4\omega$ , we can observe transfer resonances involving vibrational states up to n = 4 within our experimental resolution (see Methods). In Fig. 2(d-f), we decrease the spinmotion coupling to  $g = 0.521\omega$ , and the Frank-Condon coefficients  $FC_{n_-,n_+}$  also change, in turn affecting the transfer rate and reducing the number of observed resonances compared to Fig. 2(a-c).

In this case, the FGR in Eq.(4) underestimates both the experimental and numerical results, showing that the system is already in a non-perturbative regime.

Strongly adiabatic regime - When the electronic coupling  $V_x$  is comparable to the activation barrier  $\lambda/4$ and larger than the relaxation rate  $\gamma$ , the dynamics cannot be simply described in terms of weakly coupled wavefunction localized on the donor and acceptor site. In this regime, the population evolution features an initial coherent oscillation between the donor and acceptor states before the eventual equilibration in the acceptor state, as shown in Fig. 3(b-c). Here, the density plots of the experimental and theoretical  $P_D(t)$  are plotted as a function of  $\Delta E$ , showing good agreement. In this regime, the evolution cannot be fitted with an exponential function as in the non-adiabatic case. Therefore, to extract the effective transfer rate, we use the inverse lifetime of the donor population as proposed in Refs. [28, 38]:

$$k_T^{-1} = \frac{\int t P_D(t) dt}{\int P_D(t) dt}.$$
(5)

In Fig. 3(a) the transfer rates are extracted using Eq. (5) by interpolating and integrating both the experimental data and the numerical results (see Methods). We show the transfer rates extracted from the data for two sets of parameters that have nearly equal spin-phonon coupling g and electronic coupling  $V_x$ , but different relaxation rates  $\gamma$ . We report the results in units of  $\gamma$ , showing that the transfer rate is proportional to the relaxation rate  $(k_T \propto \gamma)$  [28]. A few comments are in order: (i) for  $\Delta E < 2\omega$ , the transfer rate  $k_T$  does not exhibit distinct resonances as opposed to the transfer rate in the non-adiabatic regime. (ii) for  $\Delta E > 2\omega$ , the characteristic peaked structure of the non-adiabatic regime is recovered, which can be explained by the localization of the initial state in the upper surface, as suggested by Ref. [28]; (iii) For  $\Delta E > 3\omega$ , the envelope of the transfer rate shows a decrease as a function of  $\Delta E$ . This is sometimes called the "inverted regime" of electron transfer, where, at both high and low temperatures, the reaction counterintuitively becomes slower despite the transfer becoming more exothermic. This can be explained by the decreasing Frank-Condon factor  $\mathrm{FC}_{n_-,n_+}$  as a function of  $\Delta E$ and can also be observed in the non-adiabatic regime (see Fig. 2).

**Optimal transfer** - When  $\Delta E$  is set on a resonance, sweeping  $V_x/\gamma$  allows one to pinpoint an optimal transfer regime [38]. In Fig. 4, we report the transfer rate measured as a function of  $V_x/\gamma$ , setting  $\Delta E = 2\omega$ . The data exhibit a distinct optimal transfer rate at  $V_x/\gamma \sim 3.3$ , in good agreement with the numerical predictions based on Eq. (2). It is worth noting that, for small  $V_x/\gamma$ , the transfer rate varies quadratically as predicted by Eq. (4). Beyond the optimum, the transfer rate is less sensitive to  $V_x/\gamma$ . This robustness has been suggested to be important for fast transfer in photosynthetic complexes [38, 39]. In particular, the presence of an optimal relaxation rate underscores the crucial role of dephasing in transport phenomena that was previously pointed out in solid-state [40] and atomic systems [22], as well as in biomolecules [41–44].

In conclusion, our experiment demonstrates the re-



FIG. 4. **Optimal transfer:** Transfer rate  $k_T$  as a function of  $V_x/\gamma$ , with  $(\Delta E, g, \gamma) = (2, 0.80, 0.11)\omega$  and detuning  $\delta/2\pi = -4$  kHz. The numerical results (solid curve) include spin decoherence ( $\gamma_z = 0.0013\omega$ ) and motional dephasing ( $\gamma_m = 0.0013\omega$ ). The optimal transfer is located at  $V_x/\gamma \sim 3.3$ , in agreement with the theoretical prediction of Eq. (2). Error bars are calculated using bootstrapping (see Methods).

markable flexibility of the trapped-ion platform to perform direct analog quantum simulations of models relevant to chemical physics, including an engineered environment. These simulations are performed through careful tuning of both the Hamiltonian of the trapped-ion system and its engineered reservoir by using seven simultaneous laser tones and two different atomic species. This toolbox allowed us to investigate relevant regimes of a paradigmatic ET model with tunable dissipation at low temperatures, where the interplay of quantum effects and interactions with the environment is crucial in determining the dynamics. The observed time-resolved dynamics of the donor-acceptor population and the measured transfer rate in both the non-adiabatic and adiabatic regimes agree with the numerics with independently calibrated parameters and identify an optimal transfer regime that has been suggested to be relevant for ET in photosynthetic complexes [38].

We note that, from a quantum optics perspective, our system simulates a variant of the Rabi model [29] with tunable dissipation, ranging from weak to ultra-strong coupling regimes. In this regard, the Rabi model with dissipation can be investigated by measuring motional observables in addition to the spin degrees of freedom, which is an inherent capability of the trapped-ion platform. In the Supplementary Information, we measure the average phonon population in the steady state of the evolution under Eq. (2) and observe spin-phonon correlations depending on the donor-acceptor energy separation.

Our setup can be extended to include multiple sites encoded in separate ions interacting via a spin-hopping Hamiltonian. This will enable the study of the dynamics of Frenkel-type excitons [45] to investigate the role of coherence and exciton delocalization in the energy transfer process in biomolecules and photosynthetic complexes [46–48]. In this setting, ancillary cooling ions can provide multiple bosonic modes to engineer structured spectral density functions [34] leading to the simulation of colored baths and non-Markovian dynamics [49, 50]. In addition, the trapped-ion platform naturally offers the possibility to include tunable anharmonic couplings among different bosonic modes [51] that can be used to study the effects of anharmonicity on energy transfer [52], a crucial but often overlooked feature of realistic molecular systems. Finally, increasing the number of ions and bosonic modes in this setup will also allow dissipative engineering of correlated many-body phonon and spin states [53–56]. Our experiment is a stepping-stone toward the use of quantum devices to provide new insights into open questions in chemical and biological physics and to shed light on the underlying principles of biochemical processes.

# ACKNOWLEDGMENTS

The authors acknowledge Yoshitaka Tanimura for insightful discussions and Fabrizio Minganti, Diego Fallas-Padilla, and Marcello Dalmonte for suggestions and careful reading of the manuscript. We acknowledge April Sheffield for her early contribution to the experimental setup. GP acknowledges the support of the Welch Foundation Award C-2154, the Office of Naval Research Young Investigator Program (grant no. N00014-22-1-2282), the NSF CAREER Award (grant No. PHY-2144910), the Army Research Office (W911NF22C0012), and the Office of Naval Research (grant No. N00014-23-1-2665). The authors acknowledge that this material is based upon work supported by the U.S Department of Energy, Office of Science, Office of Nuclear Physics under the Early Career Award No. DE-SC0023806. The isotopes used in this research were supplied by the U.S. Department of Energy Isotope Program, managed by the Office of Isotope R&D and Production. HP acknowledges support from NSF under Grant No. PHY-2207283. Work at the Center for Theoretical Biological Physics was supported by the NSF (Grant PHY-2019745). JNO was also supported by the NSF grant PHY-2210291. PGW was also supported by the D. R. Bullard-Welch Chair at Rice University, Grant No. C0016.

 A. J. Daley, I. Bloch, C. Kokail, S. Flannigan, N. Pearson, M. Troyer, and P. Zoller, Practical quantum advantage in quantum simulation, Nature 607, 667 (2022).

- [2] M. Reiher, N. Wiebe, K. M. Svore, D. Wecker, and M. Troyer, Elucidating reaction mechanisms on quantum computers, Proceedings of the National Academy of Sciences **114**, 7555 (2017), https://www.pnas.org/doi/pdf/10.1073/pnas.1619152114.
- [3] N. Maskara, S. Ostermann, J. Shee, M. Kalinowski, A. M. Gomez, R. A. Bravo, D. S. Wang, A. I. Krylov, N. Y. Yao, M. Head-Gordon, M. D. Lukin, and S. F. Yelin, Programmable simulations of molecules and materials with reconfigurable quantum processors (2023), arXiv:2312.02265 [quant-ph].
- [4] S. Mostame, J. Huh, C. Kreisbeck, A. J. Kerman, T. Fujita, A. Eisfeld, and A. Aspuru-Guzik, Emulation of complex open quantum systems using superconducting qubits, Quantum Information Processing 16, 44 (2016).
- [5] M. Kang, H. Nuomin, S. N. Chowdhury, J. L. Yuly, K. Sun, J. Whitlow, J. Valdiviezo, Z. Zhang, P. Zhang, D. N. Beratan, and K. R. Brown, Seeking a quantum advantage with trapped-ion quantum simulations of condensed-phase chemical dynamics, Nature Reviews Chemistry 8, 340 (2024).
- [6] H. Frauenfelder and P. G. Wolynes, Rate theories and puzzles of hemeprotein kinetics, Science 229, 337 (1985), https://www.science.org/doi/pdf/10.1126/science.4012322.
- [7] R. Zhuravel, H. Huang, G. Polycarpou, S. Polydorides, P. Motamarri, L. Katrivas, D. Rotem, J. Sperling, L. A. Zotti, A. B. Kotlyar, J. C. Cuevas, V. Gavini, S. S. Skourtis, and D. Porath, Backbone charge transport in doublestranded dna, Nature Nanotechnology 15, 836 (2020).
- [8] P. G. Wolynes, Imaginary time path integral Monte Carlo route to rate coefficients for nonadiabatic barrier crossing, The Journal of Chemical Physics 87, 6559 (1987), https://pubs.aip.org/aip/jcp/articlepdf/87/11/6559/18968099/6559\_1\_online.pdf.
- [9] C. Zheng, J. A. McCammon, and P. G. Wolynes, Quantum simulation of nuclear rearrangement in electron transfer reactions, Proceedings of the National Academy of Sciences 86, 6441 (1989), https://www.pnas.org/doi/pdf/10.1073/pnas.86.17.6441.
- [10] J. E. Lawrence and D. E. Manolopoulos, Path integral methods for reaction rates in complex systems, Faraday Discuss. 221, 9 (2020).
- [11] E. R. Bittner and P. J. Rossky, Quantum decoherence in mixed quantum-classical systems: Nonadiabatic processes, The Journal of Chemical Physics 103, 8130 (1995), https://pubs.aip.org/aip/jcp/articlepdf/103/18/8130/19079262/8130\_1\_online.pdf.
- [12] G. D. Scholes, G. R. Fleming, L. X. Chen, A. Aspuru-Guzik, A. Buchleitner, D. F. Coker, G. S. Engel, R. van Grondelle, A. Ishizaki, D. M. Jonas, J. S. Lundeen, J. K. McCusker, S. Mukamel, J. P. Ogilvie, A. Olaya-Castro, M. A. Ratner, F. C. Spano, K. B. Whaley, and X. Zhu, Using coherence to enhance function in chemical and bio-physical systems, Nature **543**, 647 (2017).
- [13] L. Wang, M. A. Allodi, and G. S. Engel, Quantum coherences reveal excited-state dynamics in biophysical systems, Nature Reviews Chemistry 3, 477 (2019).
- [14] Y. Tanimura, Numerically "exact" approach to open quantum dynamics: The hierarchical equations of motion (HEOM), The Journal of Chemical Physics 153, 020901 (2020), https://pubs.aip.org/aip/jcp/articlepdf/doi/10.1063/5.0011599/15575870/020901\_1\_online.pdf.
- [15] N. Makri, Modular path integral methodology for real-time quantum dynamics, The

Journal of Chemical Physics **149**, 214108 (2018), https://pubs.aip.org/aip/jcp/articlepdf/doi/10.1063/1.5058223/15551048/214108\_1\_online.pdf.

- [16] S. Kundu and N. Makri, Real-time path integral simulation of exciton-vibration dynamics in light-harvesting bacteriochlorophyll aggregates, The Journal of Physical Chemistry Letters 11, 8783 (2020).
- [17] D. J. Gorman, B. Hemmerling, E. Megidish, S. A. Moeller, P. Schindler, M. Sarovar, and H. Haeffner, Engineering vibrationally assisted energy transfer in a trapped-ion quantum simulator, Phys. Rev. X 8, 011038 (2018).
- [18] J. Whitlow, Z. Jia, Y. Wang, C. Fang, J. Kim, and K. R. Brown, Quantum simulation of conical intersections using trapped ions, Nature Chemistry 15, 1509 (2023).
- [19] C. H. Valahu, V. C. Olaya-Agudelo, R. J. MacDonell, T. Navickas, A. D. Rao, M. J. Millican, J. B. Pérez-Sánchez, J. Yuen-Zhou, M. J. Biercuk, C. Hempel, T. R. Tan, and I. Kassal, Direct observation of geometric-phase interference in dynamics around a conical intersection, Nature Chemistry 15, 1503 (2023).
- [20] C. S. Wang, N. E. Frattini, B. J. Chapman, S. Puri, S. M. Girvin, M. H. Devoret, and R. J. Schoelkopf, Observation of wave-packet branching through an engineered conical intersection, Phys. Rev. X 13, 011008 (2023).
- [21] A. Potočnik, A. Bargerbos, F. A. Y. N. Schröder, S. A. Khan, M. C. Collodo, S. Gasparinetti, Y. Salathé, C. Creatore, C. Eichler, H. E. Türeci, A. W. Chin, and A. Wallraff, Studying light-harvesting models with superconducting circuits, Nature Communications 9, 904 (2018).
- [22] C. Maier, T. Brydges, P. Jurcevic, N. Trautmann, C. Hempel, B. P. Lanyon, P. Hauke, R. Blatt, and C. F. Roos, Environment-assisted quantum transport in a 10qubit network, Phys. Rev. Lett. **122**, 050501 (2019).
- [23] K. Sun, C. Fang, M. Kang, Z. Zhang, P. Zhang, D. N. Beratan, K. R. Brown, and J. Kim, Quantum simulation of polarized light-induced electron transfer with a trapped-ion qutrit system, The Journal of Physical Chemistry Letters 14, 6071 (2023), pMID: 37358521, https://doi.org/10.1021/acs.jpclett.3c01166.
- [24] C. Sparrow, E. Martín-López, N. Maraviglia, A. Neville, C. Harrold, J. Carolan, Y. N. Joglekar, T. Hashimoto, N. Matsuda, J. L. O'Brien, D. P. Tew, and A. Laing, Simulating the vibrational quantum dynamics of molecules using photonics, Nature 557, 660 (2018).
- [25] A. J. Leggett, S. Chakravarty, A. T. Dorsey, M. P. A. Fisher, A. Garg, and W. Zwerger, Dynamics of the dissipative two-state system, Rev. Mod. Phys. 59, 1 (1987).
- [26] A. Garg, J. N. Onuchic, and V. Ambegaokar, Effect of friction on electron transfer in biomolecules, The Journal of Chemical Physics 83, 4491 (1985).
- [27] P. G. Wolynes, Dissipation, tunneling, and adiabaticity criteria for curve crossing problems in the condensed phase, The Journal of Chemical Physics 86, 1957 (1987), https://pubs.aip.org/aip/jcp/articlepdf/86/4/1957/18963608/1957\_1\_online.pdf.
- [28] F. Schlawin, M. Gessner, A. Buchleitner, T. Schätz, and S. S. Skourtis, Continuously parametrized quantum simulation of molecular electron-transfer reactions, PRX Quantum 2, 010314 (2021).
- [29] D. Lv, S. An, Z. Liu, J.-N. Zhang, J. S. Pedernales, L. Lamata, E. Solano, and K. Kim, Quantum simulation of the quantum rabi model in a trapped ion, Phys.

Rev. X 8, 021027 (2018).

- [30] R. Marcus and N. Sutin, Electron transfers in chemistry and biology, Biochimica et Biophysica Acta (BBA) - Reviews on Bioenergetics 811, 265 (1985).
- [31] T. D. Lee, F. E. Low, and D. Pines, The motion of slow electrons in a polar crystal, Phys. Rev. 90, 297 (1953).
- [32] G. Clos, D. Porras, U. Warring, and T. Schaetz, Timeresolved observation of thermalization in an isolated quantum system, Phys. Rev. Lett. 117, 170401 (2016).
- [33] G. X. Wang, Y. K. Wu, R. Yao, W. Q. Lian, Z. J. Cheng, Y. L. Xu, C. Zhang, Y. Jiang, Y. Z. Xu, B. X. Qi, P. Y. Hou, Z. C. Zhou, L. He, and L. M. Duan, Simulating the spin-boson model with a controllable reservoir in an ion trap (2024), arXiv:2402.07461 [quant-ph].
- [34] A. Lemmer, C. Cormick, D. Tamascelli, T. Schaetz, S. F. Huelga, and M. B. Plenio, A trapped-ion simulator for spin-boson models with structured environments, New Journal of Physics 20, 073002 (2018).
- [35] C. Schneider, D. Porras, and T. Schaetz, Experimental quantum simulations of many-body physics with trapped ions, Reports on Progress in Physics 75, 024401 (2012).
- [36] M. Cetina, L. Egan, C. Noel, M. Goldman, D. Biswas, A. Risinger, D. Zhu, and C. Monroe, Control of transverse motion for quantum gates on individually addressed atomic qubits, PRX Quantum 3, 010334 (2022).
- [37] K. D. Demadis, C. M. Hartshorn, and T. J. Meyer, The localized-to-delocalized transition in mixed-valence chemistry, Chemical Reviews 101, 2655 (2001).
- [38] S. S. Skourtis, A. J. R. Da Silva, W. Bialek, and J. N. Onuchic, New look at the primary charge separation in bacterial photosynthesis, The Journal of Physical Chemistry 96, 8034 (1992), https://doi.org/10.1021/j100199a038.
- [39] J. N. Onuchic and P. G. Wolynes, Classical and quantum pictures of reaction dynamics in condensed matter: resonances, dephasing, and all that, *The Journal of Physical Chemistry*, The Journal of Physical Chemistry **92**, 6495 (1988).
- [40] D. E. Logan and P. G. Wolynes, Dephasing and anderson localization in topologically disordered systems, Phys. Rev. B 36, 4135 (1987).
- [41] M. B. Plenio and S. F. Huelga, Dephasing-assisted transport: quantum networks and biomolecules, New Journal of Physics 10, 113019 (2008).
- [42] P. G. Wolynes, Some quantum weirdness in physiology, Proceedings of the National Academy of Sciences 106, 17247 (2009).
- [43] P. Rebentrost, M. Mohseni, I. Kassal, S. Lloyd, and A. Aspuru-Guzik, Environment-assisted quantum transport, New Journal of Physics 11, 033003 (2009).
- [44] A. W. Chin, A. Datta, F. Caruso, S. F. Huelga, and M. B. Plenio, Noise-assisted energy transfer in quantum networks and light-harvesting complexes, New Journal of Physics 12, 065002 (2010).
- [45] S. J. Jang and B. Mennucci, Delocalized excitons in natural light-harvesting complexes, Rev. Mod. Phys. 90, 035003 (2018).
- [46] A. Ishizaki and G. R. Fleming, Theoretical examination of quantum coherence in a photosynthetic system at physiological temperature, Proceedings of the National Academy of Sciences 106, 17255 (2009).
- [47] F. Levi, S. Mostarda, F. Rao, and F. Mintert, Quantum mechanics of excitation transport in photosynthetic complexes: a key issues review, Reports on Progress in

Physics 78, 082001 (2015).

- [48] A. Mattioni, F. Caycedo-Soler, S. F. Huelga, and M. B. Plenio, Design principles for long-range energy transfer at room temperature, Phys. Rev. X 11, 041003 (2021).
- [49] I. de Vega and D. Alonso, Dynamics of non-markovian open quantum systems, Rev. Mod. Phys. 89, 015001 (2017).
- [50] P. M. Harrington, E. J. Mueller, and K. W. Murch, Engineered dissipation for quantum information science, Nature Reviews Physics 4, 660 (2022).
- [51] S. Ding, G. Maslennikov, R. Hablützel, and D. Matsukevich, Cross-kerr nonlinearity for phonon counting, Phys. Rev. Lett. **119**, 193602 (2017).
- [52] C. Zhang, M. Gruebele, D. E. Logan, and P. G. Wolynes, Surface crossing and energy flow in manydimensional quantum systems, Proceedings of the National Academy of Sciences **120**, e2221690120 (2023), https://www.pnas.org/doi/pdf/10.1073/pnas.2221690120.
- [53] S. F. Huelga, A. Rivas, and M. B. Plenio, Nonmarkovianity-assisted steady state entanglement, Phys. Rev. Lett. 108, 160402 (2012).
- [54] A. González-Tudela and D. Porras, Mesoscopic entanglement induced by spontaneous emission in solid-state quantum optics, Phys. Rev. Lett. **110**, 080502 (2013).
- [55] D. Kienzler, H.-Y. Lo, B. Keitch, L. de Clercq, F. Leupold, F. Lindenfelser, M. Marinelli, V. Negnevitsky, and J. P. Home, Quantum harmonic oscillator state synthesis by reservoir engineering, Science 347, 53 (2015), http://science.sciencemag.org/content/347/6217/53.full.pdf.
- [56] G. Morigi, J. Eschner, C. Cormick, Y. Lin, D. Leibfried, and D. J. Wineland, Dissipative quantum control of a spin chain, Phys. Rev. Lett. **115**, 200502 (2015).
- [57] C. Tamm, N. Huntemann, B. Lipphardt, V. Gerginov, N. Nemitz, M. Kazda, S. Weyers, and E. Peik, Cs-based optical frequency measurement using cross-linked optical and microwave oscillators, Phys. Rev. A 89, 023820 (2014).
- [58] D. T. C. Allcock, W. C. Campbell, J. Chiaverini, I. L. Chuang, E. R. Hudson, I. D. Moore, A. Ransford, C. Roman, J. M. Sage, and D. J. Wineland, omg blueprint for trapped ion quantum computing with metastable states, Applied Physics Letters 119, 214002 (2021), https://pubs.aip.org/aip/apl/articlepdf/doi/10.1063/5.0069544/13267942/214002\_1\_online.pdf.
- [59] C. Monroe, W. C. Campbell, L.-M. Duan, Z.-X. Gong, A. V. Gorshkov, P. W. Hess, R. Islam, K. Kim, N. M. Linke, G. Pagano, P. Richerme, C. Senko, and N. Y. Yao, Programmable quantum simulations of spin systems with trapped ions, Rev. Mod. Phys. **93**, 025001 (2021).
- [60] J. Johansson, P. Nation, and F. Nori, Qutip 2: A python framework for the dynamics of open quantum systems, Computer Physics Communications 184, 1234 (2013).
- [61] C. Flühmann, T. L. Nguyen, M. Marinelli, V. Negnevitsky, K. Mehta, and J. P. Home, Encoding a qubit in a trapped-ion mechanical oscillator, Nature 566, 513 (2019).
- [62] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2007).
- [63] H. J. Carmichael, Statistical methods in quantum optics 1: master equations and Fokker-Planck equations (Springer Science & Business Media, 2013).
- [64] T. Brandes, Chapter 7 (quantum dissipation), lectures on background to quantum information (2004)

### METHODS

### A. Ion trap setup

The experiment is based on a blade trap, where each blade features 5 segmented electrodes. We mounted the gold-coated fused silica blades on an Alumina holder. Alumina is chosen for its high thermal conductivity and low degassing rate. The blades are positioned in a  $60^{\circ}/30^{\circ}$  angle configuration to enable high optical access along the vertical direction for high-resolution imaging (0.6 NA) and along the in-plane direction orthogonal to the trap axis (0.3 NA). This configuration also breaks rotational symmetry, which allows for well-defined trap principal axes. Each electrode is biased via a gold fuzz button, which is, in turn, connected to a Kaptoninsulated wire via customized Macor holders. To shunt the RF pickup voltages on the static DC blades, we use UHV-compatible silver-filled epoxy to glue 800 pF capacitors to each static segment on one side and wire-bond the other side to a ground strip present on the blades. We use a helical resonator with a resonant frequency of 27.9 MHz and a quality factor Q = 198 to drive the rf blades, achieving a trap center-of-mass radial trap frequency of 3.363 MHz at  $V_{\text{peak}} = 420$  V. The heating rate on the radial center-of-mass (com) mode is measured to be 0.4 quanta/ms, while the tilt mode features a lower heating rate ( $\dot{n} \sim 0.03$  quanta/ms).

# B. Laser setup

A 370 nm laser red detuned from the  ${}^{2}S_{1/2} \rightarrow {}^{2}P_{1/2}$ transition passing through 3.704 and 14.748 GHz EOMs is used to produce Doppler cooling light for both isotopes  ${}^{171}\text{Yb}^{+}$  and  ${}^{172}\text{Yb}^{+}$ . This beam is placed in-plane at  $45^{\circ}$  with respect to the ion chain for projection along all three trap principal axes. In addition, two axial 370 nm beams are used for detection and optical pumping of  ${}^{171}\text{Yb}^{+}$ . They are also superimposed with two 935 nm superimposed repumper beams for both Yb<sup>+</sup> isotopes.

A pulsed 355 nm laser is used to resonantly address the  $^{171}\mathrm{Yb^+}$  ground state qubit via two-photon Raman processes. The same laser is used to generate the spin-phonon coupling. The counter-propagating Raman beams have elliptical shapes with vertical and horizontal waists  $w_z=5\,\mu\mathrm{m}$  and  $w_x=150\,\mu\mathrm{m}$  and are in  $\mathrm{lin}\pm\mathrm{lin}$  polarization configuration to maximize the coupling between the two ground-state clock states.

A 435 nm diode laser locked to a Ultra-low-expansion cavity is used to address  ${}^{2}S_{1/2} \rightarrow {}^{2}D_{3/2}$  transition (or  $|g\rangle \rightarrow |o\rangle$ ) in  ${}^{172}$ Yb<sup>+</sup> [57, 58]. The beam is aligned at 45° with respect to the magnetic field and horizontally polarized to maximize the coupling to the two  $\Delta m_{j} = 0$  transitions ( $m_{J} = \pm 1/2 \rightarrow m_{J'} = \pm 1/2$ ) separated by 8.23 MHz. The cooling is achieved by continuously driving the red sideband (RSB) of  $m_{J} = \pm 1/2 \rightarrow m_{J'} = \pm 1/2$ transitions while also using a 935 nm repumper laser that allows the transition between  $|o\rangle$  and  ${}^{3}D[3/2]_{1/2} \equiv |e\rangle$ . Two tones on the 935 nm laser separated by 113 MHz address both  ${}^{171}\mathrm{Yb^{+}}$  and  ${}^{172}\mathrm{Yb^{+}}$ . To avoid optical pumping into either of  $m_{J} = \pm 1/2$  ground states during continuous sideband cooling, we use two laser tones on the 435 nm laser to address both the  $m_{J} = \pm 1/2 \rightarrow m_{J'} = \pm 1/2$  transitions simultaneously. The effective cooling rate is highly dependent on the power of the 935 nm laser, and it is the main turning knob to tune the cooling rate  $\gamma$ .

### C. Hamiltonian derivation

In this section, we derive the mapping from the electron transfer model in Eq. (1) to the experimental trapped-ion Hamiltonian. When we apply a pair of counter-propagating Raman beams with a wavevector difference of  $\vec{k}$ , phase difference of  $\phi$ , and a beatnote frequency at  $\omega_L$  on the <sup>171</sup>Yb<sup>+</sup> trapped qubit in a dual-species chain, the system can be described by ( $\hbar = 1$ ):

$$H = \frac{\omega_{\rm hf}}{2} \sigma_z + \sum_{\nu} \omega_{\nu} a_{\nu}^{\dagger} a_{\nu} + \frac{\Omega}{2} \left( e^{\sum_{\nu} i \eta_{\nu} \left( a_{\nu} + a_{\nu}^{\dagger} \right) - i \omega_L t + i \phi} \sigma^+ + \text{h.c.} \right), \quad (6)$$

where  $\omega_{\rm hf}$  is the energy splitting between the two qubit states,  $\omega_{\nu}$  is the  $\nu$ -th collective motional frequency of the chain associated with the raising (lowering) operator,  $a_{\nu}^{\dagger}(a_{\nu})$ ,  $\Omega$  is the Rabi coupling strength, and  $\eta_{\nu} = k\sqrt{1/2m\omega_{\nu}b_{\nu}}$  is the Lamb-Dicke parameter and m is the qubit mass.  $b_{\nu}$  is the normalized motional eigenvector for the <sup>171</sup>Yb<sup>+</sup> qubit ion in the  $\nu = 1, 2$  modes, namely the com and tilt modes of the <sup>171</sup>Yb<sup>+</sup>-<sup>172</sup>Yb<sup>+</sup> crystal.

By adding and subtracting  $\sum \delta_{\nu} a_{\nu}^{\dagger} a_{\nu}$  to Eq. (6) and rotating with respect to  $\frac{\omega_{\rm hf}}{2} \sigma_z + \sum_{\nu} \mu a_{\nu}^{\dagger} a_{\nu}$ , Eq. (6) is transformed into a resonant interaction frame rotating at  $\mu = \omega_L - \omega_{\rm hf} \equiv \omega_{\nu} + \delta_{\nu}$ , where  $\delta_{\nu}$  is the detuning from the  $\nu$ -th motional mode [35]. In our experiment,  $\mu + \omega_{\nu} \gg$  $|\mu - \omega_{\nu}| = \delta_{\nu}$ , therefore a rotating-wave approximation (RWA) is justified, and terms that rotate at  $\mu + \omega_{\nu}$  can be neglected. After the RWA, the Hamiltonian is described by:

$$H_I^{\text{res}} = \frac{\Omega}{2} \left( e^{i \sum_{\nu} \eta_{\nu} \left( a_{\nu} e^{-i\mu t} + a_{\nu}^{\dagger} e^{i\mu t} \right)} e^{i(\mu t + \phi)} \sigma^+ + \text{h.c.} \right) - \sum_{\nu} \delta_{\nu} a_{\nu}^{\dagger} a_{\nu}.$$
(7)

For our setup, the detuning from the tilt mode  $(\delta_{\text{tilt}}/2\pi \equiv \delta/2\pi \text{ in the main text})$  ranges from -4 to -10 kHz, while  $\delta_{\text{com}}/2\pi \sim -160$  kHz. Therefore, we can safely neglect the contribution from the com mode, obtaining a single-mode Hamiltonian:

$$H_I^{\text{res}} = \frac{\Omega}{2} \left( e^{i\eta \left( ae^{-i\mu t} + a^{\dagger} e^{i\mu t} \right)} e^{i(\mu t + \phi)} \sigma^+ + \text{h.c.} \right)$$
$$-\delta a^{\dagger} a.$$
(8)



FIG. 5. Laser configuration and relevant level structures: (a) The diagram illustrates the laser beam setup for the experiments. The black double-sided arrows indicate the directions of the linearly polarized lights. The 355 nm Raman beam pair is in  $\ln \perp \ln p$  obtained and configuration to ensure maximum coupling to the clock qubit and minimal state-dependent Stark shift. The 435 nm beam is horizontally polarized to the magnetic field to predominately drive the two  $\Delta m_j = 0$  transitions of  ${}^2S_{1/2}$ ,  $m_j = \pm 1/2 \rightarrow {}^2D_{3/2}$ ,  $m_j = \pm 1/2$ . (b) Simplified level structures of  ${}^{171}$ Yb<sup>+</sup> and  ${}^{172}$ Yb<sup>+</sup> used in the experimental protocol.

In the experiment, we apply 4 tones to one of the Raman beams generating four beatnotes at frequencies  $\omega_r = \omega_{\rm hf} - \mu$  with phase  $\phi_r$  (red sideband or RSB),  $\omega_b = \omega_{\rm hf} + \mu$  with phase  $\phi_b$  (blue sideband or BSB),  $\omega_x = \omega_{\rm hf}$  with phase  $\phi_x$ , and  $\omega_y = \omega_{\rm hf}$  with phase  $\phi_y$ . Thus, Eq. (8) becomes

$$H_{I}^{\text{res}} = \sum_{k=r,b,x,y} \frac{\Omega_{k}}{2} \left( e^{i\eta \left( ae^{-i\mu t} + a^{\dagger}e^{i\mu t} \right)} e^{-i(\omega_{k} - \omega_{\text{hf}})t + i\phi_{k}} \sigma^{+} \right.$$
  
+h.c.)  $- \delta a^{\dagger}a,$   
(9)

The first two terms generate the spin-phonon coupling term in Eq. (1). In the Lamb-Dicke regime, where  $\eta \sqrt{\langle (a+a^{\dagger})^2 \rangle} \ll 1$ , we can expand the two terms with respect to  $\eta$  to the first order and apply rotating-wave approximation to neglect off-resonant terms rotating at  $\mu$  and  $2\mu$ . For  $\Omega_r = \Omega_b \equiv \Omega$ , we obtain the effective spin-phonon Hamiltonian

$$H_{\rm sp} = \frac{\eta\Omega}{2} \left( a e^{i\phi_m} + a^{\dagger} e^{-i\phi_m} \right) \left( \cos\phi_s \sigma_x + \sin\phi_s \sigma_y \right), \tag{10}$$

where the motional phase  $\phi_m \equiv \frac{\phi_b - \phi_r}{2}$  and the spin phase  $\phi_s \equiv \frac{\pi}{2} + \frac{\phi_b + \phi_r}{2}$ . We choose  $\phi_r = \phi_b = 0$  for the experiment. Hence, the Hamiltonian is further simplified to

$$H_{\rm sp} = \frac{\eta\Omega}{2} \sigma_y \left( a + a^{\dagger} \right). \tag{11}$$

The two remaining terms follow the same form, differed by only the phase difference  $\phi_k$  with k = x, y and, in the Lamb-Dicke regime, generate the Hamiltonian:

$$H_k = \frac{\Omega_k}{2} \left( \cos \phi_k \sigma_x + \sin \phi_k \sigma_y \right). \tag{12}$$

By substituting Eq. (11) and Eq. (12) with  $\phi_x = 0$ and  $\phi_y = \pi/2$  into Eq. (9), we obtain

$$H_I^{\text{res}} = \frac{\Omega_y}{2}\sigma_y + \frac{\Omega_x}{2}\sigma_x + \frac{\eta\Omega}{2}\sigma_y(a^{\dagger} + a) - \delta a^{\dagger}a \quad (13)$$

As explained in the next section, to map the ET model in Eq. (1) to Eq. (13), we apply a rotation  $U_x(\pi/2) = \exp(-i\sigma_x\pi/4)$  to the qubit initialized in  $|\downarrow\rangle_z$  prior to the simulation (see Fig. 6). This rotates the qubit to  $|\uparrow\rangle_y$ . At the end of the evolution, we apply another rotation  $U_x(\pi/2)$  to perform the mapping  $|\uparrow\rangle_y \leftrightarrow |\uparrow\rangle_z$  and  $|\downarrow\rangle_y \leftrightarrow |\downarrow\rangle_z$ . Therefore, to realize Eq. (1), the parameter mappings are  $\Omega_y = \Delta E$ ,  $\Omega_x/2 = V_x$ ,  $\eta\Omega = g$ , and  $\delta = -\omega$ .

### D. Experimental sequence

The experimental procedure is summarized in Fig. 6. Our system consists of a  $^{171}$ Yb<sup>+</sup> ion acting as the qubit and a  ${}^{172}$ Yb<sup>+</sup> ion acting as the coolant. Initially, we use the standard Doppler cooling technique on both ions to prepare the temperature of the trapped dual-species chain near the Doppler limit. We then perform the Raman resolved-sideband cooling protocol on the radial center-of-mass and tilt modes, followed by an optical pumping pulse, to prepare the system in  $|\downarrow\rangle_z \langle\downarrow|_z \otimes \rho$ , where  $\rho = \sum_n e^{-n\omega/k_B T} |n\rangle \langle n|$  is the thermal phonon density matrix of the tilt mode and  $k_B T = \omega / \log(1 + \omega)$  $1/\bar{n}_0$ ) is the associated temperature. The initial tilt mode average phonon  $\bar{n}_0$  ranges between 0.1 and 0.3, which is similar to the bath temperature  $\bar{n}$ . To transform the system from the qubit basis  $\sigma_z$  to the  $\sigma_y$  basis, we apply a global rotation  $U_x(\pi/2) = \exp(-i\sigma_x\pi/4)$ . The state of the system becomes  $|D\rangle \langle D| \otimes \rho$ , where  $|D\rangle \equiv |\uparrow\rangle_{y}$  here.

To prepare the motional population from  $\rho$  to  $\rho_{-}$  defined in the main text, we use the optical dipole force from two Raman beatnotes,  $\omega_r = \omega_{\rm hf} - \mu$  and  $\omega_b = \omega_{\rm hf} + \mu$ , which have the same Rabi coupling strength of  $\Omega^{\rm displace} = \Omega/2$ , with  $\Omega$  defined in Eq. (11) and  $\phi_r = \phi_b = 0$ . We point out that this is the same drive that generates the spin-phonon term in Eq. (13) but with half the Rabi coupling strength.

In the ordinary interaction frame, where we rotate Eq. (6) with respect to  $\frac{\omega_{\rm hf}}{2}\sigma_z + \omega_{\nu}a^{\dagger}a$  and use rotating-wave approximations to neglect terms that rotate at  $2\omega_{\nu}$ , the subsequent derivation modifies Eq. (11) to

$$H_{\rm displace}^{\rm eff} = \frac{\eta \Omega^{\rm displace}}{2} \sigma_y \left( a e^{i\delta t} + a^{\dagger} e^{-i\delta t} \right), \qquad (14)$$

which causes a spin-dependent displacement of the motional state. Under this operation, the system evolves as  $U(t) = \mathcal{D}(\alpha(t)) |\uparrow\rangle_y \langle\uparrow|_y + \mathcal{D}(-\alpha(t)) |\downarrow\rangle_y \langle\downarrow|_y \equiv \mathcal{D}(\alpha(t)) |D\rangle \langle D| + \mathcal{D}(-\alpha(t)) |A\rangle \langle A|$ , where  $\mathcal{D}$  is the displacement operator in position-momentum phase space and  $\alpha(t) = \alpha_0(1 - e^{-i\delta t})$  with  $\alpha_0 = \eta \Omega^{\text{displace}}/2\delta = g/4\delta$ [59]. Hence, the applied pulse duration is  $t_{\text{displace}} = \pi/\delta$ to get the displacement of  $\alpha(t_{\text{displace}}) = g/2\delta = -g/2\omega$ onto  $\rho$ .

With the system being in the desired initial state  $|D\rangle \langle D| \otimes \rho_{-}$ , we address the <sup>171</sup>Yb<sup>+</sup> with the 4 Raman beatnotes discussed in the previous section to generate  $H_I^{\text{res}}$  in Eq. (13) that maps to the ET unitary model. Simultaneously, we apply the continuous resolved sideband cooling protocol on the <sup>172</sup>Yb<sup>+</sup> 's narrow linewidth optical transition to sympathetically cool the tilt mode of the system at the rate  $\gamma$  and effectively realize an engineered phonon dissipation. By varying the simulation time  $t_{\text{sim}}$ , we can measure the time-dependent evolution of the system. Prior to the measurement, we rotate the system back to the qubit basis with another global rotation  $U_x(\pi/2)$ .

To measure the average spin excitation, we use spindependent fluorescence, where only the spin in state  $|\uparrow\rangle_z$ , now representing  $|D\rangle$  after the  $\pi/2$  global rotation, scatters photons. We use an objective lens with a numerical aperture of 0.6 to collect the scattered photons into the photomultiplier tube. The average state discrimination fidelity between  $|D\rangle$  and  $|A\rangle$  is 99.5%.

Alternatively, we can measure the average phonon excitation  $\langle a^{\dagger}a \rangle$  by performing an optical pumping pulse to put the spin state of the system to  $|\downarrow\rangle_z$  and a resonant Raman blue sideband (BSB) transition drive,  $H^{\text{BSB}} =$  $i(\eta\Omega/2)(a\sigma^- - a^{\dagger}\sigma^+)$ , before the average spin excitation measurement. The phonon-number distribution that represents the diagonal elements of the final phonon density matrix of the system,  $\rho_m$ , can be extracted by fitting the spin evolution under the resonant Raman blue sideband transition drive with

$$P_{|\uparrow\rangle_z}(t) = \frac{1}{2} \sum_n p(n) \left[ 1 - e^{-\alpha_m t} \cos(\sqrt{n+1}\eta \Omega t) \right], \quad (15)$$



FIG. 6. **Experimental protocol:** The motional population of the system is initially cooled down by Doppler cooling and Raman pulsed sideband cooling sequences to  $\rho$ . The spin degree of freedom is prepared to  $|\downarrow\rangle_z$  in the qubit basis with optical pumping. Then a  $\pi/2$  pulse along x direction to rotate the spin state  $|\downarrow\rangle_z$  to  $|D\rangle \equiv |\uparrow\rangle_y$ . Following this, the phonon state is displaced with an optical dipole force from a spin-phonon coupling drive to prepare the system for the initial state of the ET theory,  $|D\rangle \langle D| \otimes \rho_{-}$ . After that, the experimental pulses are performed. Before the spin or phonon measurement, another  $\pi/2$  pulse along x axis projects the final spin state back into the qubit basis.

where p(n) denotes the phonon-number state population,  $\alpha_m$  is a parameter to capture the decoherence rate of the spin-phonon evolution, and t is the drive time [29]. Hence, we can compute  $\langle a^{\dagger}a \rangle = \text{Tr} (\rho_m a^{\dagger}a)$ .

# E. System calibration

We independently calibrate the parameters of the laser tones used to realize both the unitary and the dissipative terms in Eq. (2). The spin-phonon coupling and phonon terms,  $\frac{\eta\Omega}{2}\sigma_y(a^{\dagger} + a) - \delta a^{\dagger}a$ , are calibrated by adjusting the Rabi frequency  $\eta\Omega_n$  and detunings  $\delta_n$  of the red and blue sideband Raman laser beatnotes from the tilt mode sideband resonances for n = r, b.

We calibrate the spin-phonon coupling and the detuning by preparing the <sup>171</sup>Yb<sup>+</sup> qubit in the z basis and applying the Hamiltonian in Eq. (14) with  $\eta\Omega$  as the spin-phonon Rabi coupling strength. The hopping period corresponds to  $2\pi/|\delta|$ , which we use to estimate  $\delta$ . We then drive each tone on resonance to the tilt mode separately while setting the other tone off-resonant to estimate the effective  $\eta\Omega$  for the experiments (see Fig. 7(a,b)). In order to compensate for the ac-Stark shift due to the off-resonant excitation of the carrier transition by the red and blue sidebands, we use the following procedure: we first balance the Rabi coupling strengths of the red and blue sideband resonant drives ( $\delta_n = 0$ ) to the tilt mode separately. Then we turn on both tones simultaneously with the same detuning,  $\delta_r = \delta_b = \delta$ , from



FIG. 7. Hamiltonian and dissipation experimental calibration. (a) Spin dynamics from the red and blue sideband Raman laser beatnotes with a common detuning from the tilt mode,  $\delta/2\pi = -10$  kHz, and equal Rabi coupling strengths,  $\eta\Omega = 0.55|\delta|$ . The hopping period corresponds to  $2\pi/|\delta|$ . (b) The same spin-phonon drives with one tone on resonance and another detuned from the tilt mode. Red circles (experimental data) and solid curve (numerics) correspond to  $\{\delta_r/2\pi, \delta_b/2\pi\} = \{0, -10\}$  kHz, and the blue counterparts correspond to  $\{\delta_r/2\pi, \delta_b/2\pi\} = \{-10, 0\}$  kHz. (c) Spin dynamics undergoes a carrier drive along x in the  $\sigma_y$  basis. The Rabi coupling strength is set to  $\Omega_x/2\pi = 0.99$  kHz. Together with another tone of the same frequency beatnote and phase difference of  $\pi/2$ , we generate the spin operation terms in the electron transfer Hamiltonian. The numerical results represented by solid curves in (a)-(c) include spin decoherence ( $\gamma_z/2\pi = 10$  Hz) and motional dephasing ( $\gamma_m/2\pi = 5$  Hz). (d) The evolution of the average tilt mode phonon occupation number of the dual-species ion crystal via continuous resolved sideband cooling on  $^{172}$ Yb<sup>+</sup> with 435 nm and 935 nm beams. The exponential constant determines the cooling rate. Here,  $\gamma/2\pi = 0.23$  kHz, and the steady state average phonon occupation number is  $\bar{n} = 0.2$ .

the tilt mode resonances. Using a Ramsey sequence, we adjust the ratio of the powers and a common frequency shift of the two laser tones to compensate for the undesired ac-Stark shift up to 0.25 kHz accuracy. For the  $\Omega_x$  and  $\Omega_y$  Rabi frequencies in Eq. (13), we adjust the power of the two carrier transition tones independently (see Fig. 7(c)).

The frequency of the 435 nm red sideband resonance of the <sup>172</sup>Yb<sup>+</sup> optical transition is found by using a scheme similar to Quantum logic spectroscopy (QLS) via spin-state measurements of  $^{171}$ Yb<sup>+</sup>. The pulse sequence consists of a series of three  $\pi$ -pulses on both ions, namely  $BSB\pi(355nm) \rightarrow RSB\pi(435nm) \rightarrow BSB\pi(355)$ nm), while varying the frequency of the 435 nm  $RSB\pi$ pulse. The 355 nm light is kept on and out of resonance with the <sup>171</sup>Yb<sup>+</sup> qubit during 435 nm illumination to account for the differential stark shift on the 435 nm cooling transition. Another method to quantify the 435 nm cooling transition frequency is to replace the initial SBC pulses of the tilt mode with a finite amount of 435 nm CSBC pulse. We then scan the RSB frequency of both the Zeeman  $\Delta m_i = 0$  transitions of <sup>172</sup>Yb<sup>+</sup> parking at the  $2\pi$  time of tilt mode BSB evolution. By observing the contrast of the BSB population at  $2\pi$  time, we estimate the stark shifted frequencies of the 435 nm RSB pulses during sympathetic cooling.

To estimate the cooling rate with 435 nm and 935 nm beams on  $^{172}{\rm Yb^+}$ , we carry out the following procedure: we first perform Doppler cooling on the dual-species chain; then, we employ continuous sideband cooling on the tilt mode through  $^{172}{\rm Yb^+}$ , followed by Raman sideband cooling on the com mode through  $^{171}{\rm Yb^+}$ ; subsequently, we optically pump the spin state of  $^{171}{\rm Yb^+}$  to  $|\downarrow\rangle_z$  and perform a phonon distribution measurement on tilt mode via a resonant BSB drive to estimate the aver-



FIG. 8. Cooling rate versus 935 nm power: An optimal 935 nm power is observed with the measured RSB Rabi frequency of about 3.4 kHz for each 435 nm tone. The solid curve is the theoretical results using the steady state solution of the master equation of a simplified three-level system ( $|g\rangle$ ,  $|o\rangle$ , and  $|e\rangle$ ) with  $\gamma \approx \Gamma_{935}\rho_{ee}$ , where  $\Gamma_{935}$  is the scattering rate and  $\rho_{ee}$  is the steady state population of  $|e\rangle$ . Here, we use the 935 nm detuning from  $|e\rangle$ ,  $\Delta_{935} = 2\pi \times 5.5$  MHz for the theory.

age tilt mode phonon. By varying the cooling time and measuring the corresponding average phonon, we can obtain the cooling rate  $\gamma$  and the average phonon  $\bar{n}$  in Eq. (2) with an exponential fit, as shown in Fig. 7(d). The cooling rate can be adjusted by changing the 935 nm repumper power as it is non-monotonically dependent on the 935 nm power exhibiting an optimum, as shown in Fig. 8.

# F. Numerical simulations

We simulate Eq. (2) using a Python package based on QUTIP [60], which allows us to include experimental imperfections that induce different types of dephasing in our system. As the experiment is performed in the rotated basis  $(z \leftrightarrow y)$ , fluctuations in the laser intensity and in the detuning cause effective spin decoherence, while trap frequency fluctuations and the heating rate of tilt motional mode cause motional dephasing. Therefore, when comparing the numerics with the experimental results, we introduce two additional dissipative processes, which modify Eq. (2) to

$$\frac{\partial \rho}{\partial t} = -i[H_{\rm s},\rho] + \gamma(\bar{n}+1)\mathcal{L}_a[\rho] + \gamma\bar{n}\mathcal{L}_{a^{\dagger}}[\rho] + \sum_{k=z,m} \gamma_k \mathcal{L}_{c_k}[\rho],$$
(16)

where the jump operator  $c_z = \sigma_y$  and its corresponding rate  $\gamma_z$  account for spin decoherence while the jump operator  $c_m = aa^{\dagger} + a^{\dagger}a$  and its corresponding rate  $\gamma_m$  account for motional dephasing [61]. We determine these dephasing rates by comparing numerical results to experimental data, finding that  $\gamma_z/2\pi \sim (0-10)$  Hz, and  $\gamma_m/2\pi \sim 5$  Hz.

### G. Data analysis

In the non-adiabatic regime, the transfer dynamics can be well described by an exponential decay. Due to the finite bath temperature  $\bar{n} \sim 0.1 - 0.3$ , the spin population transfer is not complete from  $|D\rangle$  to  $|A\rangle$ . Therefore, the transfer rates are extracted from an exponential function with the rates and final populations as the fitting parameters. The uncertainties of the rates are the corresponding standard errors of the fits.

On the other hand, the spin evolutions in the adiabatic regime feature complex oscillatory decays that an analytical model cannot describe. As mentioned in the main text, we use the inverse lifetime of the donor population in Eq. (5) to determine the transfer rates [28, 38]. This definition considers  $t \to \infty$ . Therefore, there is a correction we need to consider when we use this formula for a finite experimental time. In the case of no electronic coupling,  $V_x = 0$ , the donor population does not evolve,  $P_D(t) = 1$ , because it is in an eigenstate of the system. However, Eq. 5 still evaluates a non-zero transfer rate between t = 0 and  $t = t_{sim}$  as  $k_0 = \frac{2}{t_{sim}}$ . This contribution to the transfer rate only goes to zero if one evaluates Eq. 5 for  $t \to \infty$ . Since  $P_D$  reaches the steady state within our experimental resolution in a finite time  $t_{sim}$  ranging from 4 ms to 10 ms, we calculate the transfer rates by subtracting  $k_0$  as:

$$k_T = \frac{\int_0^{t_{\rm sim}} P_D(t)dt}{\int_0^{t_{\rm sim}} t P_D(t)dt} - k_0.$$
(17)

To numerically evaluate the integrals, we interpolate the

evolution  $P_D(t)$  data. We also use Eq. (17) to estimate the numerical transfer rates.

To estimate the errors of the transfer rate, we follow a resampling procedure. We consider the experimental error of each time-step of the  $P_D(t)$  measurements as the standard deviation of a normal distribution centered at the mean measured value. We then randomly sample the distributions at each time-step, and we estimate the error of the transfer rate by taking the standard deviation of the rates obtained from the resampled datasets by using Eq. (17). The process is repeated for all adiabatic transfer dynamics.

### SUPPLEMENTARY INFORMATION

# H. Phonon Steady State Characterization

In this section, we discuss the steady state of the Lindbladian master equation [62], focusing in particular on the properties of the phonon population. From Eq. (2), the expectation an observable O satisfies:

$$\partial_t \langle O \rangle = \operatorname{Tr} \left[ -iO\left[H,\rho\right] + O\gamma\left(\bar{n}+1\right) \mathcal{D}_a\left(\rho\right) + O\gamma\bar{n}\mathcal{D}_{a^{\dagger}}\left(\rho\right) \right] \\ = -i\langle [O,H] \rangle + \gamma\left(\bar{n}+1\right) \langle a^{\dagger}Oa - \frac{1}{2} \left\{ O, a^{\dagger}a \right\} \rangle \\ + \gamma \bar{n} \langle aOa^{\dagger} - \frac{1}{2} \left\{ O, aa^{\dagger} \right\} \rangle.$$
(18)

Using the bosonic commutation relation, the number operator  $n = a^{\dagger}a$  satisfies:

$$\partial_t \langle n \rangle = -i \frac{g}{2} \langle \sigma_z \left( a^{\dagger} - a \right) \rangle + \gamma \left( \bar{n} - \langle n \rangle \right).$$
 (19)

The creation/annihilation operators satisfy:

$$\partial_t \langle a^{\dagger} \rangle = i \frac{g}{2} \langle \sigma_z \rangle + (i\omega - \gamma/2) \langle a^{\dagger} \rangle, \partial_t \langle a \rangle = -i \frac{g}{2} \langle \sigma_z \rangle - (i\omega + \gamma/2) \langle a \rangle.$$
(20)

To obtain steady-state solutions we set LHS of Eq. (19), (20) equal to zero leading to:

$$n_{\rm ss} = \bar{n} - \frac{i}{2} \frac{g}{\gamma} \langle \sigma_z \left( a^{\dagger} - a \right) \rangle_{\rm ss}, \qquad (21)$$

$$\langle a^{\dagger} \rangle_{\rm ss} = -\frac{ig}{2i\omega - \gamma} \langle \sigma_z \rangle_{\rm ss}, \ \langle a \rangle_{\rm ss} = -\frac{ig}{2i\omega + \gamma} \langle \sigma_z \rangle_{\rm ss},$$
(22)

which immediately gives a expression of steady-state reaction coordinate y in terms of steady-state donor population  $P_D$ :

$$y_{\rm ss} = \frac{y_0}{2} \left( a + a^{\dagger} \right) = -\frac{2\omega g}{4\omega^2 + \gamma^2} (2P_D^{ss} - 1)y_0.$$
(23)

To quantify spin-phonon correlation we can compare the exact steady-state  $n_{\rm ss}$  in Eq. (21), with the one calculated assuming that spin and phonon are in an uncorrelated state, namely  $\langle \sigma_z (a^{\dagger} - a) \rangle_{\rm ss} = \langle \sigma_z \rangle_{\rm ss} \langle a^{\dagger} - a \rangle_{\rm ss}$ .



FIG. 9. Steady-state characterization: Phonon population in the steady state (blue circles) as a function of  $\Delta E$ using  $(V_x, g, \gamma) = (0.19, 1.91, 0.038)\omega$ ,  $\delta/2\pi = -4$  kHz and  $\bar{n} = 0.2$ . The average phonon population is extracted by fitting the first six phonon states. The error bars are the standard deviation from the mean. The dark red solid line is the exact prediction given by Eq. (21) while the dark green solid line is the prediction given by Eq. (24). At low  $\Delta E$ , there is spin-phonon correlation in the steady state, which decreases monotonically as  $\Delta E$  is increased. Here, we also consider a motional dephasing of  $\gamma_m = 0.0013\omega$ .

This leads to:

$$n_{\rm ss}^{\rm un} = \bar{n} + \frac{g^2}{4\omega + \gamma^2} \left(2P_D^{ss} - 1\right)^2.$$
 (24)

In Fig. 9, we report measurements of the steady state  $\langle a^{\dagger}a \rangle_{ss}$  as a function of  $\Delta E$ . In order to measure the steady-state phonon population, after the evolution has reached its steady state, the sympathetic cooling is turned off, the qubit is reset via optical pumping, and then a resonant BSB Hamiltonian  $H^{BSB} = i(\eta\Omega/2)(a\sigma^{-}-a^{\dagger}\sigma^{+})$  is applied to the system. The resulting spin evolution is fitted to extract the average phonon population  $\langle n \rangle_{ss} = \langle a^{\dagger}a \rangle_{ss}$  in the steady state. The measured values are in agreement with the exact solution  $\langle n \rangle_{ss}$  in Eq. (21), confirming the presence of spin-phonon correlations in the system's steady state.

To get an intuitive understanding of the effect of dissipation due to the Lindbladian, we shall assume the spin is either in  $|\uparrow\rangle$  or  $|\downarrow\rangle$  such that the Hamiltonian (1) can be reduced to:

$$H_p = \omega a^{\dagger} a \pm \frac{g}{2} (a + a^{\dagger}). \tag{25}$$

The corresponding master equation becomes

$$\partial_t \rho = -i[H_p, \rho] + \gamma(\bar{n}+1)\mathcal{D}_a[\rho] + \gamma \bar{n}\mathcal{D}_{a^{\dagger}}[\rho].$$
(26)

We can define displaced bosonic creation/annihilation operators  $b \equiv a + \alpha$ ,  $b^{\dagger} \equiv a^{\dagger} + \alpha^{*}$  with  $\alpha$  being a complex constant to be determined. It can be shown that if we set

$$\alpha = \pm \frac{2g\omega}{4\omega^2 + \gamma^2} \pm \frac{ig\gamma}{4\omega^2 + \gamma^2},\tag{27}$$

Eq. (26) then becomes:

$$\partial_t \rho = -i[\omega b^{\dagger} b, \rho] + \gamma(\bar{n} + 1)\mathcal{D}_b[\rho] + \gamma \bar{n}\mathcal{D}_{b^{\dagger}}[\rho], \quad (28)$$

which takes the form of a simple damped oscillator with steady state being a thermal vibrational state characterized by  $\bar{n}$ . Undoing the displacement gives us the steadystate solution of Eq.(26):

$$\rho_{ss} = D(-\alpha)\rho_{th}D(\alpha), \ \rho_{th} = \frac{e^{-\beta\omega a^{\dagger}a}}{1 - e^{-\beta\omega}}, \qquad (29)$$

where D is the displacement operator and  $1/\beta = \omega/\log(1+1/\bar{n})$ . In the experiment  $\omega \gg \gamma$  such that  $\alpha \to \pm g/2\omega$ . Depending on the spin, the system is effectively pumped to the ground state of the left/right displaced harmonic well, as shown in Fig. 1.

### I. Ohmic Bath and Lindbladian Formalism

In this section, we show that the derivation of a Lindbladian master equation for the system in Eq. (1) in contact with an Ohmic bath is equivalent to the dissipative spin-boson model realized in this experiment under certain conditions. More formally, we will establish the equivalence between Eq. (2) and the spin-boson Hamiltonian  $H_{ET}$  in Eq. (1.3) of Ref. [26], namely:

$$H_{ET} = H_{\rm s} + H_{\rm sb} + H_{\rm b}.$$
 (30)

The system Hamiltonian  $H_{\rm s}$  is given by Eq. (1). The bath Hamiltonian  $H_{\rm b} = \sum_n \omega_n \Gamma_n^{\dagger} \Gamma_n$  is described by a collection of infinite harmonic oscillators with  $\Gamma_n(\Gamma_n^{\dagger})$ being the annihilation(creation) operator of the *n*-th bosonic mode. The reaction coordinate of the system is linearly coupled to the position coordinate of the bath via

$$H_{\rm sb} = S \otimes B, \ S \equiv a + a^{\dagger}, \ B \equiv K + K^{\dagger},$$
 (31)

with  $K^{\dagger} \equiv \sum_{n} c_{n} \Gamma_{n}^{\dagger}$  being a linear combination of bath operators and  $c_{n}$  being the coupling coefficients of the *n*-th mode. The coupling coefficients  $c_{n}$  and the bath frequencies determine the bath spectral density function  $J(\omega) = \sum_{n} c_{n}^{2} \delta(\omega - \omega_{n})$ . We take  $J(\omega)$  to be Ohmic [26]:

$$J(\omega) = \eta \omega \exp(-\omega/\omega_c), \ \omega_c \to \infty, \tag{32}$$

which corresponds to a classical damped oscillator with  $\eta$  being the linear damping coefficient. Note that in this section, we use  $\omega$  as the frequency variable for the spectral density function J and  $\omega_0$  as the bosonic mode frequency in Eq. (1).

To obtain the reduced dynamics of the system density matrix  $\rho(t)$ , we shall first change into the interaction picture of  $H_0 = H_s + H_b$ . Let us denote a generic operator O in the interaction picture of  $H_0$  as

$$\tilde{O}(t) = U^{\dagger}(t) OU(t), \ U(t) = \exp(-iH_0 t),$$
 (33)

the master equation in the interaction picture is then given by:

$$\partial_t \widetilde{\chi}(t) = -i[\widetilde{H}_{\rm sb}(t), \widetilde{\chi}(t)], \qquad (34)$$

where  $\chi$  is the system-bath density matrix. Explicitly integrating this equation and inserting the expression for  $\tilde{\chi}$  back leads to

$$\partial_t \widetilde{\chi} (t) = -i [\widetilde{H}_{\rm sb} (t), \chi (0)] - \int_0^t dt' [\widetilde{H}_{\rm sb} (t), [\widetilde{H}_{\rm sb} (t'), \chi (t')]]$$
(35)

Assuming the system-bath coupling is sufficiently weak, and the bath is kept at thermal equilibrium,  $\chi(t)$  becomes separable (Born approximation):

$$\chi(t) = \rho(t) \otimes R_0, \quad R_0 = e^{-\beta H_{\rm b}} / \text{Tr}(e^{-\beta H_{\rm b}}), \qquad (36)$$

where  $\beta = 1/k_B T$ . Taking partial trace with respect to the bath degrees of freedom on Eq. (35) gives an equation for  $\tilde{\rho}$ :

$$\partial_{t}\widetilde{\rho}(t) = -\int_{0}^{t} dt' C(\tau) \left[ \widetilde{S}(t) \widetilde{S}(t') \widetilde{\rho}(t') - \widetilde{S}(t') \widetilde{\rho}(t') \widetilde{S}(t) \right] + C(-\tau) \left[ \widetilde{\rho}(t') \widetilde{S}(t') \widetilde{S}(t) - \widetilde{S}(t) \widetilde{\rho}(t') \widetilde{S}(t') \right], \quad (37)$$

where  $\tau \equiv t - t'$  and  $C(\tau)$  is the temporal correlation function of the bath, namely

$$C(\tau) = \operatorname{Tr}\left[\widetilde{B}(\tau) BR_{0}\right]$$
  
= 
$$\int_{0}^{\infty} d\omega J(\omega) \left[\operatorname{coth}\left(\beta\omega/2\right)\cos\left(\omega\tau\right) - i\sin\left(\omega\tau\right)\right].$$
(38)

 $S, \tilde{S}, B, \tilde{B}$  follow the definitions in Eq. (31),(33). By introducing a displaced bosonic operator  $b = a + a_0$  with spin dependent constant  $a_0 = \frac{g}{2\omega_0}\sigma_z$ , we can evaluate the interaction frame system operator  $\tilde{S}(t)$ :

$$\widetilde{S}(t) = \left(a^{\dagger}e^{i\omega_0 t} + ae^{-i\omega_0 t}\right) + 2a_0\left(\cos\omega_0 t - 1\right).$$
(39)

Assuming the bath correlation function is strongly peaked around  $\tau = 0$  with a correlation time  $\tau_r$  much smaller than the typical time scale of the system's dynamics,  $C(\tau)$  can be approximated as  $\delta(\tau)$ , which yields the replacement  $\rho(t') \rightarrow \rho(t)$ . Another important effect of this approximation is the extension of the integration limit from t to  $\infty$  of the integral in Eq. (37) (Markovian approximation)[63]. Eq. (37) then results in the Schrodinger picture master equation:

$$\partial_{t}\rho\left(t\right) = -i\left[H_{s},\rho\left(t\right)\right] - \int_{0}^{\infty} d\tau \left[C\left(\tau\right)\left[S\widetilde{S}\left(-\tau\right)\rho\left(t\right) - \widetilde{S}\left(-\tau\right)\rho\left(t\right)S\right] + C\left(-\tau\right)\left[\rho\left(t\right)\widetilde{S}\left(-\tau\right)S - S\rho\left(t\right)\widetilde{S}\left(-\tau\right)\right]\right]$$

$$\tag{40}$$

Inserting Eq. (38),(39), this equation can be written in a compact form as:

$$\partial_{t}\rho(t) = -i[H_{s}, \rho(t)]$$

$$- [S(\Lambda + C_{0})\rho(t) - (\Lambda + C_{0})\rho(t)S + h.c.],$$
(41)

where  $\Lambda = \mathcal{L}_{+}a + \mathcal{L}a^{\dagger}$  and  $\mathcal{L}_{+}, \mathcal{L}$  take form of Laplace transform:

$$\mathcal{L}_{+} = \int_{0}^{\infty} d\omega J(\omega) \left[1 + \bar{n}(\omega)\right] \int_{0}^{\infty} d\tau e^{-i(\omega - \omega_{0})\tau}$$
$$\mathcal{L} = \int_{0}^{\infty} d\omega J(\omega) \bar{n}(\omega) \int_{0}^{\infty} d\tau e^{i(\omega - \omega_{0})\tau}, \qquad (42)$$

where  $\bar{n}(\omega)$  is the thermal phonon population of a mode of frequency  $\omega$  and  $C_0$  is a constant due to the scalar part of Eq. (39),

$$C_0 = 2a_0 \int_0^\infty d\tau C\left(\tau\right) \left(\cos\omega_0 \tau - 1\right). \tag{43}$$

It shall be noted that in the evaluation of  $\Lambda$ , we have applied the secular approximation, neglecting the contribution from highly oscillatory terms involving  $e^{\pm i(\omega+\omega_0)t}$ [64].

Evaluating the integrals in Eq. (42), (43) leads to the reduced master equation of the system:

$$\partial_{t}\rho = -i \left[H_{s} + H_{c}, \rho\right] - i \left[H_{n}\rho - \rho H_{n}^{\dagger}\right] + \gamma \left(\bar{n}_{0} + 1\right) \left(\mathcal{D}_{a}\left(\rho\right) + \mathcal{D}_{a}^{\prime}\left(\rho\right)\right) + \gamma \bar{n}_{0} \left(\mathcal{D}_{a^{\dagger}}\left(\rho\right) + \mathcal{D}_{a^{\dagger}}^{\prime}\left(\rho\right)\right) + i\Delta_{d}\mathcal{D}_{a}^{Im}\left(\rho\right).$$
(44)

Let us break down the different terms of the master equation (44):  $\mathcal{D}_c(\rho)$  is the Lindbladian super operator defined in (3) with coefficients,

$$\gamma = 2\pi\eta\omega_0, \ \bar{n}_0 = \bar{n}(\omega_0). \tag{45}$$

 $\mathcal{D}'_c(\rho)$  is defined as

$$\mathcal{D}_{c}^{\prime}(\rho) \equiv \frac{1}{2} \left( c^{\dagger} \rho c^{\dagger} + c \rho c - c^{\dagger} c^{\dagger} \rho - \rho c c \right), \qquad (46)$$

 $\mathcal{D}_{a}^{Im}\left(\rho\right)$  represents a super operator with imaginary coefficients

$$\mathcal{D}_{c}^{Im}\left(\rho\right) \equiv \left(c\rho c - c^{\dagger}\rho c^{\dagger}\right),\tag{47}$$

 $H_n$  represents the following non-hermitian Hamiltonian:

$$H_n = \Delta_d aa, \ \Delta_d = P\left[\int_0^\infty d\omega \frac{(2\bar{n}(\omega) + 1)J(\omega)}{\omega_0 - \omega}\right], \ (48)$$

where P stands for Principal Value, and  $H_c$  is a correc-

tion to the system Hamiltonian:

$$H_{c} = \widetilde{\omega}a^{\dagger}a + \frac{\widetilde{g}}{2}\sigma_{z}\left(a + a^{\dagger}\right),$$
  

$$\widetilde{\omega} = P\left[\int_{0}^{\infty}d\omega\frac{J\left(\omega\right)}{\omega_{0} - \omega}\right],$$
  

$$\widetilde{g} = 4gP\left[\int_{0}^{\infty}d\omega\frac{J\left(\omega\right)}{\omega_{0}^{2} - \omega^{2}}\right]$$
(49)

When the frequency of the system's oscillator  $\omega_0$  is much larger than the decay rate  $\gamma$ , under rotating wave approximation, we can effectively neglect the terms involving  $aa, a^{\dagger}a^{\dagger}$  that do not conserve the energy[63]. These include the superoperators  $\mathcal{D}'_a(\rho), \mathcal{D}'_{a^{\dagger}}(\rho), \mathcal{D}^{Im}_a(\rho)$  and the non-hermitian Hamiltonian terms  $H_n$ . The rotating wave approximation is consistent with the Born approximation, which assumes that the system-bath coupling is sufficiently small so that the system and bath can be described by a separable state. Finally, we obtain Eq. (2) in the main text, with renormalized oscillator frequency  $\omega' = \omega_0 + \tilde{\omega}$  and displacement  $g' = g + \tilde{g}$ .

We can check the validity of the Markovian approximation by comparing the time scale of the system dynamics  $\tau_{\rm s} \sim 1/\gamma$  and the width of the position correlation function estimated by the bath correlation time  $\tau_r \sim \beta$  [63]. For the approximation to be valid, we therefore require  $\tau_r \ll \tau_{\rm s}$ . Hence, for Eq. (2) to be a good description of the ET model in the weak decay regime, we require the following conditions, which are also derived in Ref. [34]:

$$\gamma \ll 1/\beta,$$
 (Markovian)  
 $\gamma \ll \omega_0.$  (RWA, Born)